



Oxidative stress and an animal neurotransmitter synthesizing enzyme in the leaves of wild growing myrtle after exposure to GSM radiation

Aikaterina L. Stefi^a, Dido Vassilacopoulou^b, Lukas H. Margaritis^c, Nikolaos S. Christodoulakis^{a,*}

^a Section of Botany, Faculty of Biology, National and Kapodistrian University of Athens, Ilisia, Athens, 15701, Hellas, Greece

^b Section of Biochemistry and Molecular Biology, Faculty of Biology, National and Kapodistrian University of Athens, Ilisia, Athens, 15701, Hellas, Greece

^c Section of Cell Biology & Biophysics, Faculty of Biology, National and Kapodistrian University of Athens, Ilisia, Athens, 15701, Hellas, Greece

ARTICLE INFO

Edited by Alessio Papini

Keywords:

Radiation

Myrtus communis

Leaf anatomy

Secondary metabolites

Environmental stress

ROS

DDC

ABSTRACT

Mobile phones turned to be the most common form of public communication. Therefore, life on our planet continues wrapped in a “cloud” of non-ionizing radiations. *Myrtus communis* L. is an evergreen shrub, common in Mediterranean formations, exposed and adapted to two seasonally separated and qualitatively different environmental stresses. Thus, we considered it important to investigate the response of this tolerant species to emitted GSM non-ionizing radiations and compare it to already available data from thoroughly investigated plant species. Although the leaves of the exposed plants present unaffected tissue arrangement, their mesophyll cells accumulate large amounts of secondary metabolites, their photosynthetic pigments are dramatically reduced, the ROS counted are significantly increased and the presence of DDC, which cannot be detected in the leaves of the control plants, is recorded in high levels. The exposed leaves seem to experience a severe oxidative stress which probably induces DDC expression and the biosynthesis of the neurotransmitter dopamine, the activation of the shikimate pathway and, eventually, the accumulation of secondary metabolites.

1. Introduction

Considering that Earth's natural radiofrequency environment has remained more or less unaltered within the lifespan of modern trees since before 1800, the major components of this environment were the radio emissions from space (galactic noise), the electric discharges within the atmosphere (atmospheric noise), and the small Radio Frequency (RF) component from the sun (Haggerty, 2010). Thus, we may assume that plants have evolved learning to use these environmental signals and intercept electromagnetic radiation (light) in order to regulate their circadian and seasonal functions (Haggerty, 2010). This is probably the main reason for which the impact of radio-frequency radiation on plants has received little attention and the physiological effects have been, rather arbitrarily, considered as negligible (Waldmann-Selsam et al., 2016).

High frequency electromagnetic fields (HF-EMF, i.e., frequencies from 300 MHz to 3 GHz, wavelengths from 1 m to 10 cm) are mainly human-produced, non-ionizing electromagnetic radiations that do not naturally occur in the environment (Vian et al., 2016). They are increasingly present because of the rapid development of wireless technology, such as Wi-Fi, various kinds of connected devices and cellular phones. Concerning the later, as the twentieth century passed away, the

number of the mobile phone users increased from about 2.2 to 5.9 billion between 2005 and 2011 (Choi, 2012). Approximately four mobile phone service providers exist in each given geographical area (Hyland, 2005). Consequently, the number of base stations was also increased to support the tremendous growth of mobile phone users (World Health Organization, 2006) thus high frequency non-ionizing electromagnetic fields (HF-EMF) are increasingly present in the environment and the living organisms of the civilised world thrive within a “cloud” of non-ionizing radiations.

Much concern is given to the effects of this radiation to human life and environmental health (Roux et al., 2006; Pietruszewski et al., 2007; Sheridan et al., 2010; Sharma and Parihar, 2014) although radio-frequency constitutes a genuine environmental stimulus for plants as well. This stimulus is able to evoke specific responses many of them being similar to those observed after a stressful treatment (Waldmann-Selsam et al., 2016). Moreover, plants proved an outstanding model for studying such interactions since their architecture (high surface/volume ratio) optimizes their interaction with the environment. They remain exposed to radiation at a constant orientation in the electromagnetic field, due to their inability to move (Vian et al., 2016), while it is possible to easily document changes over time, such as disturbed cell structure, growth and yield (Stefi et al., 2017a, 2017b), dying

* Corresponding author.

E-mail address: nchristo@biol.uoa.gr (N.S. Christodoulakis).

branches and colour changes of leaves. These damages can be considered objective and cannot be attributed to psychological or psychosomatic factors (Waldmann-Selsam et al., 2016).

During the previous century and currently, several researchers have investigated the effects of electromagnetic radiation on plants, both in the laboratory (Harte, 1950, 1972; Jerman et al., 1998; Lerchl et al., 2000; Sandu et al., 2005; Roux et al., 2006; Pietruszewski et al., 2007; Tkalec et al., 2005, 2009; Beaubois et al., 2007; Haggerty, 2010; Pesnya and Romanovsky, 2013; Sharma and Parihar 2014; Cammaerts and Johansson, 2015; Kumar et al., 2015; Stefi et al., 2016, 2017a, 2017b), and on the field (Bernatzky, 1986; Volkrodt, 1987, 1991; Selga and Selga, 1996; Balodis et al., 1996; Haggerty, 2010). For both types of studies, most of the times, detrimental effects have been reported, even for weak radiation exposure (Halgamuge, 2017).

It has long ago been documented that mediterranean plants – both the evergreen sclerophylls (Mitrakos and Christodoulakis, 1981) and the seasonally dimorphics (Christodoulakis, 1989) – are exposed to two seasonally separated and qualitatively different environmental stresses (Mitrakos, 1980). In the summer, the exclusionary high temperatures and the arid environment, force plant metabolism to shift from the photosynthetic to other, secondary metabolic pathways to relieve the cell from the oxidative stress caused by the ionized P680 pigments. During winter, frost is the other stressing condition which seems to be treated in a different way by the evergreen sclerophylls compared to the seasonally dimorphics.

In spite of the rather hostile climate, the Mediterranean evergreen sclerophyllous plants are considered to be very productive (Mooney et al., 1970; Di Castri and Mooney, 1973; Mooney and Dunn, 1976). This is true only when the environmental conditions are mild. Yet, the Mediterranean climate is not so generous (Mitrakos, 1980; Mitrakos and Christodoulakis, 1981) and the periods of favorable conditions, when growth can be promoted, are limited and unpredictable (Margaris, 1981). Eventually, Mediterranean plant formations fall short in biomass production. This is mainly due to the stressing conditions of the long and arid Mediterranean summer. Among them is the excess of solar radiation during the summer months. Excessive solar radiation causes photodamage, disables photosynthesis and strongly reduces growth and productivity (Di Castri and Mooney, 1973; Margaris, 1981). This is the reason why an investigation on the effect of the recently, massively introduced, additional EM radiation, produced by the numerous cellphone towers settled throughout the urban areas and the countryside, covering 95% of the Greek territories, seems more than interesting. Moreover, the global debate on the effects of the EM radiation on animal and plant life is continuously boosted and new data are always welcomed.

Myrtle (family: Myrtaceae) is an evergreen sclerophyllous species well adapted to the Mediterranean climatic conditions. It is a shrub or small tree thriving in typical Mediterranean formations, common all over Greece's lowlands thus very often exposed to non-ionizing electromagnetic radiation produced by the numerous phone masts of the continuously expanding cellular communication system.

The production and accumulation of secondary metabolites, phenolics in particular, is a common feature of the mediterranean plants (Christodoulakis, 1992, Fahn and Cutler, 1992). Secondary metabolites do have a multi-protective role and stressing conditions enhance the biosynthesis of these metabolites which then appear abundant in the vacuoles all leaf cells of the plant (Mitrakos and Christodoulakis, 1981).

After exposure to the summer drought stress, myrtle follows the habit of almost all mediterranean species (Nikolakaki and Christodoulakis, 2007), concerning the activation of the shikimic acid pathway. This metabolic pathway eventually leads stressed plants to produce various secondary metabolites whose biosynthesis is shared by animal and plant organisms (Tocher and Tocher, 1972; Iriti, 2013) including animal neurotransmitters i.e. acetylcholine in *Urtica* spp (Jaffe, 1970; Barlow and Dixon, 1973; Odjakova and Hadjiivanova, 1997). In myrtle, these metabolites are accumulated either within the

schizolysigenous gland cavities of the leaves or, mainly, in the form of phenolic tannin precursors or various types of tannins, within the vacuoles of all living cells, mostly those of the leaves.

Therefore, a) willing to exploit the numerous pieces of data on the stress response of the Mediterranean evergreen sclerophylls, b) having in mind that non-ionizing radiation proves to be a stressing factor for the plants by interfering with their secondary metabolic pathways, c) considering myrtle as a plant of wide distribution within the areas of Mediterranean climate, we set up a series of experiments to investigate the response of this plant to the radiation produced from a Global System for Mobile communications (GSM – mobile telephone) unit and work out on the results hoping to contribute to the reply on a global question mark.

2. Materials and methods

2.1. Plant material and radiation setup

Two myrtle individuals growing in a formation of Mediterranean evergreen sclerophylls, at the western slope of mount Hymettus (37°58'00.80"N, 23°47'07.35"E, altitude 259 m), north-East of Athens metropolitan area, were selected and young leaves of the previous vegetative period (autumn 2016) were marked with plastic tags each year.

Soon as the leaves reached their maturity (20 days after their emergence), the radiation procedure was applied. Leaves were irradiated with a GSM-1800 MHz mobile phone for 30 min, in talk mode, every Monday, Wednesday and Friday, at 08:00 am, from **02 Nov, 2016 until 21 Dec 2016** (1st experiment – 49 days) and **20 Feb 2017 until 10 Apr 2017** (2nd experiment – 50 days). The hand set (Motorola C115) was mounted on a gooseneck holder with a clamp at the end of the flexible extension arm which facilitated positioning the apparatus in touch with the exposed leaves. A tiny, remote controlled, battery operated transistor radio (Panasonic RF-H830) was also mounted facing the phone in order to provide continuous sound, simulate the talking mode and activate the handset to transmit towards the regional phone mast. The radio was tuned to a distant radio station; thus, the sound produced was actually full of radio parasites, a practically constant noise at high volume. The whole construction was tightly fixed on a heavy duty photographic tripod (Fig. 2 in Plate 2). The average E-field intensity value was **27 V/m**, for 6 min (according to ICNIRP, 1998). Maximum value of 58 V/m was recorded while background records were as low as 0,650 V/m. Measurements were taken with a NARDA SRM3000 (Germany) spectrum analyser while the GSM unit was transmitting. Supplementary measurements, for cross-checking, were taken with a broadband field meter (TES-92, 50 MHz - 3.5 GHz, Electromagnetic radiation detector – TES Electrical Electronic Corp. Taipei, Taiwan, R.O.C.). Values recorded were identical.

Investigation of the leaves was scheduled and executed as indicated in Fig. 1.

2.2. Leaf anatomy – microscopy

At the end of each experiment and when the plants had been exposed to twenty-two (22) doses of radiation, five leaves from the control and five from the exposed plants were cut-off, in random, a piece from the centre of each leaf was detached (the central nerve removed), cut into numerous small pieces (1 × 1 mm) and fixed in phosphate buffered 3% glutaraldehyde (pH 6.8) at 0 °C for 2 h. The tissue was post fixed, overnight, in 1% osmium tetroxide in phosphate buffer. A few pieces from the fixed tissue were dehydrated in graded acetone series, critical point dried, coated with gold and viewed with a JEOL JSM-6530 Scanning Electron Microscope. The rest of the tissue was dehydrated in graded ethanol series and embedded in Durcupan ACM (Fluka, Steinheim, Switzerland). Semithin sections, obtained from a LKB Ultratome III, were placed on glass slides and stained with 0.5 toluidine blue O (in 1% borax solution), as a general stain, for light microscopic




Plate 2. (Figs. 2–7): Experiment set-up, gland count on fresh whole mounts and cross sections from the two leaf types (both experiments) of myrtle. **Fig. 2.** The GSM handset, the marked myrtle individual (1, at the bottom) and the leaves under exposure are demonstrated; **Fig. 3.** Low magnification micrograph of a leaf whole mount. Glands are marked with circles; **Fig. 4.** Cross section of a control leaf (1st experiment). A gland can be observed just below the adaxial epidermis; **Fig. 5.** Cross section of an exposed leaf (1st experiment). The mesophyll cells, impregnated with osmiophillic material and a gland, just below the adaxial epidermis, can be observed; **Fig. 6.** Cross section of a control leaf (2nd experiment). A gland and a crystal can be observed just below the adaxial epidermis; **Fig. 7.** Cross section of an exposed leaf (2nd experiment). The mesophyll cells, impregnated with osmiophillic material and a gland, just below the adaxial epidermis and a few crystals below the adaxial epidermis, can be observed. **ad ep:** adaxial epidermis; **ab ep:** abaxial epidermis; **pp:** palisade parenchyma; **sp:** spongy parenchyma; **ct:** conductive tissue; **gl:** secretive structure; **cr:** crystal.

observations. Ultrathin sections were placed on 100 mesh grids, double stained with uranyl acetate and lead citrate (Reynolds, 1963) and viewed with a Phillips EM-300 Transmission Electron Microscope.

The whole procedure was repeated at the end of each experiment and the leaf tissue was subjected to identical treatment, i.e. fixed, dehydrated, embedded, sectioned, stained and observed. The results were thoroughly recorded for an oncoming evaluation and comparison. Literature for double fixation is cited in detail by Christodoulakis et al., (2009) and Christodoulakis et al., (2010).

The procedures for protein extraction and determination of protein concentration where repeated: 1) the 1st day, after the first radiation dose, 2) at the middle of the exposure period when the leaves had been

exposed to 11 doses of radiation and 3) at the end of the experiment, when the radiation process was fulfilled (22 doses) (Fig. 1).

2.3. Histochemistry

Fresh leaves were sectioned by hand or with a cryotome (Leica CM1850, Germany) at –10 °C and stained with histo-chemical reagents: (a) osmium tetroxide for unsaturated lipids; (b) antimony trichloride (SbCl₃) for terpene containing steroids; (c) vanillin/HCl for flavonoids; (d) concentrated H₂SO₄ for sesquiterpenes; (e) Wagner’s reagent for alkaloids; (f) Dittmar’s reagent for alkaloids; (g) iodine–iodide solution for alkaloids; (h) potassium bichromate for tannins; (i)

1 dose/1 st day			11 doses/24 days			22 doses/49 days		
preliminary 1st experiment (2 Nov 2016 to 21 Dec 2016) – anatomical investigation								
			25 November 2016 leaf morphology			21 December 2016 leaf morphology, tissue fixation & microscopy (OM, SEM, TEM)		
2nd experiment (20 Feb 2017 to 10 Apr 2017) – anatomical and biochemical investigation								
20 February 2017			15 March 2017 leaf morphology			21 April 2017 leaf morphology, tissue fixation & microscopy (OM, SEM, TEM)		
immediately	1 hr after exposure	24 hrs after exposure	immediately	1 hr after exposure	24 hrs after exposure	immediately	1 hr after exposure	24 hrs after exposure
ROS pigments DDC			ROS pigments DDC			ROS pigments DDC		
						1		

Fig. 1. Timetable for the investigation of the myrtle leaves (both experiments).

DMB (dimethoxy benzaldehyde or veratral aldehyde) for phenolic tannin precursors; (j) ferric chloride for polyphenols; (k) alcoholic vanillin/HCl (vanillin test) for phenolic compounds; (l) 4-nitrosophenol in concentrated H_2SO_4 for monoterpenic phenols; (m) nitroso reaction for catechol tannins. All stains were matched by controls. The literature for histochemical reactions is cited in detail by Christodoulakis et al. (2009) and Christodoulakis et al. (2010).

2.4. Protein extraction and determination of protein concentration

Leaves from both control and exposed samples were detached and grinded in liquid nitrogen. Protein extraction was performed with RIPA lysis buffer (Thermo Fisher Scientific Inc. – NYSE: TMO), samples were centrifuged ($12,000 \times g$, 30 min, 4°C) and the supernatants were stored at -20°C . Total protein concentration was determined according to the method of Bradford (Bradford, 1976), using Bovine Serum Albumin (BSA) as standard.

2.5. SDS-polyacrylamide gel electrophoresis and immunoblotting

SDS-polyacrylamide gel electrophoresis and immunoblotting SDS-PAGE was performed on a “Biorad Mini Protean” electrophoresis apparatus as described by Laemmli (1970), using a 12% polyacrylamide slab gel. Electrophoresis was carried out at 120 V for 2 h at room temperature. Following electrophoresis, samples were transferred onto nitrocellulose membrane (Towbin et al., 1979). Immunological detection of the proteins was performed according to Batteiger’s method (Batteiger et al., 1982) and subsequent immunological detection was performed using the antibodies described below.

2.6. Antibodies

A specific polyclonal L-Dopa DeCarboxylase (DDC) antibody raised against the purified human protein (Mappouras et al., 1990) and the DDC antibody raised from *Ceratitis capitata* (lab. of Prof. Dr D. Vassilacopoulou, University of Athens, GR – Bossinakou and Fragoulis, 1996) were used for immunodetection.

2.7. Pigments protocol

Chlorophyll pigments, from approximately 50 mg leaves, were extracted with 1 ml 80% acetone, overnight, at 4°C . Supernatant was transferred to a 1-ml glass cuvette for measurement in a Shimadzu Europe UV-1700 spectrophotometer. Absorbance was recorded throughout the visible spectrum, from 400 to 700 nm. We focused at both 663.6 and 646.6 nm, corresponding to *chlorophyll* –a and *chlorophyll* –b respectively and at 625 nm for the absorbance of *chlorophyll* –c. Quantification of pigment content was calculated using molar extinction coefficients specifically for this method (Gechev, 2013). *Chlorophyll* –a, $\epsilon_{663.6} = 76.79$ and $\epsilon_{646.6} = 18.58$; *chlorophyll* –b, $\epsilon_{663.6} = 9.79$ and $\epsilon_{646.6} = 47.04$. The total *chlorophyll* content is calculated using the following formula and normalize per fresh weight: $\text{Chlorophyll } a + b = 19,54A_{646.6} + 8,29A_{663.6}$. For beta carotenes and xanthophylls no formula is available for quantification; only the absorbance value is utilized.

2.8. Total ROS estimation

ROS levels were measured using 10 mM of the oxidant sensitive fluorescent acetyl ester CM- H_2DCFDA (5-(and-6)-chloromethyl-2', 7'-dichlorodihydrofluorescein diacetate) dissolved in DMSO. CM- H_2DCFDA is a general oxidative stress indicator that can enter cells by penetrating the cell membrane through passive diffusion. Inside the cell its acetate groups are cleaved by intracellular esterases and oxidation by ROS leads to the formation of fluorescent DCF product, which can be detected via fluorometry. The procedure was the following: Tissues

were homogenized in liquid nitrogen, placed in a 2 mL tube with 1 mL of 10 mM Tris–HCl, pH 7.2 and centrifuged at $12,000 \times g$ for 20 min at 4°C . Supernatant transferred to a fresh 2 mL tube and subsequently, the samples were placed continuously in incubator for 30 min with CM- H_2DCFDA diluted in 10 mM Tris–HCl at 37°C in the incubator. Then, the ester was removed and further incubation for 20 min in a Tris–HCl followed. Finally, tissues were washed three times with PBS buffer solution.

Measurement of fluorescence was made at the supernatant with VersaFluor Fluorometer System (Bio-Rad, 170- 2402, Hercules, CA) with excitation filter at 490 nm and emission at 520 nm. For the recovery experiments, various time points were tested as trials before finalizing the most promising values. In every set of experiments, duplicated samples were used for the exposed samples and the same run included control samples. Total ROS was expressed as fluorescent units/ μg of protein extracts (Sunkar, 2010).

3. Results

3.1. Morphology and microscopy

Control and exposed leaves were carefully examined for their external morphology. Exposed leaves, during the radiation process and at the end of the experiment, did not seem to be affected since their magnitude, shape, epidermal tissue arrangement and colour, appeared identical to their control counterparts.

Schizolysigenous gland distribution, microscopically measured on fresh leaves (Fig. 3 in Plate 2) (glands/ mm^2 : $8,2 \pm 0,8$ for the control and $9,1 \pm 0,6$ for the exposed leaves of the 1st experiment; $9,1 \pm 0,7$ for the control and $9,4 \pm 0,7$ for the exposed leaves of the 2nd experiment) as well as stomatal frequency (stomata/ mm^2 : 451 ± 28 for the control and 459 ± 35 for the exposed leaves of the 1st experiment; 438 ± 31 for the control and 445 ± 39 for the exposed leaves of the 2nd experiment), measured at low magnification OM (Fig. 3 in Plate 2) and SE micrographs, do not seem to vary significantly in the two leaf types.

Cross sections of both leaf types were investigated in detail (Fig. 4–7 in Plate 2). An obvious difference has to do with the leaf compactness and the accumulation of secondary metabolites in the cells of the mesophyll tissues (compare Figs. 4–7 in Plate 2). Exposed leaves appear more compact while palisade and spongy parenchyma cells accumulate large amounts of osmophilic material (Figs. 5 and 7 in Plate 2).

Cross sections of fresh leaves histochemically treated with DMB (dimethoxy–benzaldehyde), a reagent used to localise phenolic tannin precursors, reacted positively giving a vivid red colour for the cells of both adaxial and abaxial epidermis (Fig. 8 in Plate 3) as well as for the secretory cells of the glands (Figs. 8 and 9 in Plate 3). An intense brown tincture was obtained, in the mesophyll cells, after using potassium bichromate for tannins. Ferric chloride for polyphenols stained the epidermal and most of the mesophyll cells with a dark blue colour, nitroso – reaction for catechol tannins stained the mesophyll brown while a deep purple colour was obtained with alcoholic vanillin/HCl, a stain for phenolic compounds in general.

Scanning electron microscope observations confirmed the findings from the light micrographs since the mesophyll cells of the exposed leaves appear impregnated with a dense material (compare Figs. 14 and 15 in Plate 4).

A clear image was obtained from the Transmission Electron Microscopic observations. The mesophyll cells of the exposed leaves are tightly “packed” with secondary metabolites (sm in Figs. 16 and 17 in Plate 4) while the mature chloroplasts of these cells demonstrate their productive past by exhibiting the large starch granules they have accumulated (st in Figs. 16 and 17 in Plate 4). All membranes and organelles of the cells from the exposed leaves (i.e. mitochondria) seem well preserved without any distinguishable deformations.




Plate 3. (Figs. 8–13): Cross sections of fresh myrtle leaves treated with histochemical reagents. **Fig. 8** and **Fig. 9**, treated with DMB. **Fig. 10** and **Fig. 11**, treated with $K_2Cr_2O_7$. **Fig. 12**, treated with $FeCl_3$ solution. **Fig. 13**, treated with alcoholic vanillin/HCl. White arrows point at the secretory, epithelial cells of the glands.

3.2. Pigments and ROS estimation

For the photosynthetic pigments the measurements were executed three times: once after the first exposure to radiation, then in the middle (24 days) of the radiation treatment and, finally, after the end of the exposure series (49 days). Each time the pigments were measured immediately after irradiation, one hour after irradiation and 24 h after irradiation. The absorption graphs are presented in Fig. 18, 20 and 22 in [Plate 5](#) (left column).

It seems that, as time goes by, the absorbance graph dives to lower values. The minimum absorbance is observed 24 h after irradiation. Surprisingly a new peak appears, in every graph, at about 535 nm (Fig. 18, 20 and 22 in [Plate 5](#)), an area in the band where none of the photosynthetic pigments absorbs. This confined region of the band is “reserved” for the absorbance of the anthocyanins (537–542 nm – [Merzlyak et al., 2008](#)). Photosynthetic pigments do not seem to be affected during or immediately after irradiation. Yet they are reduced one hour after radiation and present a significant downfall, 24 h after their exposure (Fig. 18, 20 and 22 in [Plate 5](#)).

Concerning measurements of Reactive Oxygen Species (ROS) (Figs. 19, 21 and 23 in [Plate 5](#)) it seems that immediately after irradiation the difference between control and exposed leaves remains within the limits of the statistical error. Yet, as time lapses, the oxidative stress seems to increase. One hour after irradiation, ROS counts are almost 30% more in the exposed leaves, in all three cases, while after 24 h the increase of ROS climbs to about 50% compared to the control

leaves (Figs. 19, 21 and 23 in [Plate 5](#)).

3.3. Secondary metabolites of special interest

Our investigations with SDS-PAGE electrophoresis are presented in Figs. 24–26 in [Plate 6](#). Samples from the control leaves were compared to samples from their exposed counterparts. Exposed leaves treated for immunoblotting immediately after irradiation, do not present any detectable alteration compared to the control ones, concerning the expression of DDC. On the other hand, exposed leaves, one hour after irradiation, seem to express DDC at unexpectedly high levels. This can clearly be demonstrated in Figs. 24–26 in [Plate 6](#) where the expression of DDC appears cross checked with both antibodies (left column for anti-human DDC Ab and right column for anti-*Ceratitidis capitata* DDC Ab). In Fig. 24 in [Plate 6](#) DDC immunodetection appears at approximately 55 kDa at least one hour after irradiation (lane L3 in both columns). The expression of DDC remains at high levels in the exposed leaves even 24 h after irradiation (Fig. 24 in [Plate 6](#), lane L4 in both columns). After 11 exposures (22 days) (Fig. 25 in [Plate 6](#)) and 22 exposures (49 days) (Fig. 26 in [Plate 6](#)) the same results appear for the expression of DDC one hour and 24 h after irradiation, for both antibodies used.




Plate 4. (Figs. 14–17): Scanning and transmission electron micrographs from myrtle leaves. Scanning EM: **Fig. 14.** The leaf of a control plant. Mesophyll cells appear free from inclusions (yellow arrows). **Fig. 15.** The leaf of an exposed plant, 24 hours after the last irradiation (22 exposures). Most of the mesophyll cells appear filled with dense material (white arrows). Transmission EMs from the mesophyll of **exposed leaves:** **Fig. 16.** Palisade cells obliquely sectioned. The central vacuole appears full of secondary metabolites (**sm**). Chloroplasts (**chl**) exhibit their productive past through the numerous starch grains (**st**). **Fig. 17.** Palisade cell with the central vacuole loaded with osmiophilic secondary metabolites (**sm**). Chloroplasts with starch and oil droplets and a few mitochondria can be observed. **abep:** abaxial epidermis; **pp:** palisade parenchyma; **sp:** spongy parenchyma; **ct:** conductive tissue; **st:** starch; **sm:** secondary metabolites; **cw:** cell wall; **oil d:** oil droplet; **chl:** chloroplast; **m:** mitochondrion; **is:** intercellular space.

4. Discussion

4.1. Microscopy

Irradiated mature leaves of myrtle appear unaffected after the experiment. Stomatal frequency remains the same in both leaf types. The frequency of the oil glands seems not to be affected as well. Yet, the production and accumulation of phenolic compounds within the vacuoles of the cells of the exposed leaves indicates the presence of an environmental stress.

Considering the results of the histochemical reactions as well as their osmiophilic nature, the texture and the dark colour of the vacuole inclusions as seen in the transmission electron micrographs, we may assume that the compounds accumulated are of phenolic nature. They must be either phenolic tannin precursors or condensed phenolics, at various degrees of condensation (i.e. hydrolysable tannins or non-recyclable tannins). The accumulation of phenolics, even as dead-end products, has long ago been considered as a very common feature of the mediterranean plants (Dement and Mooney, 1974; Orshan, 1989; Dafni, 1991; Fahn and Cutler, 1992; Christodoulakis, 1992; Meletiyou-Christou et al., 1994; Stephanou and Manetas, 1997) – and not only (Ramakrishna and Ravishankar, 2011) – when sustaining environmental stress (drought or high temperature, high light intensity).

During our investigation, the environmental conditions were chosen, on purpose, to be moderate. No drought, no high temperatures, no high light intensity (winter months). In spite of these moderate conditions the accumulation of phenolics in the cells of the exposed leaves was remarkable compared to their control counterparts. Since the only stressing factor was radiation, we may assume that plants actually suffer when exposed to GSM radiation.

4.2. ROS

Increased activities of H_2O_2 and ROS in cells provide clear evidence for the vulnerability of plants to oxidative stress upon EMR exposure (Jayasanka and Asaeda, 2014). Reactive oxygen species are highly active compounds and excess availability in cells may lead to oxidation of

membranes and various cellular components that ultimately result in the oxidative destruction of the cells (Mittler 2002; Hung et al., 2005). Oxidative stress is probably responsible for the production of phenolics which play an important role for plants under stress conditions (Weidner et al., 2009).

4.3. Absorbance

The measurement of photosynthetic pigments can provide basic information on the physiological status of a plant. High-pressure liquid chromatography (HPLC) is the “state of the art” method for this purpose since it provides an accurate determination of a variety of photosynthetic pigments simultaneously. Yet this technique has a drawback compared to conventional spectroscopic techniques since it is more prone to structural modification of pigments during extraction, thus generating erroneous results (Hu et al., 2013). In the current investigation, the extraction method used has already been cross checked in previous investigations (Stefi et al., 2016, 2017a, 2017b) for not producing any artifactual conversion of chlorophylls.

An interesting detail of this investigation is that besides the absorption of the photosynthetic pigments, which falls significantly 24 h after the last dose, a new peak appears: that of the anthocyanins. Anthocyanin absorption maxima *in vivo*, as estimated with spectrophotometry of intact anthocyanic leaves and micro-spectrophotometry of vacuoles in the leaf cross-sections, were found between 537 nm and 542 nm, showing a red shift of 5–20 nm compared to the corresponding maxima in acidic water-methanol extracts (Merzlyak et al., 2008). Anthocyanins are synthesized in order to absorb the excessive solar energy and prevent photo damage. They appear in stressed leaves (Merzlyak et al., 2008). In our investigation, they are present even in the control leaves probably because of the “Mediterranean habits” of these leaves which start their life “red coloured” and obtain their normal green colour at maturity (Manetas et al., 2002; Liakopoulos et al., 2006). Although it is not known whether anthocyanins can scavenge reactive oxygen in living plant cells (Gould et al., 2002) some data support the hypothesis that “red cells” have elevated antioxidant capabilities *in vivo* (Gould et al., 2002).




Plate 5. (Figs. 19–23): Values for the concentration of the photosynthetic pigments (left column, Figs. 18, 20 and 22) and Reactive Oxygen Species – ROS (right column, Figs. 19, 21 and 23) from the myrtle leaves investigated after exposure to 1 dose (first row, 18 and 19), 11 doses (second row 20 and 21) and 22 doses (third row, 22 and 23).

Therefore, we may assume that since the stressing conditions of the Mediterranean climate were not present in our experiment, the drop of the photosynthetic pigment absorbance is due to radiation while the presence of anthocyanins is an indication that some other, non-detectable stressing factors have an impact on the leaves.

4.4. Secondary metabolite biosynthesis

L-DOPA is considered among the secondary metabolites whose biosynthesis is shared by animal and plant organisms (Tocher and Tocher, 1972; Iriti, 2013). These metabolites are involved, in concert with some phytohormones, in some regulatory functions in plant physiology such as growth, carbohydrate metabolism and some stress responses (Kulma and Szopa, 2007). The rapid, impressive increase of




Plate 6. (Figs. 24–26): Expression of DDC in control and irradiated myrtle leaves. Left column: Immunodetection with anti-human DDC antibody. Right column: Immunodetection with anti-*Ceratitis capitata* DDC antibody. Immunodetection after, **Fig. 24.** the first exposure (immediately, one hour, 24 hours); **Fig. 25.** eleven exposures (immediately, one hour, 24 hours); **Fig. 26.** twenty-two exposures (immediately, one hour, 24 hours).

DDC expression in the leaves of the exposed plants indicates the urgent need for the elimination of the volume of its substrate, that is L-DOPA, through decarboxylation of L-DOPA to another well-known mammal neurotransmitter (Kulma and Szopa, 2007), the catecholamine dopamine, an 3,4-dihydroxy-substituted phenyl ring possessing amine (Odjakova and Hadjiivanova, 1997). Dopamine is a secondary metabolite which can regulate, as most catecholamines, carbohydrate metabolism, serve as a precursor of some alkaloids, protect the plants against certain pathogens and is involved in nitrogen detoxification (Kulma and Szopa, 2007). Thus, the observed fluctuation of the DDC expression patterns could reflect a possible gene expression regulation event which governs the phenomenon revealed through our western blot experiments (Figs. 24–26 in Plate 6).

It can also help the plants defend themselves against detrimental climatic conditions, environmental pollutants, high solar irradiance and water deficit (Iriti, 2013). Considering all the above we may presume that the exposed leaves produce DDC and, consequently dopamine, because they experience an inevitable, severe environmental stress due to their exposure to radiation.

What seems very interesting in the whole experiment is that photosynthetic pigments, ROS and DDC expression appear not to be affected immediately but one hour after exposure and the detected expression continues to reflect the stress situation even 24 h after irradiation. Yet, it seems that before the next exposure, a recovery mechanism “resets” the cellular behaviour so that photosynthetic pigments return to their normal concentration, ROS drops down to normal and DDC expression is suppressed. This appears to be a marvellous mechanism of recovery which gives the plant the ability to sustain long term stress conditions although avoidance of the detrimental effects cannot be unlimited. The results of the current investigation are supported by a similar approach on the significance of the DDC gene family in poppy (*Papaver somniferum*) where the authors suggest that valuable clues may be provided by identifying the temporal, spatial, developmental and environmental cues that control expression of the various

gene family members (Faccini and De Luca et al., 1994).

Our results are in total agreement with the various responses that have been observed after exposure to EMF at the cellular, molecular and whole plant scale. It is beyond any doubt that numerous metabolic activities (ROS metabolism, Krebs cycle, pentose phosphate pathway, chlorophyll content, terpene synthesis etc.) are modified, gene expression is altered and growth reduced (stem elongation and dry weight) by exposure to EM radiation (Vian et al. 2016; Stefi et al. 2016, 2017a, 2017b, 2017c). These changes occur not only in the tissues directly exposed but also systemically in distant tissues. Although the long-term impact of these changes is rather unknown, it has to be a matter of speculation since the documents we have support the aspect that non-ionizing EM radiation cannot be considered as harmless. On the contrary, it proves to be a genuine environmental factor that readily evokes changes in plant metabolism (Vian et al. 2016).

Before coming to an end, we have to emphasize that, since a) ICNIRP (International Commission On Non-Ionizing Radiation Protection, an independent scientific organization responsible for providing advice on health hazard of non-ionizing radiation exposure) has set the SAR (Specific Absorption Rate) limit for the general public at 2 W per Kilogram (W/Kg) b) the EUROPEAN PARLIAMENT Directorate General for Research (European Parliament 2001) defined that Mobile phones must be low power radio devices transmitting and receiving microwave radiation at frequencies between 900 MHz and 1800 MHz, c) this type of radiation is considered as non-thermal, weak RF-EMF (Waldmann-Selsam et al. 2016; Halgamuge, 2017) and d) the cellular phone used (Motorola C115) has an FCC approval (09 June 2004 - FCC ID value: IHDT56EX1) with SAR 0.31 W/kg (head) or 0.44 W/kg (body) measured at 1800 MHz, we may assume that the stress response observed in the leaves of *M. communis* has a causal relationship with RF-EMF and renders a strong evidence for the non-thermal origin of the RF-EMF effects.

5. Conclusions

Concluding, we may resume our data from young plants of *Arabidopsis thaliana*, *Gossypium hirsutum* and *Zea mays* grown under continuous exposure to non-ionizing radiation and the detrimental effects of this radiation on plant development, biomass production, photosynthetic pigments and chloroplast ultrastructure (Stefi et al. 2016, 2017a, 2017b, 2017c) along with the severe stress signs observed in radiated mature leaves of *Myrtus communis* individuals and contribute a huge “con” to the serious speculation on the effects of non-ionizing radiation to plant life.

Funding

A part of this research was funded by the Hellenic Foundation for Research and Innovation (ΕΛΙΔΕΚ)  according to the framework for the financial support of doctoral candidates who already implement their doctoral theses in Greek Universities.

Acknowledgements

We thank Professor Maria Triantafyllou and Professor Aikaterini Kouli for their collaboration.

References

- Baladis, V.G., Brumelis, K., Kalvickis, O., Nikodemus, D., Tjarve, V.Z., 1996. Does the skrunda radio location station diminish the radial growth of pine trees? *Sci. Tot. Environ.* 180, 57–64.
- Barlow, R.B., Dixon, R.O.D., 1973. Choline acetyltransferase in the nettle *Urtica dioica* L. *Biochem. J.* 132, 15–18.
- Batteiger, B., Newhall, W.J., Jones, R.B., 1982. The use of Tween 20 as a blocking agent in the immunological detection of proteins transferred to nitrocellulose membranes. *J. Immunol. Methods* 55, 297–307.
- Beaubois, E., Girard, S., Lallechere, S., et al., 2007. Intercellular communication in plants: evidence for two rapidly transmitted systemic signals generated in response to electromagnetic field stimulation in tomato. *Plant. Cell Environ.* 30, 834–844.
- Bernatzky, A., 1986. Elektromagnetischer Smog – Feind des Lebens. *Der Naturarzt* 11, 22–25. <http://www.diwellenbrecher.de/pdf/bernatzky.pdf>.
- Bossinakou, K.S., Fragoulis, E.G., 1996. Purification and characterisation of L-DOPA decarboxylase from pharate pupae of *Ceratitis capitata*. A comparison with the enzyme purified from the white prepupae. *Comp. Biochem. Physiol. B: Biochem. Mol. Biol.* 113 (2), 213–220.
- Bradford, M.M., 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Ann. Biochem.* 72, 248–254.
- Cammaerts, M.C., Johansson, O., 2015. Effect of man-made electromagnetic fields on common Brassicaceae *Lepidium sativum* (cress d'Alinois) seed germination: a preliminary replication study. *Fyton* 84, 132–137.
- Choi, J., 2012. Key Global Telecom Indicators for the World Telecommunication Service Sector. International Telecommunication Union (ITU), Geneva, Switzerland.
- Christodoulakis, N.S., 1992. Structural diversity and adaptations in some Mediterranean evergreen Sclerophyllous species. *Environ. Exp. Bot.* 32, 295–305.
- Christodoulakis, N.S., Kogia, D., Mavroedi, D., Fasseas, C., 2010. Anatomical and cytochemical investigation on the leaf of *Teucrium polium* L., a pharmaceutical shrub of the Greek phrygic formations. *J. Biol. Res.* 14, 199–209.
- Christodoulakis, N.S., Lampri, P.N., Fasseas, C., 2009. Structural and cytochemical investigation on the leaf of silverleaf nightshade (*Solanum eleagnifolium*), a drought resistant alien weed of the Greek Flora. *Aust. J. Bot.* 57, 432–438.
- Christodoulakis, N.S., 1989. An anatomical study of seasonal dimorphism in the leaves of *Phlomis fruticosa*. *Ann. Bot.* 63, 389–394.
- Dafni, A., 1991. Sclerophyll, chemical defense and the evolution of the Mediterranean maquis species. *Fl. Veg. Mundi.* 9, 167–174.
- Dement, W.A., Mooney, H.A., 1974. Seasonal variation in the production of tannins and cyanogenic glucosides in the chaparral shrub, *Heteromeles arbutifolia*. *Oecologia (Berl.)* 15, 65–76.
- Mediterranean type ecosystems. Origin and structure. In: di Castri, F., Mooney, H.A. (Eds.), *Ecological Studies No 7*. Springer Verlag, Berlin.
- European Parliament, 2001. Directorate General for Research-Directorate A, STOA - Scientific and Technological Options Assessment Briefing Note N° 05/2001 EN PE nr. 297.563 Electromagnetic Fields and Health. <http://www.home-biology.gr/images/emfsafetylimits/EuropeanParliamentSTOA.pdf>.
- Faccini, P.J., De Luca, V., 1994. Differential and tissue-specific expression of a gene family for tyrosine/dopa decarboxylase in opium poppy. *J. Biol. Chem.* 269, 26684–26690.
- Fahn, A., Cutler, D.F., 1992. XEROPHYTES. *ENCYCLOPEDIA OF PLANT ANATOMY* – Gebrüder Borntraeger, D-1000 Berlin. pp: 44–46, 77–100. .
- Gechev, T., 2013. A simple and powerful approach for isolation of *Arabidopsis* mutants with increased tolerance to H₂O₂. *Induced Cell Death Methods in Enzymology*, vol. 527 Elsevier Inc <http://dx.doi.org/10.1016/B978-0-12-405882-8.00011-8>. ISSN 0076-6879.
- Gould, K.S., McKelvie, J., Markham, K.R., 2002. Do anthocyanins function as antioxidants in leaves? Imaging of H₂O₂ in red and green leaves after mechanical injury. *Plant Cell Environ.* 25, 1261–1269.
- Haggerty, K., 2010. Adverse influence of radio frequency background on trembling aspen seedlings: preliminary observations. *Int. J. For. Res.*, 836278. <http://dx.doi.org/10.1155/2010/836278>.
- Halgamuge, M.N., 2017. Review: weak radiofrequency radiation exposure from mobile phone radiation on plants. *Electromagn. Biol. Med.* 36, 213–235. <http://dx.doi.org/10.1080/15368378.2016.1220389>.
- Harte, C., 1972. Auslösung von Chromosomenmutationen durch Meterwellen in Pollenmutterzellen von *Oenothera*. *Chromosoma* 36, 329–337.
- Harte, C., 1950. Mutationsauslösung durch Ultrakurzwellen. *Chromosoma* 3, 140–147.
- Hu, X., Tanaka, A., Tanaka, R., 2013. Simple extraction methods that prevent the artifactual conversion of chlorophyll to chlorophyllide during pigment isolation from leaf samples. *Plant Methods* 9, 19–32.
- Hung, S.H., Yu, C.W., Lin, C., 2005. Hydrogen peroxide functions as a stress signal in plants. *Bot. Bull. Acad. Sin.* 46, 1–10.
- Hyland, G., 2005. How Exposure to Mobile Phone Base Station Signals can Adversely Affect Humans. Available from: <http://www.tetrawatch.net/papers/hyland-2005.pdf>.
- ICNIRP, 1998. Guidelines for limiting exposure to time-varying electric, magnetic and electromagnetic fields (up to 300 GHz). *Health Phys.* 74 (4), 494–522.
- Iriti, M., 2013. Plant neurobiology, a fascinating perspective in the field of research on plant secondary metabolites. *Int. J. Mol. Sci.* 14 (6), 10819–10821.
- Jaffe, M.J., 1970. Evidence for the Regulation of Phytochrome-mediated Processes in Bean Roots by the Neurohumor, Acetylcholine. *Plant Physiol.* 46, 768–777.
- Jayasanka, S., Asaeda, T., 2014. The significance of microwaves in the environment and its effect on plants. *Environ. Rev.* 22 (3), 220–228.
- Jerman, I., Berden, M., Ruzic, R., Skarja, M., 1998. Biological effects of TV set electromagnetic fields on the growth of spruce seedlings. *Electromagn. Biol. Med.* 17, 31–42.
- Kulma, A., Szopa, J., 2007. Catecholamines are active compounds in plants. *Plant Sci.* 172, 433–440.
- Kumar, A., Pal Singh, H., Batish, D.R., Kaur, S., Kumar-Kohli, R., 2015. EMF radiations (1800 MHz)-inhibited early seedling growth of maize (*Zea mays*) involves alterations in starch and sucrose metabolism. *Protoplasma* 253 (4), 1043–1049.
- Laemmli, U.K., 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680–685.
- Lerchl, D., Lerchl, A., Hantsch, P., et al., 2000. Studies on the effects of radio frequency fields on conifers. *Kurzmitteilung auf der Tagung der Bioelectromagnetics Society in München*. http://www.boomaantastingen.nl/EMF_and_conifers%5B1%5D.pdf.
- Liakopoulos, G., Nikolopoulos, D., Klouvatou, A., Vekkos, K.-A., Manetas, Y., Karabourniotis, G., 2006. The photoprotective role of epidermal anthocyanins and surface pubescence in young leaves of Grapevine (*Vitis vinifera* L.). *Ann. Bot.* 98 (1), 257–265.
- Manetas, Y., Drinia, A., Petropoulou, Y., 2002. High contents of anthocyanins in young leaves are correlated with low pools of xanthophyll cycle components and low risk of photoinhibition. *Photosynthetica* 40 (3), 349–354.
- Mappouras, D.G., Stiakakis, J., Fragoulis, E.G., 1990. Purification and characterization of L-dopa decarboxylase from human kidney. *Mol. Cell. Biochem.* 94, 147–156.
- Margaris, N.S., 1981. Adaptive strategies in plants dominating Mediterranean-type ecosystems. In: Di Castri, F., Goodall, D.W., Specht, R. (Eds.), *Mediterranean-type Shrublands*. Elsevier, Amsterdam, pp. 239–242.
- Meleti-Christou, M.S., Rhizopolou, S., Diamantoglou, S., 1994. Seasonal changes of carbohydrates, lipids and nitrogen content in sun and shade leaves from four Mediterranean evergreen sclerophylls. *Environ. Exp. Bot.* 34, 129–140.
- Merzlyak, M.N., Chivkunova, O.B., Solovchenko, A.E., Naqvi, K.R., 2008. Light absorption by anthocyanins in juvenile, stressed, and senescing leaves. *J. Exp. Bot.* 59, 3903–3911.
- Mitrakos, K., 1980. A theory for Mediterranean plant life. *Acta Oecol. – Oec. Plant.* 1 (15), 245–252.
- Mitrakos, K., Christodoulakis, N.S., 1981. Leaf structural diversity in Mediterranean evergreen Sclerophylls. In: Margaris, N.S., Mooney, H.A. (Eds.), *Components of Productivity of Mediterranean-climate Regions-Basic and Applied Aspects*. Dr W. Junk Publishers pp 21–25.
- Mittler, R., 2002. Oxidative stress, antioxidants and stress tolerance. *Trends. Plant Sci.* 7, 405–410.
- Mooney, H.A., Dunn, E.L., Shropshire, F., Song, L., 1970. Vegetation comparison between the Mediterranean climatic areas of California and Chile. *Flora* 159, 80–96.
- Mooney, H.A., Dunn, E.L., 1976. Convergent evolution of the mediterranean climate evergreen sclerophyll shrubs. *Evolution* 24, 292–303.
- Nikolaki, A., Christodoulakis, N.S., 2007. Secretory structures and cytochemical investigation of the leaf of *Phlomis fruticosa*, a seasonally dimorphic subshrub. *Secreting activity of the leaf-originating calluses*. *Flora* 202, 429–436.
- Odjakova, M., Hadjiivanova, C., 1997. Animal neurotransmitter substances in plants. *Bulg. J. Plant Physiol.* 23, 94–102.
- Orshan, G., 1989. *Plant Pheno-Morphological Studies in Mediterranean Type Ecosystems*. Springer, Netherlands. <http://dx.doi.org/10.1007/978-94-009-3107-7>. eBook, ISBN: 978-94-009-3107-7.
- Pesnya, D.S., Romanovsky, A.V., 2013. Comparison of cytotoxic and genotoxic effects of plutonium-239 alpha particles and mobile phone GSM 900 radiation in the *Allium cepa* test. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 750, 27–33.
- Pietruszewski, S., Muszyński, S., Dziwulska, A., 2007. Electromagnetic fields and

- electromagnetic radiation as non-invasive external stimulants for seeds (selected methods and responses). *Int. Agrophys.* 21, 95–100 ISSN: 0236-8722.
- Ramakrishna, A., Ravishankar, A.-G., 2011. Influence of abiotic stress signals on secondary metabolites in plants. *Plant. Sign. Behav.* 6 (11), 1720–1731. <http://dx.doi.org/10.4161/psb.6.11.17613>.
- Reynolds, E., 1963. The use of lead citrate at high pH as an electron-opaque stain in electron microscopy. *J. Cell Biol.* 17, 19–58.
- Roux, D., Vian, A., Girard, S., et al., 2006. Electromagnetic fields (900 MHz) evoke consistent molecular responses in tomato plants. *Physiol. Plant* 128, 283–288.
- Sandu, D.D., Goiceanu, C., Ispas, A., Creanga, I., Miclaus, S., Creanga, D.E., 2005. A preliminary study on ultra high frequency electromagnetic fields effect on black locust chlorophylls. *Acta. Biol. Hung.* 56, 109–117.
- Selga, T., Selga, M., 1996. Response of *Pinus sylvestris* L. needles to electromagnetic fields. Cytological and ultrastructural aspects. *Sci. Total. Environ.* 180, 65–73.
- Sharma, S., Parihar, L., 2014. Effect of mobile phone radiation on nodule formation in the leguminous plants. *Curr. World. Environ.* 9, 145–155.
- Sheridan, S., Kalkstein, L., Kalkstein, A., Greene, S., 2010. Heat-related mortality and heat watch-warming systems in the United States: recent developments. Abstracts from the ISEE 22nd Annual Conference, Seoul, Korea, 28 August – 1 September 2010, Climate Change and Environmental Health, O-29A1-1, Epidemiology.
- Stefi, A.L., Margaritis, L.H., Christodoulakis, N.S., 2016. The effect of the non-ionizing radiation on cultivated plants of *Arabidopsis thaliana* (Col.). *Flora* 223, 114–120.
- Stefi, A.L., Margaritis, L.H., Christodoulakis, N.S., 2017a. The effect of the non-ionizing radiation on exposed, laboratory cultivated upland cotton (*Gossypium hirsutum* L.) plants. *Flora* 226, 55–64.
- Stefi, A.L., Margaritis, L.H., Christodoulakis, N.S., 2017b. The effect of the non-ionizing radiation on exposed, laboratory cultivated maize (*Zea mays* L.) plants. *Flora* 233, 22–30.
- Stefi, A.L., Margaritis, L.H., Christodoulakis, N.S., 2017c. The aftermath of long-term exposure to non-ionizing radiation on laboratory cultivated pine plants (*Pinus halepensis* M.). *Flora* 234, 173–186.
- Stephanou, M., Manetas, Y., 1997. The effects of seasons, exposure, enhanced UV-B radiation, and water stress on leaf epicuticular and internal UV-B absorbing capacity of *Cistus creticus*: a Mediterranean field study. *J. Exp. Bot.* 48 (11), 1977–1985.
- Sunkar, R., 2010. Review: MicroRNAs with macro-effects on plant stress responses. *Semin. Cell Dev. Biol.* 21, 805–811.
- Tkalec, M., Malarić, K., Pevalek-Kozlina, B., 2005. Influence of 400, 900, and 1900 MHz electromagnetic fields on *Lemna minor* growth and peroxidase activity. *Bioelectromagnetics* 26, 185–193.
- Tkalec, M., Malarić, K., Pavlica, M., Pevalek-Kozlina, B., Vidaković-Cifrek, Ž., 2009. Effects of radiofrequency electromagnetic fields on seed germination and root meristematic cells of *Allium cepa* L. *Mutat. Res. Genet. Toxicol. Environ. Mutagen.* 672, 76–81.
- Tocher, R.D., Tocher, C.S., 1972. Dopa decarboxylase in *Cytisus scoparius*. *Phytochemistry* 11, 1661–1667.
- Towbin, H., Staehelin, T., Gordon, J., 1979. Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: procedure and some applications. *Proc. Natl. Acad. Sci. U. S. A.* 76, 4350–4354.
- Vian, A., Davies, E., Gendraud, M., Bonnet, P., 2016. Plant responses to high frequency electromagnetic fields. *Biomed. Res. Int.* 1830262 <http://dx.doi.org/10.1155/2016/1830262>. 13 pages.
- Volkrodt, W., 1987. Wer ist am Waldsterben schuld? Mikrowellensmog der Funk- und Nachrichtensysteme. *Raum Zeit.* 26, 53–62.
- Volkrodt, W., 1991. Droht den Mikrowellen ein ähnliches Fiasko wie der Atomenergie? *Wetter-Boden-Mensch* 4, 16–23.
- Waldmann-Selsam, C., Balmori-de la Puente, A., Breunig, H., Balmori, A., 2016. Radiofrequency radiation injures trees around mobile phone base stations. *Sci. Total Environ.* 572, 554–569.
- Weidner, S., Karolac, M., Karamać, M., Kosińska, A., Amarowicz, R., 2009. Phenolic compounds and properties of antioxidants in grapevine roots (*Vitis vinifera* L.) under drought stress followed by recovery. *Acta Soc. Bot. Pol.* 78, 97–103.
- World Health Organization, 2006. Electromagnetic Fields and Public Health. Available from: <http://www.who.int/mediacentre/factsheets/fs304/en/index.html>.