

SCIENTIFIC STUDIES 2021

A META-INTEGRATIVE QUALITATIVE STUDY ON THE HIDDEN THREATS OF SMART BUILDINGS/CITIES AND THEIR ASSOCIATED IMPACTS ON HUMANS AND THE ENVIRONMENT

Raveendran R, Tabet Aoul KA. A Meta-Integrative Qualitative Study on the Hidden Threats of Smart Buildings/Cities and Their Associated Impacts on Humans and the Environment. *Buildings*. 2021; 11(6):251. <https://doi.org/10.3390/buildings11060251>

Abstract

Smart buildings deploying 5G and the Internet of Things (IoT) are viewed as the next sustainable solution that can be seamlessly integrated in all sectors of the built environment. The benefits are well advertised and range from inducing wellness and monitoring health, amplifying productivity, to energy savings. Comparatively, potential negative risks are less known and mostly relate to cyber-security threats and radiation effects. This meta-integrative qualitative synthesis research sought to determine the possible underlying demerits from developing smart buildings, and whether they outweigh the possible benefits. The study identified five master themes as threats of smart buildings: a surfeit of data centers, the proliferation of undersea cables, the consternation of cyber-security threats, electromagnetic pollution, and E-waste accumulation. Further, the paper discusses the rebound impacts on humans and the environment as smart buildings' actualization becomes a reality. The study reveals that, although some aspects of smart buildings do have their tangible benefits, the potential repercussions from these not-so-discussed threats could undermine the former when all perspectives and interactions are analyzed collectively rather than in isolation.

Excerpt

4.4. Master Theme 4: Electromagnetic Pollution

This is generally an identified problem with the deployment of smart buildings using 5G and IoT devices. Many countries have called to ban 5G in general until impartial research data can be made available, and several researchers around the globe have submitted a "5G appeal" [105]. The radiation effects could range from causing headaches, insomnia, to DNA alteration, along with the possibility of creating other biological damages such as hormonal imbalances, reproductive issues, tumors, nerve damage, and eye damage [65,106]. Belpomme (2015) concluded from a comprehensive study that the EMF effect could worsen health conditions related to oxidative stress, a deficit in melatonin metabolism, and is more reflected among electro-sensitive people [107]. Several studies by bio-chemical and medical researchers found that high frequencies can significantly change the heart rate, chromatin (DNA complex and proteins), and melatonin, as well as other hormonal changes [59,71,108]. Kojima et al., (2018) revealed that though most of the effects were thermally related as millimeter frequencies are quickly absorbed by water, it can induce damaging effects without the heating of the tissues, i.e., nonthermal effects that are more dangerous [109].

As a guideline, the Federal Communication Commission (FCC) adopted the SAR (specific absorption rate) limit value of 1.6 W/Kg for 1 g of tissue approved by ANSI and IEEE [110]. However, current FCC regulations check only the SAR value, which is only a measure of the thermal effects; on the contrary, several studies have concluded that evaluations other than SAR are necessary to fully understand the impact of biological effects other than the thermal effect [111,112]. Scientific evidence suggests that even radiation limits well below the

regulatory standards cause severe damage to health even from 2G and 3G [67,105]. Hardell (2017) pointed out that the World Health Organization (WHO) and International Agency for Research on Cancer (IARC) have only classified the risk from wireless cellphones as carcinogenic 2B (for instance, potentially cancerous).

Buildings are generally subjected to electromagnetic radiation (EMR) pollution from two sources: extremely-low-frequency (ELF) and high-frequency wireless devices. Leukemia in children, immunization loss, genes and DNA alteration, cancers, and tumors have been associated with increased exposure from these indoor sources since the 1960s [112]. A smart building that is operated wirelessly with very high frequencies (up to 300 GHz) can put the occupants at risk, particularly the most vulnerable. Moreover, humans have natural bio-electromagnetism

[113] in them, and cells, tissues, and skin regeneration, including the sleep process, rely on natural frequencies from 0 to 30 Hz [114,115,116]. It has been reported that, regardless of the frequency level, being exposed to artificial frequencies is detrimental to human health [112].

Furthermore, 5G cellular networks deploy many small cells placed at shorter distances on poles and buildings [117], which can easily aggravate, to a greater extent, the biological effects [68,72]. Hence, many scientists, health professionals, and environmentalists have enquired about the potential problems of continually being in a smart building with numerous IoT devices emitting radiation at high frequencies, including bio-wearable devices [73,94].

Electropollution radiation can also be a hazard to the living organisms of the ecosystem [61,66,69,70]. This problem is specifically crucial for organisms (living on land and in water) that depend strongly on Earth's natural electromagnetic field for their nutrition and survival [63,112]. The most significant example of such phenomena is the collapse of bee colonies as their navigation is affected by wireless radiation, making them unable to return to their hives or even find food [62,118]. A study spanning almost a decade by Selsam et al., (2016) found out that trees are significantly damaged by radiation, particularly those situated near cellular base stations, and the damage intensifies with aging [119].

Open access paper: <https://www.mdpi.com/2075-5309/11/6/251>

STOCHASTIC DOSIMETRY ASSESSMENT OF THE HUMAN RF-EMF EXPOSURE TO 3D BEAMFORMING ANTENNAS IN INDOOR 5G NETWORKS

Bonato M, Dossi L, Chiaramello E, Fiocchi S, Tognola G, Parazzini M. Stochastic Dosimetry Assessment of the Human RF-EMF Exposure to 3D Beamforming Antennas in indoor 5G Networks. *Applied Sciences*. 2021; 11(4):1751. <https://doi.org/10.3390/app11041751>

Abstract

The deployment of near future 5G networks will introduce modifications in the population's exposure levels to radio-frequency electromagnetic fields (RF-EMFs). The present work aimed to face the challenge of studying the exposure variability in the presence of an access point (AP) at 3.7 GHz with 64 patch elements uniform planar array antenna and 3D beamforming capability. The novelty introduced in the methodology of the exposure's evaluation was the combining of traditional computational methods with a new approach based on stochastic

dosimetry, called polynomial chaos kriging method, in order to estimate the exposure levels for 1000 different antenna beamforming patterns with low computational efforts. The simulations were evaluated considering a child model and computing the specific absorption rate (SAR) in different tissues. The analysis of the results highlighted a high exposure variability scenario depending on the beamforming patterns of the array antenna and identified the ranges of elevation and azimuth angles of the main antenna beam that may cause the highest levels of exposure.

Open access paper: <https://www.mdpi.com/2076-3417/11/4/1751>

EMPIRICAL STUDY ON SPECIFIC ABSORPTION RATE OF HEAD TISSUES DUE TO INDUCED HEATING OF 4G CELL PHONE RADIATION

Bindhu Christopher, Sheena Mary Y, Mayeen Uddin Khandaker, P.J. Jojo. Empirical study on specific absorption rate of head tissues due to induced heating of 4G cell phone radiation. Radiation Physics and Chemistry. Volume 178, 2021. doi: 10.1016/j.radphyschem.2020.108910.

Abstract

Exposures to electromagnetic radiation mainly from the extended use of mobile phones may initiate biological damages in the human body at the macromolecular level. Several studies on human and animal models have shown significant changes in the functions of neural cells. Present empirical study analyses the thermal changes and the specific absorption rates (SAR) of brain, eye and skin tissues due to prolonged exposure to mobile phone radiation. A phantom, simulating human head with skin, skull and brain was used for the study. The Phantom was exposed to radiation for longer durations (600 s and more) and the temperature variations at different specific points were studied with sensitive thermocouple probes. SAR (1 g of contiguous tissue) values were determined using the variations of temperature and other parameters. The average rise in brain temperature was found to be 0.10 ± 0.05 °C at 30 mm deep in the brain and the estimated SAR was 0.66 ± 0.35 Wkg⁻¹. The increase in temperature for the eye socket was 0.03 ± 0.02 °C with SAR 0.15 ± 0.08 Wkg⁻¹. The average rise in temperature for skin was 0.14 ± 0.05 °C and the SAR was 0.66 ± 0.42 Wkg⁻¹. Although the measured SAR lie within the safe limit of 2 Wkg⁻¹ recommended by the international regulatory body, considering the tremendous growth in the number of mobile phone users and prolonged use of mobile phone in communication purposes, the cumulative effects could be a real concern for human health.

<https://www.sciencedirect.com/science/article/pii/S0969806X20302085>

ASSESSMENT OF DNA DAMAGE IN CHICK EMBRYO BRAINS EXPOSED TO 2G AND 3G CELL PHONE RADIATION USING ALKALINE COMET ASSAY TECHNIQUE

D'Silva MH, Swer RT, Anbalagan J, Bhargavan R..J Clin of Diagn Res. 2021; 15(1):AC01- AC04. doi: 10.7860/JCDR/2021/47115/14441.

Abstract

Introduction The cellular phones/mobile phones have emerged as the fastest growing man-made phenomenon ever discovered in the history. Controversies still exist among the scientific community regarding the ill-effects of Radiofrequency Radiation (RFR) exposure

from cell phones on biological tissues. The present study will provide an insight into the basic mechanisms by which RF fields interact with developing brain in an embryo.

Aim To assess the possible Deoxyribonucleic Acid (DNA) damage in developing brain of chick embryo following chronic exposure to Ultra-High Frequency/Radiofrequency Radiation (UHF/RFR) emitted from 2G and 3G cell phone.

Materials and Methods Fertilised hen eggs were divided into three groups. Experimental Group A (exposed to 2G radiation, 24 eggs), Experimental Group B (exposed to 3G radiation, 24 eggs) and Group C sham exposed control group (24 eggs). After the completion of scheduled duration of exposure (72 minutes per day), the chick embryos were sacrificed from 9th-12th day and the brains were dissected out. The chick embryo brains were then subjected to alkaline comet assay technique to assess the DNA damage. The results were statistically compared using one-way Analysis of Variance (ANOVA).

Results In the present study, the exposure of chick embryo brains to 2G and 3G cell phone radiation caused increased mean comet length ($p<0.001$), mean tail length ($p<0.001$), mean percentage of DNA in the tail ($p<0.001$) and mean tail moment ($p<0.01$) suggestive of increased DNA damage.

Conclusion The present study concludes that the RFR exposure caused significant increase in DNA damage in developing brain of chick embryos with changes more pronounced in 3G exposure group

Open access paper:

https://jcd.r.net/article_fulltext.asp?issn=0973709x&year=2021&volume=15&issue=1&page=AC01&issn=0973-709x&id=14441

SPECIFIC ELECTROMAGNETIC RADIATION IN THE WIRELESS SIGNAL RANGE INCREASES WAKEFULNESS IN MICE

Liu L, Deng H, Tang X, Lu Y, Zhou J, Wang X, Zhao Y, Huang B, Shi Y. Specific electromagnetic radiation in the wireless signal range increases wakefulness in mice. *Proc Natl Acad Sci U S A*. 2021 Aug 3;118(31):e2105838118. doi: 10.1073/pnas.2105838118.

Abstract

Electromagnetic radiation (EMR) in the environment has increased sharply in recent decades. The effect of environmental EMR on living organisms remains poorly characterized. Here, we report the impact of wireless-range EMR on the sleep architecture of mouse. Prolonged exposure to 2.4-GHz EMR modulated by 100-Hz square pulses at a nonthermal output level results in markedly increased time of wakefulness in mice. These mice display corresponding decreased time of nonrapid eye movement (NREM) and rapid eye movement (REM). In contrast, prolonged exposure to unmodulated 2.4-GHz EMR at the same time-averaged output level has little impact on mouse sleep. These observations identify alteration of sleep architecture in mice as a specific physiological response to prolonged wireless-range EMR exposure.

Excerpts

"Prolonged radiation of mice (Pos9) using the Pulse64W regimen, but not the Conti8W regimen, results in statistically significant increase of wakefulness (Figs. 2, 4, and 5). Notably, these two regimens have the same time-averaged radiation level over the 12-h sleep period, suggesting a key role for pulse modulation."

"A distinct implication of this study is the possibility that prolonged exposure to modulated 2.4-GHz radiation, such as the wireless signal, might also increase wakefulness for humans. However, assessing this possibility may take extraordinary effort because of the complex issues for human volunteers. Consistent with our conclusion on the Conti8W group, previous studies found no significant effect on human sleep by unmodulated radiation (33, 36). Whole-night exposure to Wi-Fi was found to affect the sleep microstructure of humans but had no significant acute effects on the sleep macrostructure (42). This result is consistent with prior studies on the effect of mobile phone and base station radiation on human sleep (24, 28, 33). Another study on human subjects found no significant effects of 900-MHz EMR with global system for mobile communications modulation, either on conventional sleep parameters or on power spectra (30)."

"In a collective exposure scenario, the average power density at close proximity is about 0.037 W/m² for a smartphone, 0.013 W/m² for a laptop, and 0.13 W/m² near the Wi-Fi router (1). These values are considerably lower than the time- and whole-body-averaged general public exposure limit of 10 W/m² or occupational exposure limit of 50 W/m² for 2–300 GHz suggested by International Commission on Non-Ionizing Radiation Protection (43). In our experiments, the measured spatial averaged power density for Conti8W is 36.80 ± 0.92 W/m². Pulse64W is expected to have the same power density. Importantly, the effective EMR dose for inducing a biological response in mice is likely to be different from that in humans. Therefore, the relatively high EMR dose of the Pulse64W regimen that causes increased wakefulness in mice could be markedly reduced in humans. An epidemiological survey among those who work under either very high or very low doses of wireless radiation may reveal some clues.

In this study, 2.4-GHz EMR is modulated by 100-Hz square pulses, which have sharp edges and thus might have some unanticipated impact on neural activity in the brain. Additional experiments should be performed to examine whether other modulation functions such as sinusoidal modulation can induce similar increase of wakefulness in mice. In addition, other modulation frequencies such as 10 and 1,000 Hz should be investigated to answer the question of whether increased wakefulness is specific to certain modulation frequencies. Finally, both the intensity and the frequency of the carrier EMR (2.4 GHz in this study) should be scrutinized."

Open access paper: <https://www.pnas.org/doi/10.1073/pnas.2105838118>

EXPOSURE TO ELECTROMAGNETIC FIELDS FROM SUBMARINE POWER CABLES CAN TRIGGER STRENGTH-DEPENDENT BEHAVIOURAL AND PHYSIOLOGICAL RESPONSES IN EDIBLE CRAB

Scott K, Harsanyi P, Easton BAA, Piper AJR, Rochas CMV, Lyndon AR. *Cancer pagurus* (L.). Journal of Marine Science and Engineering. 2021; 9(7):776. <https://doi.org/10.3390/jmse9070776>

Abstract

The current study investigated the effects of different strength Electromagnetic Field (EMF) exposure (250 μ T, 500 μ T, 1000 μ T) on the commercially important decapod, edible crab (*Cancer pagurus*, Linnaeus, 1758). Stress related parameters were measured (L-Lactate, D-Glucose, Total Haemocyte Count (THC)) in addition to behavioural and response parameters (shelter preference and time spent resting/roaming) over 24 h periods. EMF strengths of 250 μ T were found to have limited physiological and behavioural impacts. Exposure to 500 μ T and 1000 μ T were found to disrupt the L-Lactate and D-Glucose circadian rhythm and alter THC. Crabs showed a clear attraction to EMF exposed (500 μ T and 1000 μ T) shelters with a significant reduction in time spent roaming. Consequently, EMF emitted from MREDS will likely affect crabs in a strength-dependent manner thus highlighting the need for reliable in-situ measurements. This information is essential for policy making, environmental assessments, and in understanding the impacts of increased anthropogenic EMF on marine organisms.

Conclusions

The results obtained from this study, combined with the data from previous work [13,41], suggests that increased physiological stress will occur if *C. pagurus* is exposed to EMF of 500 μ T or above with data obtained at 1000 μ T, 2.8 mT and 40 mT confirming this trend. This is mirrored in the behavioural trends noted, which showed an attraction to EMF sources at the same levels despite the physiological ramifications.

This suggests that a working limit of a maximum of 250 μ T could result in minimal physiological and behavioural changes within this species and should be considered during MRED design and implementation. Additional research is required to further identify sensitivities to EMF in different life stages and conditions within this species and benthic crustaceans in general. 250 μ T = 2,500 Milligauss

Open access paper: <https://www.mdpi.com/2077-1312/9/7/776>

THE PROTECTIVE EFFECTS OF EMF-LTE AGAINST DNA DOUBLE-STRAND BREAK DAMAGE IN VITRO AND IN VIVO

Hee Jin, Kyuri Kim, Ga-Young Park, Minjeong Kim, Hae-June Lee, Sangbong Jeon, Ju Hwan Kim, Hak Rim Kim, Kyung-Min Lim, Yun-Sil Lee. 2021. *International Journal of Molecular Sciences* 22, 10: 5134. doi: 10.3390/ijms22105134.

Abstract

With the rapid growth of the wireless communication industry, humans are extensively exposed to electromagnetic fields (EMF) comprised of radiofrequency (RF). The skin is considered the primary target of EMFs given its outermost location. Recent evidence suggests that extremely low frequency (ELF)-EMF can improve the efficacy of DNA repair in human cell-

lines. However, the effects of EMF-RF on DNA damage remain unknown. Here, we investigated the impact of EMF-long term evolution (LTE, 1.762 GHz, 8 W/kg) irradiation on DNA double-strand break (DSB) using the murine melanoma cell line B16 and the human keratinocyte cell line HaCaT. EMF-LTE exposure alone did not affect cell viability or induce apoptosis or necrosis. In addition, DNA DSB damage, as determined by the neutral comet assay, was not induced by EMF-LTE irradiation. Of note, EMF-LTE exposure can attenuate the DNA DSB damage induced by physical and chemical DNA damaging agents (such as ionizing radiation (IR, 10 Gy) in HaCaT and B16 cells and bleomycin (BLM, 3 μ M) in HaCaT cells and a human melanoma cell line MNT-1), suggesting that EMF-LTE promotes the repair of DNA DSB damage. The protective effect of EMF-LTE against DNA damage was further confirmed by attenuation of the DNA damage marker γ -H2AX after exposure to EMF-LTE in HaCaT and B16 cells. Most importantly, irradiation of EMF-LTE (1.76 GHz, 6 W/kg, 8 h/day) on mice in vivo for 4 weeks reduced the γ -H2AX level in the skin tissue, further supporting the protective effects of EMF-LTE against DNA DSB damage. Furthermore, p53, the master tumor-suppressor gene, was commonly upregulated by EMF-LTE irradiation in B16 and HaCaT cells. This finding suggests that p53 plays a role in the protective effect of EMF-LTE against DNA DSBs. Collectively, these results demonstrated that EMF-LTE might have a protective effect against DNA DSB damage in the skin, although further studies are necessary to understand its impact on human health.

Open access paper: <https://www.mdpi.com/1422-0067/22/10/5134>

PRELIMINARY STUDY: THE ELECTROMAGNETICS RADIATION FROM EPIPREMNUM AUREUM (MONEY PLANT) USING FREQUENCY DETECTOR

F. A. A. Shahrien, Z. H. Murat, R. S. S. A. Kadir. Preliminary Study: the Electromagnetics Radiation (EMR) from Epipremnum Aureum Using Frequency Detector. 2021 6th IEEE International Conference on Recent Advances and Innovations in Engineering (ICRAIE), 2021, pp. 1-5, doi: 10.1109/ICRAIE52900.2021.9703990.

Abstract

Epipremnum Aureum or money plant is a popular indoor plant known to emanate positive energy flow into space. Energy or aura is essentially the frequency radiated from a body, including from plants, and scientifically known as Electromagnetics Radiation (EMR). This preliminary study is to investigate the EMR pattern from money plant using frequency detector. The EMR are measured in a common sized room during five specific time range each day for three weeks with the presence of money plants and for two weeks without money plants (absence). The seven observation points of the EMR are taken around the money plant and in the middle of the room. The results shows that the EMR increases by 24% in the presence of money plant. This finding asserts that money plant radiates and influence the EMR frequency to increase in the surrounding.

<https://ieeexplore.ieee.org/document/9703990>

ASSESSMENT OF VARIED MOBILE NETWORK TOPOLOGIES ON HUMAN EXPOSURE, MOBILE COMMUNICATION QUALITY AND SUSTAINABILITY

Margot Deruyck, German Castellanos, Wout Joseph, Luc Martens, Sven Kuehn, Niels Kuster. Assessment of varied mobile network topologies on human exposure, mobile communication quality and sustainability. Final Report of Project CRR-954. Zurich, Switzerland, IT'IS Foundation. Sep 21, 2021.

Executive Summary

In October 2020, the Swiss Federal Office of Communications (OFCOM) mandated the IT'IS Foundation to evaluate various 5G network topologies regarding human exposure, mobile communication quality, and sustainability to address the questions posed by the political Motion Häberli-Koller (19.4043) dated July 30, 2020. The study was conducted jointly with the IMEC WAVES group of the Department of Information Technology of Ghent University, Belgium, and it supplements an earlier project performed by IT'IS for the Swiss Federal Office for the environment in 2019 [1].

Statement of work. Prior to the study, OFCOM specified a number of endpoints in the statement of work (SoW), each of which is addressed below. The study was conducted using a mobile network planning tool developed by the IMEC WAVES group that was adapted to the specifics of the mobile network and regulatory situation in Switzerland. Using this tool, we simulated a variety of mobile networks to address the study endpoints specified by OFCOM. The simulations included 4G and 5G networks in rural, suburban, and urban environments with usage requirements extrapolated to the year 2030. We analyzed the effects of separate operators compared to a unified network, the separation of indoor and outdoor networks, and different data rates and networks optimized for low downlink exposure. Human exposure to the electromagnetic field from the mobile communication system is expressed as the exposure ratio, the induced 6-minute time-averaged electromagnetic field level divided by the safety limits in percent, separately for the uplink (exposure to a user's own mobile device) and downlink (exposure to the base station network). All mobile networks were realized to comply with an approximation to the current precautionary limits imposed by the Swiss Ordinance for Protection against Non-Ionizing Radiation, i.e., the effect of an increase of the precautionary limits was not studied.

Note that for all of the following statements, in general, the user's own mobile device (uplink) exposure ratio contributed to the user's total exposure on average with a minimum tenfold higher level than the downlink (base station) exposure. Therefore, for any active mobile device user, a reduction of the downlink exposure will always remain insignificant in terms of the overall exposure.

Which network structures are possible based on the technologies available today (4G, 5G, Wireless Local Area Network WLAN, etc.), and what influence do they have on the spatial distribution of radiation exposure of the population? In the present study, network infrastructures based on the 4G and 5G communication technologies were simulated. The use of WLAN as a supplementary link for indoor reception was not considered in the simulation models as 4G and 5G provide better spectral efficiency technology and output power control than WLAN. Therefore, the use of WLAN in terms of network quality and exposure reduction

is not considered beneficial. Other reasons such as network separation and costs may favor the use of WLAN.

Our results show that the transition from 4G to 5G will reduce human exposure in most simulated scenarios while offering a tenfold capacity.

A unified mobile network results in downlink exposure ratios similar to that of the largest user base in Switzerland; however, the unified network can serve twice as many users. Network unification would improve the uplink exposure compared to the uplink exposure in the networks of the second and third largest user base in Switzerland. A unified network would also reduce the number of required base station locations between - 13% and -50% (depending on technology and environment).

In particular, the frequently propagated approaches of the "St. Gallen model", the separation of indoor and outdoor coverage are to be analyzed in depth. We analyzed the coverage of indoor and outdoor locations as well as indoor and outdoor exposure. Our results show that a complete separation of indoor and outdoor coverage will lead to lower (factor of 4) downlink exposure outdoors and uncovered indoor locations on average. Indoor downlink exposure is not affected by the separation of indoor and outdoor coverage. Uplink exposure remained in the same range for separate indoor and outdoor networks despite additional building attenuation. We found a trend towards lower uplink exposure (factor of 4) of the 5G technology compared to 4G. This effect is likely related to the use of the MaMIMO (Massive Multiple Input Multiple Output) capabilities in 5G.

In addition, the advantages and disadvantages of using adaptive antennas, also with regard to the data rate and spatial distribution of radiation exposure, should be shown. The 5G network employing adaptive antenna systems is well suited to reduce human exposure while increasing the network capacity by a factor of 10.

Especially, in less densely populated environments, adaptive antennas improve the exposure ratio. In the rural environment, the exposure ratio is reduced by a factor of two, while in the urban and suburban environments the average exposure is not affected by the use of adaptive antennas. We found a clear trend towards lower uplink and downlink exposure ratios for the 5G networks using adaptive antennas compared to 4G networks not using adaptive antenna technology.

What influence do the various network structures have on the quality of mobile communication coverage in Switzerland? All networks in the study were planned with a user coverage of at least 95% such that a lack of coverage was compensated by additional base station sites. Our results show that coverage with 5G speeds compared to 4G speeds requires on average three times as many base stations. The coverage of only outdoor locations would primarily reduce the need for base stations in urban environments and only for 4G. 5G will result in a base station densification for outdoor coverage that is already suited for indoor coverage in many locations. Where insufficient, e.g., in large buildings, the full coverage can be obtained by supplementary 5G indoor base stations.

What influence do the various network structures have on the expansion of mobile communications networks in Switzerland (number of antenna systems)? The transition from a 4G to a 5G network with a tenfold data bandwidth requires, on average, three times more base stations in our network optimization simulations. Base station count is increased by 60% (4G) and 14% (5G) if indoor locations are also covered by the mobile network in the urban environment. The reduction decreases to 20% (4G) and 6% (5G) for the suburban environment and vanishes in the rural environment. Extending the data rate capacity by an additional factor of ten for 5% of the users resulted in only a few additional base station locations but in higher human exposure.

The study also demonstrated that future 5G networks can be realized without an increase of precautionary limits. The number of base stations is mostly driven by the data requirements and not by the exposure limits.

What influence does the number of mobile communication networks have on the radiation exposure of the population or would a single network lead to less radiation exposure than three separate networks? Our results show that a unification of the mobile network infrastructure does not change human exposure considerably compared to multiple operators. However, a unification of the network infrastructure could lead to a smaller number of required base station sites (-13% to -50% depending on the environment and communication system). Due to the user limitation per base station for 5G adaptive antennas, the possible site reduction is greater for 4G (-30% to -50%) than for 5G (-13% to -30%).

What does an ideal network structure look like in order to minimize radiation exposure for the population and at the same time ensure a good quality of cell phone coverage? To minimize human exposure to electromagnetic fields, network planning should always take into account both uplink and downlink exposure. For active users, the uplink exposure is tenfold higher than the downlink exposure. Our results also show that consideration of downlink-only exposure in the planning stage will not reduce the exposure for non-users (members of the society without their own mobile device). A network for minimizing the combined uplink and downlink exposure employs 5G technology and has a dense base station infrastructure, supplemented locally, such as in large buildings, by indoor base stations. Another important outcome is that the 5G base station density is mainly driven by the coverage requirements, i.e., lower base station count under higher limits is not expected. The same rationale leads to the conclusion that relaxed precautionary limits likely increase the uplink exposure, i.e., the overall exposure.

Although the frequencies above 6 GHz are not yet available in Switzerland, the study should also include the future use of these frequencies (millimeter waves, in Switzerland probably in the 24.25-27.5 GHz range). At present, the use of millimeter-wave technologies in Switzerland cannot be predicted. Therefore, we did not include millimeter waves in our mobile network simulation model. Based on initial roll-outs internationally, the current main application of 5G millimeter links is for the last mile. The last mile application (fixed wireless access) is not strictly limited to mobile communication. Here, wireless point-to-point links would replace copper or fiber links. Based on the point-to-point nature, human exposure to last-mile links is unlikely.

Recently, the first mobile devices (US models of Apple iPhone 12, Samsung S21) with millimeter-wave communication capabilities were placed on the market. This development is driven by the unavailability of the 3.5 GHz band in the United States (US) and the attempt to offload the majority of the data volume over millimeter waves. There is still very little use of millimeter-wave communications in the US [2]. Due to shadowing effects of the human body [3] and the highly directive beam-forming in this frequency range, it can be assumed that the usage of millimeter waves for mobile applications will also mostly be limited to line-of-sight situations. This may lower human exposure due to the highly directive and adaptive nature of the communication links required for signal quality reasons.

Study limitations. Even though the study includes the most currently advanced simulations on user exposure as a function of network topology, several assumptions were made due to the limitations of the tools and to missing information, which are described in detail in the Methods Section (Section 4). The impact of these assumptions on the results is discussed in detail in Section 6 of this report.

Remaining Knowledge Gaps. In the course of this study, we identified future work and research needs to fill the remaining knowledge gaps. As soon as a possible application of millimeter waves emerges in Switzerland, its influence on human exposure should be analyzed. In the present study, we applied harmonized, yet simplified models to analyze indoor exposure to mobile networks. To lower the uncertainties on exposure in indoor scenarios, additional indoor modeling would help to substantiate our results. They could further be strengthened by validation measurements in the up- and downlinks of real 4G and 5G networks in Switzerland. Other areas of future research include the extension of the networks with distributed MaMIMO, mixed technology networks as well as more realistic assignment and weighting of user and base station locations.

Excerpt

To the knowledge of the authors, this study is currently the most advanced study on user exposure as a function of network topology. In order to perform the study using the developed tools and available information, several assumptions were made and described in detail in the Methods Section (Section 4). The impact of these assumptions on the results are as follows:

- The use of simplified models for the propagation and environments (buildings, etc.) may not exactly represent the propagation environment present in the real mobile network environments.
- The chosen coverage goal of 95% of the users in all the study environments at any user location might be higher in some environments than in real networks.
- The selected configurations for the modeling of mobile networks may be different in the actual network implementation. A global 5 V/m downlink exposure limit and a 6 minute average of the exposure were applied to map the regulatory boundaries. This means that not all details of the current regulatory requirement with respect to mobile communication exposure could be modeled 1:1 in the simulation tool.
- Even though the precautionary limits were only approximated, the results imply that the networks can be realized under the current regulation.

- The results were based on average usage, yet due to the dominating effect of the uplink exposure, exposure ratios are strongly dependent on personal mobile device usage which was not studied in detail here.
- Base station locations in the extended set were assigned by the optimization algorithm irrespective of the actual building feasibility.
- Currently, there is relatively limited knowledge about the actual deployment and development of 5G in Switzerland.
- Only frequency ranges currently licensed for mobile communication use in Switzerland were included, i.e., millimeter-wave communication links (5G NR FR2) were neglected in the network planning.
- The statistical-based modeling of exposure and limited environmental details and resolution statements on the absolute levels of exposure have a higher level of uncertainty than relative comparisons between scenarios.
- Other sources of exposure than from the simulated mobile communication networks were not included.
- It was assumed that other mobile communication services, e.g., 2G, 3G, were not present in the analyzed scenarios.
- The currently discussed relaxation of the precautionary limits was not studied in detail; however, the simulations indicate that the 5G base station density is mainly driven by the coverage requirements, i.e., lower base station count under higher limits is not expected. Furthermore, [1] relaxed limits likely increase the uplink exposure, i.e., the overall exposure.

Open access report: <https://bit.ly/Deruyck2021>

STEREOLOGICAL STUDY ON THE EFFECT OF CARNOSINE ON PURKINJE CELLS IN THE CEREBELLUM OF RATS EXPOSED TO 900 MHZ ELECTROMAGNETIC FIELD

Arslan A, Acer N, Kesici H, Sonmez MF, Ertekin T, Gultekin M, Dagdelen U, Saracoglu OG. Stereological Study on the Effect of Carnosine on Purkinje Cells in the Cerebellum of Rats Exposed to 900 MHz Electromagnetic Field. Turk Neurosurg. 2021 Aug 28. doi: 10.5137/1019-5149.JTN.35313-21.2.

Abstract

Aim: Electromagnetic fields have been known to reduce the number of Purkinje neurons in the cerebellum, whereas carnosine has been known to protect them. This study aimed to evaluate the impact of carnosine on Purkinje neurons in rats exposed to a 900 Mhz electromagnetic field.

Material and methods: This study evaluated 24 rats divided into the following three different groups: a control group, a group exposed to the electromagnetic field, and a group that was injected with carnosine while being exposed to the electromagnetic field. The electromagnetic field group was exposed to a 900 Mhz electromagnetic field for an hour daily over 28 days. Thereafter, stereological analysis was performed histologically on cerebellar sections, and the number of Purkinje cells were counted.

Results: The electromagnetic field group had remarkably fewer Purkinje cell compared to control. The electromagnetic field group plus 20 mg of carnosine had significantly more total Purkinje cells compared to the electromagnetic field group (p 0.05).

Conclusion: The present study showed that electromagnetic field exposure decreases the number of Purkinje cell, whereas carnosine protected the cerebellum from neural damage induced by electromagnetic field exposure.

Open access paper: http://www.turkishneurosurgery.org.tr/pdf/JTNEPUB_35313_online.pdf

EXPOSURE TO THE ELECTROMAGNETIC FIELD ALTERS THE TRANSCRIPTOMIC PROFILE IN THE PORCINE ENDOMETRIUM DURING THE PERI-IMPLANTATION PERIOD

Kozłowska W, Drzewiecka EM, Paukšto L, Zmijewska A, Wydorski PJ, Jastrzebski JP, Franczak A. Exposure to the electromagnetic field alters the transcriptomic profile in the porcine endometrium during the peri-implantation period. *J Physiol Pharmacol*. 2021 Dec;72(6). doi: 10.26402/jpp.2021.6.04.

Abstract

A low-frequency electromagnetic field (EMF) is an environmental pollutant that may influence female reproduction. This research was undertaken to test the hypothesis that EMF causes alterations in the transcriptomic profile of the endometrium. This study investigated the in vitro effects of EMF treatment (50 Hz, 2 h) on global transcriptome alterations in the endometrium isolated from pigs during the peri-implantation period. The control endometrium was not treated with EMF. The EMF treatment altered the expression of 1561 transcriptionally active regions (TARs) in the endometrium. In the group of 461 evaluated DEGs, 156 were up-regulated (34%), 305 were down-regulated (66%) and 341 (74%) had known biological functions. A total of 210 long noncoding RNAs (lncRNAs) with changes in expression profiles, and 146 predicted RNA editing sites were also evaluated. Exposure to EMF changes the expression of genes encoding proteins that are involved in proliferation and metabolism in endometrial tissue. These results provide useful inputs for further research into the impact of EMF on molecular changes in the uterus during the peri-implantation period and, consequently, pregnancy outcome.

Open access paper:

https://www.jpp.krakow.pl/journal/archive/12_21/pdf/10.26402/jpp.2021.6.04.pdf

PSYCHOLOGICAL DRIVERS OF INDIVIDUAL DIFFERENCES IN RISK PERCEPTION: A SYSTEMATIC CASE STUDY FOCUSING ON 5G

Renato Frey. Psychological Drivers of Individual Differences in Risk Perception: A Systematic Case Study Focusing on 5G. *Psychol Sci*. 2021 Oct;32(10):1592-1604. doi: 10.1177/0956797621998312.

Abstract

What drives people's perceptions of novel risks, and how malleable are such risk perceptions? Psychological research has identified multiple potential drivers of risk perception, but no studies have yet tested within a unified analytic framework how well each of these drivers accounts for individual differences in large population samples. To provide such a framework, I harnessed the deployment of 5G-the latest generation of cellular network technology. Specifically, I conducted a multiverse analysis using a representative population sample in Switzerland (Study 1; N = 2,919 individuals between 15 and 94 years old), finding that

interindividual differences in risk perceptions were strongly associated with hazard-related drivers (e.g., trust in the institutions regulating

5G, dread) and person-specific drivers (e.g., electromagnetic hypersensitivity)-and strongly predictive of people's policy-related attitudes (e.g., voting intentions). Further, a field experiment based on a national expert report on 5G (N = 839 individuals in a longitudinal sample between 17 and 79 years old) identified links between intraindividual changes in psychological drivers and perceived risk, thus highlighting potential targets for future policy interventions.

Excerpt

... three hazard-related drivers stood out in terms of their associations with interindividual differences in risk perceptions: Dread of 5G was strongly associated with higher risk perceptions, whereas trust in the institutions regulating 5G and objective knowledge about 5G were strongly associated with lower risk perceptions. Two more generic, person-specific drivers (electromagnetic hypersensitivity and gender) were further predictors of interindividual differences. Crucially, changes in risk perception that occurred within individuals across time were primarily associated with intraindividual changes in dread and trust, whereas changes in perceived benefit were primarily associated with intraindividual changes in trust and objective knowledge. Finally, people's policy-related attitudes proved to hinge strongly on perceived risk (inversely) and benefit (positively) as well as to be sensitive to respective changes in these two dimensions that occurred across time.

Open access paper: <https://journals.sagepub.com/doi/10.1177/0956797621998312>

NEW TECHNIQUE TO STUDY THE EFFECT OF FIFTH GENERATION (5G) RADIATION ANTENNA ON HUMAN BODY

G. G. Emam, G. M. Amer, W. A. Mohamed, K. A. Mustafa. 2021 13th Biomedical Engineering International Conference (BMEiCON), 2021, pp. 1-12, doi:10.1109/BMEiCON53485.2021.9745216.

Abstract

With the invention of the 5G network, the world has witnessed response to change in communication networks and is working on an unprecedented scale. This raises the question of whether high frequencies and constant exposure to nonparticulate radiation (EMF) radiation have a negative impact on human health. Some scientists believe that these waves don't cause effects on the physical body. In this paper, simulation was done by using MATLAB R2019a. To do this simulation the tissues were divided into 5 groups according to the type of tissues which are nerve, connective, epithelial and muscle tissues. The specific absorption rate (SAR) and power density (PD) in different frequencies (20, 30, 40, 60, 70, 80, 100 GHz) for more than 40 tissues are calculated. The comparison is done with other researches which have used the same techniques but most of these papers used the effect of the above frequencies on two or three tissues. In this paper, about 40 types of tissue are used for the simulation, and the simulation results show that the nerve tissues are the most affected tissue.

Excerpt

Tissues have four major types which are connective tissues, nerve tissues, epithelial tissues, and muscle tissues. It is found that the nerve tissues are the most tissue affected by the

chosen frequencies than other tissues understudy. The connective tissue and the epithelia tissue are affected by radiation in the same range. The least affected is muscle tissues. It was found that the breast fat is the most effective (affected) by radiation although it is epithelia tissues. We can refer to this as abnormal jump that we the breast fat is outside the body.

Conclusions

This article examines the influence of 5G networks on the human body. The importance of EMF exposure on various organs in downlinks at frequencies above 20 GHz is highlighted. Following the simulation, we discovered that 5G resulted in increased EMF exposure due to the usage of stronger directionality in electromagnetic beams and smaller cell size.

To do this simulation the tissues were divided into 5 groups according to the type of tissues which are nerve, connective, epithelial and muscle tissues. The specific absorption rate (SAR) and power density (PD) in different frequencies (20, 30, 40,60,70,80,100 GHz) for more than 40 tissues are calculated. Results show that the nerve tissues are the most tissue affected by the chosen frequencies, then the connective tissues and the epithelia tissue, while muscle tissues were the least affected type.

<https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9745216&isnumber=9745198>

OCCUPATIONAL EXPOSURES TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS AND THE RISK OF CANCER

Rohan Mate, Ken Karipidis, Michael Abramson, Geza Benke, Sarah Loughran. Volume 13, Supplement:S207-S208. 2022. doi: 10.1016/j.shaw.2021.12.1400.

Abstract

Introduction: Previous cohort and case-control studies have investigated possible associations between occupational exposure to radiofrequency (RF) electromagnetic fields (EMF) and a range of cancers. The findings to date give no consistent evidence of a causal relation between occupational RF EMF exposure and cancer, however previous studies have too many deficiencies to rule out an association and further research is warranted. A key concern across all previous studies is the quality of the RF EMF exposure assessment.

Methods: We plan to overcome previous methodological shortcomings by applying a newly developed job exposure matrix (JEM) to three separate case-control studies investigating glioma, follicular lymphoma and multiple myeloma, respectively. We further plan to investigate the validity of the JEM by conducting personal exposure measurements of workers in the highest exposure occupations identified by the JEM. We will also investigate how the level of occupational RF EMF exposure correlates with the level of risk perception to environmental RF EMF exposure.

Results: Initial results from this long-term project will be presented on the possible risk of occupation RF EMF and cancer. We will also present initial results on the agreement between local measurements and the highest exposure occupations identified by the JEM.

Conclusion: Our investigation will add to the body of knowledge on whether occupational RF EMF is a risk factor for cancer outcomes.

<https://www.sciencedirect.com/science/article/pii/S2093791121014992>

EFFECT EXPOSURE OF MOBILE PHONE RADIATION ON BLOOD PARAMETERS IN RATS

Al Abas, MAA, AL-Hakkak, ZMM (2021). Effect Exposure of Mobile Phone Radiation on Blood Parameters in Rats. Iraqi Journal of Science. 62(11): 4225–4231. doi: 10.24996/ijs.2021.62.11(SI).2.

Abstract

The aim of the research was to assess effects of short and long-period exposure to radiation from mobile phone on blood indices in experimental rats. In this study forty mature female rats were used; the animals were divided into two experimental group, each group consists of twenty animals. Short-period group of rats were exposed to cell phone radiation for different duration 30 m, 60 m, and 90 m per day for six weeks. Long-period group of rats were exposed to radiation from mobile phone for different duration 2h, 4h, and 6h per day for three months. The study noticed that there was significant ($P \geq 0.05$) elevation in total white blood cells and the study demonstrated significant increment ($P \geq 0.05$) in percentage of lymphocytes (71.1%) of rats which exposed to radiation from mobile phone for short-period in (90 minutes) compared to the control group (42.64%). While the study revealed that there was a significantly ($P \geq 0.05$) lower percentage of neutrophils (15.36%) in rats that were exposed to mobile phone radiation for long-period in (6 hours) compared with the control group (52.12%). The study recorded that there was a significant ($P \geq 0.05$) elevation in total red blood corpuscles and packed cell volume of rats exposed to radiation from mobile phone for short and long-period in different times compared with control. On the other hand, the research indicated that there was a significant ($p \leq 0.05$) decrement in mean corpuscles volume (MCV), mean corpuscle hemoglobin (MCH), mean corpuscle hemoglobin concentration and red blood cell distribution width standard deviation (RDW-SD) of rats to radiation from mobile phone for short and long-period in different times than control group.

Conclusions

It can be concluded that hematological parameters of rats were significantly affected after short and long-period exposure to radiation from mobile phone. The hematological analysis showed a significant elevation in values of white blood cells, lymphocytes, total red blood corpuscles and packed cell volume of rats after exposure to short and long-period to electromagnetic radiation from mobile phone. However, the study confirms that values of mean corpuscles volume and mean corpuscle hemoglobin concentration and red blood cell distribution width standard deviation were significantly decreased in rats after exposure to mobile phone electromagnetic radiation.

Open access paper:

<https://ijs.uobaghdad.edu.iq/index.php/eijs/article/download/3742/1879>

E-FIELD STRENGTH MEASUREMENTS OF A 5G BASE STATION IN 28 GHZ BAND FOR EMF EXPOSURE ASSESSMENT

Sen Liu, Teruo Onishi, Masao Taki, Miwa Ikuyo, Kazuhiro Tobita, Soichi Watanabe, Yukihiisa Suzuki2021 IEEE USNC- URSI Radio Science Meeting (Joint with AP-S Symposium), 2021, pp. 49-50, doi: 10.23919/USNC- URSI51813.2021.9703567.

Abstract

This paper presents the preliminary measurement results of the electric field (E-field) strength resulting from a fifth-generation (5G) base station operating in 28 GHz band. Three different measurement cases are discussed. The time-averaged (1 min) total E-field strength in all cases is no more than 91 dB μ V/m.

Conclusion

This paper presents the measurement results of the E-field strength from a 5G base station operating in 28 GHz band. Three different situations are considered, and the total E-field strength is much lower than the limit value of 155.76 dB μ V/m (61.4 V/m) in the corresponding band [4] for all cases. It is also found that although the instantaneous E-field strength during data traffic reaches 97.74 dB μ V/m, the RMS E-field strength during data traffic is only 0.4 dB higher than that without data traffic-on in this measurement. Although horn antennas have the advantage of discriminating downlink and uplink in TDD, an omnidirectional antenna will be used, and the corresponding results will be compared with those in this work in the future. In addition, comprehensive measurements and verifications will also be conducted.

<https://ieeexplore.ieee.org/document/9703567>

LESSONS LEARNED FROM A DISTRIBUTED RF-EMF SENSOR NETWORK

Sam Aerts, Günter Vermeeren, Matthias Van den Bossche, Reza Aminzadeh, Leen Verloock, Arno Thielens, Philip Leroux, Johan Bergs, Bart Braem, Astrid Philippron, Luc Martens, Wout Joseph. 2022. Lessons Learned from a Distributed RF-EMF Sensor Network. Sensors 22, no. 5: 1715. doi: 10.3390/s22051715.

Abstract

In an increasingly wireless world, spatiotemporal monitoring of the exposure to environmental radiofrequency (RF) electromagnetic fields (EMF) is crucial to appease public uncertainty and anxiety about RF-EMF. However, although the advent of smart city infrastructures allows for dense networks of distributed sensors, the costs of accurate RF sensors remain high, and dedicated RF monitoring networks remain rare. This paper describes a comprehensive study comprising the design of a low-cost RF-EMF sensor node capable of monitoring four frequency bands used by wireless telecommunications with an unparalleled temporal resolution, its application in a small-scale distributed sensor network consisting of both fixed (on building façades) and mobile sensor nodes (on postal vans), and the subsequent analysis of over a year of data between January 2019 and May 2020, during which slightly less than 10 million samples were collected. From the fixed nodes' results, the potential errors were determined that are induced when sampling at lower speeds (e.g., one sample per 15 min) and measuring for shorter periods of time (e.g., a few weeks), as well as an adequate resolution (30 min) for diurnal and weekly temporal profiles which sufficiently preserves short-term variations. Furthermore, based on the correlation between the sensors, an adequate density of 100 sensor nodes per km² was deduced for future networks. Finally,

the mobile sensor nodes were used to identify potential RF-EMF exposure hotspots in a previously unattainable area of more than 60 km². In summary, through the analysis of a small number of RF- EMF sensor nodes (both fixed and mobile) in an urban area, this study offers invaluable insights applicable to future designs and deployments of distributed RF-EMF sensor networks.

Open access paper: <https://www.mdpi.com/1424-8220/22/5/1715>

CHANGES IN PATTERN OF PRESENTATION OF PATIENTS WITH UNILATERAL VESTIBULAR SCHWANNOMA OVER TWO DECADES: INFLUENCE OF CELL PHONE USE IN EARLY DIAGNOSIS

Adeline Hephzibah, Sanjeev Kumar Pandey, V Rupa, Ranjith K Moorthy, Vedantam Rajshekhar. Changes in pattern of presentation of patients with unilateral vestibular schwannoma over two decades: Influence of cell phone use in early diagnosis. J Clin Neurosci. 2021 Dec;94:102-106. doi: 10.1016/j.jocn.2021.10.004.

Highlights

- Utility of cell phone usage in early detection of VS has not been studied.
- Cell phone usage aided in detecting hearing loss in 50% of patients.
- Majority of patients with VS still have severe hearing loss and large tumors.

Abstract

We aimed to study whether the increased use of cell phone in south Asia over the past two decades has impacted presentation of patients with vestibular schwannoma (VS). In this observational cohort study, data on cell phone use and severity of hearing loss and tumor size was collected using a questionnaire through a patient interview (n = 62) in consecutive patients with VS managed between 2017 and 2020. Association between cell phone use and severity of hearing loss and tumor size were studied and compared with our data published earlier when telephone use was not widely prevalent. 71% of the 62 patients (aged between 15 and 81 years) had large or giant VS. Pure tone audiometry showed moderately severe or profound hearing loss in 77.4%. Sixty (96.7%) patients used cell phones. 50% of patients complained of difficulty in using their cell phone because of hearing loss. Compared to the earlier era, a higher proportion consulted an ENT surgeon first for hearing loss (59.7% vs 26%; p = 0.0005) and had small or medium sized tumors (29% vs 8%; p = 0.008) with a smaller mean tumor size (3.3 vs 3.9 cm; p = 0.03). Increased cell phone use has led to earlier diagnosis of VS and smaller tumours in recent years when compared to an earlier era. However, the vast majority of patients continue to present with severe to profound hearing loss and large tumors. <https://pubmed.ncbi.nlm.nih.gov/34863422/>

Conclusions

Increased cell phone use has led to earlier diagnosis of VS and smaller tumours in recent years when compared to data from the same institution published 15 years back. A greater proportion of patients now present with mild to moderate hearing loss, have smaller tumours and consult an otolaryngologist for their problem. However, the vast majority of patients continue to present with severe to profound hearing loss and large tumors necessitating patient and physician education to further improve awareness regarding the need for early screening of patients with unilateral/asymmetrical sensorineural hearing loss with PTA and neuroimaging.

SHOULD SAR GUIDELINES INCLUDE VARIABILITY?

K. Masumnia-Bisheh, C. Furse. Should SAR Guidelines Include Variability? 2021 IEEE International Symposium on Antennas and Propagation and USNC-URSI Radio Science Meeting (APS/URSI), 2021, pp. 1945-1946, doi: 10.1109/APS/URSI47566.2021.9703740.

Abstract

Variations in 1-g and 10-g specific absorption rate (SAR) values due to variation in tissue dielectric properties are determined. To accomplish this goal, stochastic finite difference time domain (S-FDTD) and Monte Carlo methods are compared. A 3D head model with variable dielectric properties is exposed to a half wave dipole antenna at 835 and 1900 MHz. The results show that these variations can significantly affect the peak 1-g and 10-g SAR values and should be considered in SAR guidelines for cell phone assessment.

Conclusions

Variations in 1-g and 10-g SAR values caused by dielectric properties variation of the tissues are calculated in this paper. The results are obtained using S-FDTD and Monte Carlo methods for a head model adjacent to a half wave dipole antenna at 835 MHz and 1900 MHz frequencies. S-FDTD results were sufficiently close to the Monte Carlo results at 835 MHz, that we conclude that the more efficient S-FDTD approach can provide the assessment for this study.

At both frequencies, the standard deviation of the peak 1-g SAR is within 50% of the mean SAR, and the standard deviation of the peak 10-g SAR is approximately 40% of the mean. A typical interpretation of the standard deviation for a normally distributed random variable, is that we can have a 95% confidence that the values will fall within two standard deviations of the mean. The significance of these results is that the effect of tissue property variation is too large to be ignored in SAR guidelines. The normal, expected variance in SAR should be considered in SAR guidelines. We recommend the next step in this research is to model realistic cell phones, at current (5G) frequencies, to determine the mean and standard deviation of the 1-g and 10-g SAR levels. If these follow the same trends we are seeing from a dipole antenna, it is then imperative that the SAR guidelines be revisited, and that an assessment of the effect of variation in tissue electrical properties be included in them.

<https://ieeexplore.ieee.org/document/9703740>

COMPARING THE EFFECTS OF LONG-TERM EXPOSURE TO EXTREMELY LOW-FREQUENCY ELECTROMAGNETIC FIELDS WITH DIFFERENT VALUES ON LEARNING, MEMORY, ANXIETY, AND B-AMYLOID DEPOSITION IN ADULT RATS

Faraji N, Salehi I, Alizadeh A, Pourgholaminejad A, Komaki A, Taheri Azandaryani M, et al . Comparing the Effects of Long-term Exposure to Extremely Low-frequency Electromagnetic Fields With Different Values on Learning, Memory, Anxiety, and β -amyloid Deposition in Adult Rats. BCN. 2021; 12 (6) :849-860

Abstract

Introduction: Extremely Low-Frequency Electromagnetic Fields (ELF-EMFs) have gathered significant consideration for their possible pathogenicity. However, their effects on the nervous system's functions were not fully clarified. This study aimed to assay the impact of

ELF-EMFs with different intensities on memory, anxiety, antioxidant activity, β -amyloid ($A\beta$) deposition, and microglia population in rats.

Methods: Fifty male adult rats were randomly separated into 5 groups; 4 were exposed to a flux density of 1, 100, 500, and 2000 microtesla (μT), 50 Hz frequency for one h/day for two months, and one group as a control group. The control group was without ELF-EMF stimulation. After 8 weeks, passive avoidance and Elevated Plus Maze (EPM) tests were performed to assess memory formation and anxiety-like behavior, respectively. Total free thiol groups and the index of lipid peroxidation were evaluated. Additionally, for detection of $A\beta$ deposition and stained microglia in the brain, anti- β -amyloid and anti-Iba1 antibodies were used.

Results: The step-through latency in the retention test in ELF-EMF exposure groups (100500 & 2000 μT) was significantly greater than the control group ($P < 0.05$). Furthermore, the frequency of the entries into the open arms in ELF-EMF exposure groups (especially 2000 μT) decreased than the control group ($P < 0.05$). No $A\beta$ depositions were detected in the hippocampus of different groups. An increase in microglia numbers in the 100, 500, and 2000 μT groups was observed compared to the control and one μT group.

Conclusion: Exposure to ELF-EMF had an anxiogenic effect on rats, promoted memory, and induced oxidative stress. No $A\beta$ depositions were detected in the brain. Moreover, the positive impact of ELF-EMF was observed on the microglia population in the brain.

<http://bcn.iums.ac.ir/article-1-1740-en.html>

AUDITORY EFFECTS OF MICROWAVE RADIATION

James C. Lin. Auditory Effects of Microwave Radiation. Springer, 2021. ISBN 978-3-030-64543-4 ISBN 978-3-030-64544-1 (eBook). doi:10.1007/978-3-030-64544-1

This book examines the human auditory effects of exposure to beams of high-power microwave pulses, which research results have shown can cause a cascade of health events when aimed at a human subject or the subject's head. The book details multidisciplinary investigations using physical theories and models, physiological events and phenomena, and computer analysis and simulation. Coverage includes brain anatomy and physiology, dosimetry of microwave power deposition, microwave auditory effect, interaction mechanisms, and shock/pressure wave induction. The book will be welcomed by scientists, academics, health professionals, and practicing engineers as an important contribution to the continuing study of the effects of microwave pulse absorption on humans.

<https://doi.org/10.1007/978-3-030-64544-1>

EXPOSURE ASSESSMENT TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS IN OCCUPATIONAL MILITARY SCENARIOS: A REVIEW

Gallucci S, Fiocchi S, Bonato M, Chiaramello E, Tognola G, Parazzini M. Exposure Assessment to Radiofrequency Electromagnetic Fields in Occupational Military Scenarios: A Review. *Int J Environ Res Public Health*. 2022 Jan 14;19(2):920. doi: 10.3390/ijerph19020920.

Abstract

Background: Radiofrequency radiations are used in most devices in current use and, consequently, the assessment of the human exposure to the radiofrequency radiations has become an issue of strong interest. Even if in the military field there is wide use of radiofrequency devices, a clear picture on the exposure assessment to the electromagnetic field of the human beings in the military scenario is still missing.

Methods: a review of the scientific literature regarding the assessment of the exposure of the military personnel to the RF specific to the military environment, was performed.

Results: the review has been performed grouping the scientific literature by the typology of military devices to which the military personnel can be exposed to. The military devices have been classified in four main classes, according to their intended use: communication devices, localization/surveillance devices, jammers and EM directed-energy weapons.

Discussion and Conclusions: The review showed that in the exposure conditions here evaluated, there were only occasional situations of overexposure, whereas in the majority of the conditions the exposure was below the worker exposure limits. Nevertheless, the limited number of studies and the lack of exposure assessment studies for some devices prevent us to draw definitive conclusions and encourage further studies on military exposure assessment.

Conclusions

In conclusion, this study provides an overview of the current used military devices and of the exposure studies that have been performed to characterize the exposure environment in the military scenario with the aim to analyze the state of the art regarding the interaction between the soldier and the RF-EMFs emitted by military devices. The results include four different typologies of systems, each one for different intended use: communication, localization, jamming, and electronic weaponry.

The reviewed studies about the assessment of the human exposure to the radiation emitted by the above mentioned military devices suggest that in the exposure conditions here evaluated, there were only occasional situations of overexposure, but the large number of variables involved in the description of these scenarios, the wide heterogeneity of the exposure conditions and the absence of exposure evaluation for some specific devices make further studies necessary, also considering upcoming use of other technologies, such as 5G, which will dramatically change the exposure conditions also in military environment, both in terms of the time and conditions exposure and in terms of intrinsic sources characteristics, such as the used frequency ranges.

Open access paper: <https://www.mdpi.com/1660-4601/19/2/920/html>

NO ALTERATION BETWEEN INTRINSIC CONNECTIVITY NETWORKS BY A PILOT STUDY ON LOCALIZED EXPOSURE TO THE FOURTH- GENERATION WIRELESS COMMUNICATION SIGNALS (4G)

Yang Lei, Liu Qingmeng, Zhou Yu, Wang Xing, Wu Tongning, Chen Zhiye. No Alteration Between Intrinsic Connectivity Networks by a Pilot Study on Localized Exposure to the Fourth-Generation Wireless Communication Signals. *Frontiers in Public Health*. Vol 9. 2022. doi:10.3389/fpubh.2021.734370.

Abstract

Neurophysiological effect of human exposure to radiofrequency signals has attracted considerable attention, which was claimed to have an association with a series of clinical symptoms. A few investigations have been conducted on alteration of brain functions, yet no known research focused on intrinsic connectivity networks, an attribute that may relate to some behavioral functions. To investigate the exposure effect on functional connectivity between intrinsic connectivity networks, we conducted experiments with seventeen participants experiencing localized head exposure to real and sham time-division long-term evolution signal for 30 min. The resting-state functional magnetic resonance imaging data were collected before and after exposure, respectively. Group-level independent component analysis was used to decompose networks of interest. Three states were clustered, which can reflect different cognitive conditions. Dynamic connectivity as well as conventional connectivity between networks per state were computed and followed by paired sample t-tests. Results showed that there was no statistical difference in static or dynamic functional network connectivity in both real and sham exposure conditions, and pointed out that the impact of short-term electromagnetic exposure was undetected at the ICNs level. The specific brain parcellations and metrics used in the study may lead to different results on brain modulation.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.734370>

COMPARE FABRIC AND BRICK WALL AS SHIELDING MATERIAL DEPENDENCE ON ELECTROMAGNETIC FIELD

M. Pavlík, J. Zbojovský, I. Kolcunová. Compare of fabric and brick wall as shielding material dependence on electromagnetic field. 2021 IEEE 4th International Conference and Workshop Óbuda on Electrical and Power Engineering (CANDO-EPE), 2021, pp. 139-142, doi: 10.1109/CANDO-EPE54223.2021.9667841.

Abstract

The electromagnetic field is a much discussed topic, especially now with the advent of 5G networks. The general public is concerned about the possible impact of the electromagnetic field on human health. By using wireless technologies, we increase the influence of the electromagnetic field on the human body. The basic measures for the protection against electromagnetic radiation are: 1. shielding of the radiation source, 2. shielding of the workplace, 3. personal protective equipment, 4. reduction of radiation in the immediate vicinity of the radiation source. In the household it is possible to shade the “working” environment which is our house. In this paper, we focused on measuring the shielding efficiency of the electromagnetic field. Shielding effectiveness is an indicator of the shielding ability of the material. The measured objects were a brick and a shielding fabric, which can be used practically anywhere. The research focused on the frequency range from 1 GHz to 9 GHz.

The results showed the shielding ability achieved by the brick material and the shielding fabric.

Excerpts

A brick wall with 33cm thickness (Fig. 3) and a shielding fabric (Fig. 2) were placed in the middle between the antennas. The shielding fabric was attached to polystyrene. The shielding efficiency of the polystyrene on which the fabric was attached was also measured during the experiment. The shielding efficiency of the polystyrene was subtracted from the total shielding efficiency of the shielding fabric. The composition of the shielding fabric is approximately 68% cotton, 16% polyester, 16% stainless steel....

The current trend of increasing the sources of the electromagnetic field will probably not stop. We can assume that the number of sources of the electromagnetic field will continue to increase. As mentioned in the introduction, the basic measures for the protection against electromagnetic radiation are: 1. shielding of the radiation source, 2. shielding of the workplace, 3. personal protective equipment, 4. reduction of radiation in the immediate vicinity of the radiation source. However, not all are feasible. At present, however, there are many ways to protect yourself from the effects of the electromagnetic field. The basic protection against the external electromagnetic field is provided by buildings, which prevent the penetration of the electromagnetic field into the interior. However, inside the building, there are many sources of electromagnetic fields that are commonly used. It is therefore possible to additionally protect yourself, for example by using shielding materials which can be commercially purchased. Such a material is, for example, a shielding fabric.

In this paper, we focused on measuring the shielding efficiency, which is an indicator of the shielding ability of the material. The measurement was performed on the basis of [6] IEEE 299–2006 Standard: Method for Measuring the Effectiveness of Electromagnetic Shielding Enclosures, EMC Society. The research focused on the frequency range from 1 GHz to 9 GHz. The shielding ability of both materials was monitored. From the results shown in Fig. 4 we can see the shielding ability of both materials. The brick wall shading efficiency gradually increases in the monitored frequency band up to 5.5 GHz. At higher frequencies, the shielding efficiency remains approximately the same. The shielding efficiency of the shielding fabric does not change much in the monitored frequency range. The results of the research show that the shielding fabric is a more suitable material than brick wall, which could be assumed. However, it must be said that brick wall is a material that is commonly available in the house. The shielding fabric can be used to supplement and increase the shielding ability. Since we need an electromagnetic field for the functionality of, for example, a mobile phone, wifi internet, etc., it would be appropriate to use a shielding fabric when these devices are not in use. For example, at night. In futures we could other materials be used to shield the radiation, for example materials used for thermal and acoustic insulation of buildings.

<https://ieeexplore-ieee-org.libproxy.berkeley.edu/stamp/stamp.jsp?tp=&arnumber=9667841&isnumber=9667546>

EFFECT OF ELECTROMAGNETIC RADIATION ON THE LIVER STRUCTURE AND ULTRASTRUCTURE OF IN UTERO IRRADIATED RATS

Katarína Holovská, Viera Almášiová, Sandra Andrašková, Zuzana Demčišáková, Enikő Račeková, Viera Cigánková. Effect of electromagnetic radiation on the liver structure and ultrastructure of in utero irradiated rats. *Acta Vet. Brno* 2021, 90: 315-319. doi:10.2754/avb202190030315.

The aim of this study was to observe the influence of electromagnetic radiation (EMR) on the structure and ultrastructure of the rat's liver. The pregnant rats used in the experiment were exposed to a pulsed microwave radiation (frequency of 2.45 GHz; mean power density of 2.8 mW/cm²) daily for 2 h, throughout their pregnancy. After delivery, the offspring was not exposed to EMR. Samples of the liver of 5-week-old offspring were subjected to histopathological evaluation. They were processed for light and transmission electron microscopy. Our results indicated that EMR did not cause pronounced changes in the structure of the liver of the investigated offspring. The size and shape of liver lobuli was preserved and the amount of connective tissue in the liver parenchyma did not increase. However, electron microscopy revealed changes in the shape and number of microvilli at the vascular pole of hepatocytes, and formation of vesicles of various shapes and sizes.

The endothelial cells were swollen with larger fenestrations compared to the control group. The spaces of Disse were irregular and dilated. Even though these changes were only mild, further studies are needed to determine the effect of EMR and clarify its potential risk during pregnancy.

Open access paper: https://actavet.vfu.cz/media/pdf/actavet_2021090030315.pdf

POTENTIAL FACTORS AFFECTING CHRONIC CHEMICAL INTOLERANCE ASSOCIATED WITH CONSTITUTIONAL PREDISPOSITION OR LIFESTYLE AND ENVIRONMENT DURING CHILDHOOD: FROM A SIX-YEAR FOLLOW-UP STUDY

Kenichi Azuma, Iwao Uchiyama, Naoki Kunugita. Potential factors affecting chronic chemical intolerance associated with constitutional predisposition or lifestyle and environment during childhood: From a six-year follow-up study. *J Psychosom Res.* 2021 Dec;151:110665. doi: 10.1016/j.jpsychores.2021.110665.

Abstract

Objective: The Japanese chemical intolerance (JCI) study was conducted in January 2012 with a cohort of 7245 adults from population-based sampling. This study aimed to investigate the childhood constitutional and environmental factors involved in the development of chronic CI from the prospective cohort study.

Methods: In the cohort, 4683 persons were identified after six years. Self-reported questionnaires were administered to the subjects to obtain information on CI status; medical history; constitution, lifestyle, and housing environment during childhood; and recent psychosomatic states. We assessed the differences between individuals with persisting CI status during the follow-up (defined as chronic CI) and controls not having CI status during the follow-up.

Results: A total of 2500 individuals responded. Multiple logistic regression analyses revealed significant associations between chronic CI and motion sickness to car or bus and allergic

conjunctivitis during childhood. Significant associations between a possible increased risk of CI and the existence of high-voltage power lines close to housing, the use of vinyl covering in wall material, the use of strong perfume by a family member, and the experience of stinky odor of paint or wax at elementary school were observed. However, the use of carpet in floor material and plaster coating in wall material was associated with a possible decreased risk of CI.

Conclusions: Some potential constitutional predisposition from childhood, including inherent susceptibility in the autonomic nervous system may be involved in the development of CI.

<https://pubmed.ncbi.nlm.nih.gov/34763204/>

ASSOCIATION OF EXCESSIVE MOBILE PHONE USAGE WITH SLEEP QUALITY AND FATIGUE SEVERITY: AN EPIDEMIOLOGIC SURVEY IN SAUDI POPULATION

Al-khlaiwi T, Habib S. Association of Excessive Mobile Phone Usage With Sleep Quality and Fatigue Severity: an Epidemiologic Survey in Saudi Population. KMUJ. 13(2):60-5. 2021. doi: 10.35845/kmuj.2021.20531.

Abstract

OBJECTIVE: To determine the effect of mobile phone usage on sleep quality and fatigue severity through standard questionnaires in Saudi Arabia mobile phone users.

METHODS: This cross-sectional study was conducted in Riyadh, Saudi Arabia on 251 mobile phone users from September 2019 to January 2020. Volunteer mobile phone users aging 15-65 years were selected by nonprobability convenience sampling technique from the Riyadh city community. The subjects were interviewed about the knowledge, attitude and practice of mobile phone use and were assessed through Pittsburg Sleep Quality index (PSQI) and Fatigue severity scale (FSS). Spearman's correlations and Linear regression were applied for statistical analysis through SPSS V.22.

RESULTS: Mean age of study participants was 32.43±12.8 years. There was a significant increase in PSQI and FSS with increase in daily mobile usage. When participants exceeded one-hour daily usage, PSQI (<1h=3.75, 1-2 h=5.78, >2h=6.85, p= 0.001) and FSS (<1h= 2.25, 1-2h= 3.17, >2h= 3.61, p=0.008) exceeded acceptable scores.

Moreover, placing mobile phone inside bedroom was associated with the increase of PSQI (Near Pillow=6.93, Inside Bed=6.16, Outside bed=5.75, p=0.001) and FSS (Near Pillow=3.65, Inside Bed=3.39, Outside Bed=2.58, p= 0.008). There was a significant increase in PSQI (Near Pillow=6.93, Inside Bed=6.16, Outside bed=5.75, p=0.001) and FSS (Near Pillow=3.65, Inside Bed=3.39, Outside Bed=2.58, p= 0.008) its location inside bedrooms. Significant correlation was observed between high PSQI and FSS with daily usage (r=0.213, p=0.001 & r=0.171, p=0.01 respectively).

CONCLUSION: Inappropriate and excessive mobile phone usage is associated with poor sleep quality and excessive fatigue in Saudi Arabian population.

Open access paper: <https://www.kmuj.kmu.edu.pk/article/view/20531>

ELECTROMAGNETIC EXPOSURE OF PERSONNEL INVOLVED IN CARDIAC MRI EXAMINATIONS IN 1.5T, 3T AND 7T SCANNERS

Katarzyna Sklinda, Jolanta Karpowicz, Andrzej Stępniewski. Electromagnetic Exposure of Personnel Involved in Cardiac MRI Examinations in 1.5T, 3T and 7T Scanners. *Int J Environ Res Public Health*. 2021 Dec 22;19(1):76. doi: 10.3390/ijerph19010076.

Abstract

(1) Background: It has been hypothesised that a significant increase in the use of cardiac magnetic resonance (CMR), for example, when examining COVID-19 convalescents using magnetic resonance imaging (MRI), has an influence the exposure profiles of medical personnel to static magnetic fields (STmf).

(2) Methods: Static exposure to STmf (SEmf) was recorded during activities that modelled performing CMR by radiographers. The motion-induced time variability of that exposure (TVEmf) was calculated from SEMf samples. The results were compared with: (i) labour law requirements; (ii) the distribution of vertigo perception probability near MRI magnets; and (iii) the exposure profile when actually performing a head MRI.

(3) Results: The exposure profiles of personnel managing 42 CMR scans (modelled using medium (1.5T), high (3T) and ultrahigh (7T) field scanners) were significantly different than when managing a head MRI. The majority of SEMf and TVEmf samples (up to the 95th percentile) were at low vertigo perception probability (SEmf < 500 mT, TVEmf < 600 mT/s), but a small fraction were at medium/high levels.

(4) Conclusion: Even under the "normal working conditions" defined for SEMf (STmf < 2T) by labour legislation (Directive 2013/35/EC), increased CMR usage increases vertigo-related hazards experienced by MRI personnel (a re-evaluation of electromagnetic safety hazards is suggested in the case of these or similar changes in work organisation).

Open access paper: <https://www.mdpi.com/1660-4601/19/1/76>

NONEQUILIBRIUM, WEAK-FIELD-INDUCED CYCLOTRON MOTION: A MECHANISM FOR MAGNETOBIOLOGY

Ashot Matevosyan, Armen E Allahverdyan. Nonequilibrium, weak-field-induced cyclotron motion: A mechanism for magnetobiology. *Phys Rev E*. 2021 Dec;104(6-1):064407. doi: 10.1103/PhysRevE.104.064407.

Abstract

There is a long-time quest for understanding physical mechanisms of weak magnetic field interaction with biological matter. Two factors impeded the development of such mechanisms: first, a high (room) temperature of a cellular environment, where a weak, static magnetic field induces a (classically) zero equilibrium response. Second, the friction in the cellular environment is large, preventing a weak field to alter nonequilibrium processes such as a free diffusion of charges. Here we study a class of nonequilibrium steady states of a cellular ion in a confining potential, where the response to a (weak, homogeneous, static)

magnetic field survives strong friction and thermal fluctuations. The magnetic field induces a rotational motion of the ion that proceeds with the cyclotron frequency. Such nonequilibrium states are generated by a white noise acting on the ion additionally to the nonlocal (memory-containing) friction and noise generated by an equilibrium thermal bath. The intensity of this white noise can be weak, i.e., much smaller than the thermal noise intensity.

<https://pubmed.ncbi.nlm.nih.gov/35030908/>

HUMAN PSYCHOPHYSIOLOGY IS INFLUENCED BY LOW-LEVEL MAGNETIC FIELDS: SOLAR ACTIVITY AS THE CAUSE

Michael Hanzelka, Jiří Dan, Pavel Fiala, Přemysl Dohnal. Human Psychophysiology Is Influenced by Low-Level Magnetic Fields: Solar Activity as the Cause. *Atmosphere* 12, no. 12: 1600. 2021. doi: 10.3390/atmos12121600

Abstract

We evaluate the impact of changes in solar activity on three significant human psychophysiological parameters: skin conductance, electromyography (EMG), and the share of abdominal and diaphragmatic breathing in overall ventilation. Variations affecting human psychophysiology due to changes in solar activity directly document the assertion that psychology, behavior, and decision-making all reflect geomagnetic field alterations that stem from variable solar activity. The relevant experiments showed that solar processes, during which the Earth is exposed to electrically charged particles from the Sun (solar wind), exert an impact on the psychophysiological parameters of the body.

Open access paper: <https://www.mdpi.com/2073-4433/12/12/1600>

EFFECT OF EXTREMELY LOW-FREQUENCY MAGNETIC FIELDS ON LIGHT-INDUCED ELECTRIC REACTIONS IN WHEAT

Grinberg M, Mudrilov M, Kozlova E, Sukhov V, Sarafanov F, Evtushenko A, Ilin N, Vodeneev V, Price C, Mareev E. Effect of extremely low-frequency magnetic fields on light-induced electric reactions in wheat. *Plant Signal Behav.* 2022 Jan 7:2021664. doi: 10.1080/15592324.2021.2021664.

Abstract

Magnetic field oscillations resulting from atmospheric events could have an effect on growth and development of plants and on the responsive reactions of plants to other environmental factors. In the current work, extremely low-frequency magnetic field (14.3 Hz) was shown to modulate light-induced electric reactions of wheat (*Triticum aestivum* L.). Blue light-induced electric reaction in wheat leaf comprises depolarization and two waves of hyperpolarization resulting in an increase of the potential to a higher level compared to the dark one. Fluorescent and inhibitory analysis demonstrate a key role of calcium ions and calcium-dependent H⁺-ATPase of the plasma membrane in the development of the reaction. Activation of H⁺-ATPase by the increased calcium influx is suggested as a mechanism of the influence of magnetic field on light-induced electric reaction.

<https://pubmed.ncbi.nlm.nih.gov/34994282/>

HEALTH EFFECTS OF 5G BASE STATION EXPOSURE: A SYSTEMATIC REVIEW

Tasneem Sofri, Hasliza A Rahim, Mohamedfareq Abdulmalek, Khatijahhusna Abd Rani, Mohd Hafizi Omar, Mohd Najib Mohd Yasin, Muzammil Jusoh, Ping Jack Soh. Health Effects of 5G Base Station Exposure: A Systematic Review. IEEE Access. Dec 30, 2021. doi: 10.1109/ACCESS.2021.3139385.

Abstract

The Fifth Generation (5G) communication technology will deliver faster data speeds and support numerous new applications such as virtual and augmented reality. The additional need for a larger number of 5G base stations has sparked widespread public concerns about their possible negative health impacts. This review analyzes the latest research on electromagnetic exposure on humans, with particular attention to its effect on cognitive performance, well-being, physiological parameters, and Electroencephalography (EEG). While most of their results indicated no changes in cognitive function, physiological parameters, or overall well-being, the strength of the EEG alpha wave is noticed to vary depending on various aspects of cognitive functions. However, the available studies have not investigated the health effects resulting from exposure from the 5G mobile phone and base station antennas from 700 MHz to 30 GHz on the cognitive performance, well-being subjective symptoms, human physiological parameters, and EEG of adults. There is a need for such research regarding this current emerging technology. Such studies are significant in determining whether 5G technology is indeed safe for humans.

Conclusion

This work presents an analysis of exposure studies conducted using signals from 400 MHz to 1750 MHz (for 4G). From this analysis, the following conclusions are made:

- Most of the studies in literature using 2G/3G/4G showed no effects and no consistency in how exposure to these signals affected the cognitive, physiological parameters, well-being, and EEG of the volunteers.
- Most research on human cognition, physiological parameters, and well-being so far have focused on the impacts of GSM900/GSM1800/UMTS/4G MPs, GSM900/GSM1800/UMTS BSs, DECT, and Wi-Fi exposures.
- There is an absence of studies reporting the effects of 5G (700 MHz, 3.5 GHz, or 28 GHz) BS signals on adults in terms of cognitive performance, well-being, or physiological markers (heart rate, blood pressure, and body temperature).

Figure 9 and 10 illustrated the possible flowchart and schematic diagram to study the effects of 5G BS exposure signals for sub-6 GHz and mmWave bands (of up to 30 GHz) to human subjects. Data from such a study will be useful in explicitly determining the significance signal exposure from 5G BS on human health, considering their much closer proximity to users.

Open access paper: <https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9665755>

TRIGGER OF A MIGRAINE HEADACHE AMONG THAI ADOLESCENTS SMARTPHONE USERS: A TIME SERIES STUDY

Wanna Chongchitpaisan, Phongtape Wiwatanadate, Surat Tanprawate, Assawin Narkpongphan, Nipapon Siripon. Trigger of a migraine headache among Thai adolescents smartphone users: a time series study. *Environ Anal Health Toxicol*. 2021 Mar;36(1):e2021006-0. doi: 10.5620/eaht.2021006.

Abstract

The study aims to investigate the effect of modern technology in the trigger of migraine headaches in high school students by using generalized estimating equation analysis. The prospective time series study was conducted in one hundred and forty-five smartphones using high school students in Chiang Mai Province who each completed a headache diary giving a total of 12,969 data entries. The smartphone output power was measured and recorded by a smartphone application. Smartphone use, sleep quality, anxiety, and depression also were assessed. Results revealed that the prevalence of repeated headache be 13.4% and migraine 16.9%. The migraine had the strongest association with handfree device use and internet use, followed by age and anxiety. Furthermore, the strongest effect of smartphone output power in triggering a migraine was found at $\leq 1.79 \times 10^{-5}$ and $1.80-1.99 \times 10^{-5}$ mW ranges. Meanwhile, Lag_6 of daily smartphone output power exposure produced the migraine effect in a reverse dose-response manner. The smartphone electromagnetic radiation was a primary migraine trigger. The study results led to the recommendation to avoid triggers by use of smartphone talking with hand-free devices to prevent a recurrent migraine.

Conclusions

SOP, which is smartphone electromagnetic radiation, has a non-linear correlation with migraine headaches which has been called a window effect response. The result has also found migraine response to a delayed effect of SOP in the form of reverse dose-response. The delayed effect response on repeated exposure can recover and re-balance itself after exposure to electromagnetic radiation. Additionally, smartphone electromagnetic radiation has effects that fit the criteria for triggers that induce migraines. Finally, younger student, internet use and talking without hand-free devices were risk factors of migraines. It is recommended that limited time for smartphone talking with hand-free device and older age starting using smartphone be suggested in order to prevent migraine attack.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8207005/>

ANSWERS AND QUESTIONS: FORTY YEARS IN BIOELECTROMAGNETICS

Ben Greenebaum. Answers and Questions: Forty Years in Bioelectromagnetics. *Bioelectromagnetics*. 2021 Dec 2. doi: 10.1002/bem.22381.

Abstract

The work began in 1972 when three young assistant professors used a slime mold to see if electromagnetic fields would affect it. The fields did, though the effects were small and hard to tease out of the noise. The cell cycle was lengthened and there were changes in respiration. So, the next question was "how and why?" Further changes were seen using these and then other bacterial and eukaryotic cells in respiration, in ATP, in the protein replication chain, and so forth. Changes occurred even in cell extracts that lacked an intact plasma membrane.

Nerve cells showed changes in leakage of neurotransmitters and in neurite outgrowth from excised ganglia.

Based on some experiments with nerve cells, I also did some computer calculations, modeling the internal electric and magnetic fields and current densities in simplified representations of bone fractures and also of spinal cords in vertebrae. More recently, I have collaborated on some theoretical models of what fields might be doing at the cellular and molecular level, particularly with reference to the radical model. With each piece of research, my collaborators and I generally found a small piece of information about fields and biological systems; and each answer raised another set of questions, which is the way of science. Though bioelectromagnetic scientists have learned much and can say much at greater depth about what happens when an organism is exposed to a field, the fundamental question still remains: What exactly is going on here?

<https://pubmed.ncbi.nlm.nih.gov/34859455/>

WHAT EVIDENCE EXISTS ON THE IMPACT OF ANTHROPOGENIC RADIOFREQUENCY ELECTROMAGNETIC FIELDS ON ANIMALS AND PLANTS IN THE ENVIRONMENT? A SYSTEMATIC MAP PROTOCOL

Ken Karipidis, Chris Brzozek, Chhavi Raj Bhatt, Sarah Loughran, Andrew Wood. What evidence exists on the impact of anthropogenic radiofrequency electromagnetic fields on animals and plants in the environment? A systematic map protocol. *Environmental Evidence*. 10(39). 21 December 2021.

Abstract

Background Exposure to radiofrequency (RF) electromagnetic fields (EMF), particularly from telecommunications sources, is one of the most common and fastest growing anthropogenic factors on the environment. In many countries, humans are protected from excessive RF EMF exposure by safety standards that are based on guidelines by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). The ICNIRP guidelines are based on knowledge of how RF EMF affects the human body, however, there are currently no recognised international guidelines to specifically protect animals and plants. Whether the ICNIRP guidelines for humans is adequate to provide protection to the environment is a subject of active debate. This systematic map will collate all the available evidence on whether anthropogenic RF EMF has a negative effect on plants and animals in the environment. The map will also identify gaps in knowledge, recommend future research and inform environmental and radiation protection authorities.

Methods The proposed systematic map will include peer-reviewed and grey literature published in English. The EMF—Portal, PubMed and Web of Science databases will be searched using a search string prepared by the review team and tested for comprehensiveness against a list of known relevant reviews. Once duplicates are removed, retrieved articles will be screened in three stages: title, abstract, and full text. Studies will be selected with a subject population of all plants and animals, with exposures to anthropogenic RF EMF (frequency range 100 kHz–300 GHz) compared to no or lower-level exposure, and for all outcomes related to the studied populations. Kappa statistic tests will be conducted at each stage to ensure consistency of decision-making regarding the predefined inclusion/exclusion criteria. Eligible studies will then proceed to the data extraction phase,

which will extract meta-data such as bibliographic information, taxonomic information, RF EMF exposure data, outcome(s), sample size, etc. The extracted data will then be organised into a systematic map and the findings summarised by cross-tabulating key meta-data variables in heat maps, charts or other data visualization methods. The systematic map will identify gaps in knowledge, priorities for future research and potential subtopics for further analysis and/or systematic review.

Open access paper:

<https://environmentalevidencejournal.biomedcentral.com/articles/10.1186/s13750-021-00252-w>

CAN THE MICROWAVE AUDITORY EFFECT BE “WEAPONIZED”?

Foster Kenneth R., Garrett David C., Ziskin Marvin C. Can the Microwave Auditory Effect Be “Weaponized”? *Frontiers in Public Health*. 9:2142. 2021. doi:10.3389/fpubh.2021.788613.

No abstract.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.788613>

THE POTENTIAL ADVERSE EFFECT OF 2.45 GHZ MICROWAVE RADIATION ON THE TESTES OF PRENATALLY EXPOSED PERIPUBERTAL MALE RATS

Andrašková S, Holovská K, Ševčíková Z, Andrejčáková Z, Tóth Š, Martončíková M, Račková E, Almášiová V. The potential adverse effect of 2.45 GHz microwave radiation on the testes of prenatally exposed peripubertal male rats. *Histol Histopathol*. 2021 Dec 2:18402. doi: 10.14670/HH-18-402.

Abstract

In utero development of organs is easily influenced by many environmental factors. The aim of this study was to elucidate the effect of microwave radiation (MR) at a frequency of 2.45 GHz and a specific absorption rate of 1.73 W/kg on intrauterine development of testis. Pregnant albino rats were exposed to whole-body MR for 2 hours per day throughout the pregnancy. Male offspring (n=12, age 35 days) were not exposed to MR after birth. The study revealed that MR applied in utero induced apparent structural changes in the testes, such as irregular shape of seminiferous tubules, significant decrease in the diameter of seminiferous tubules ($p<0.05$) and in the height of the germinal epithelium ($p<0.01$), disorganisation of germ cells, desquamations of immature germ cells, formation of giant multinucleated cells, and significant ($p<0.01$) expansion of the interstitium. At the level of transmission electron microscopy, there were observed basement membrane irregularities in seminiferous tubules, vacuolation of the cytoplasm and adversely affected organelles in Sertoli cells, germ cells, Leydig cells, peritubular and endothelial cells. The tight junctions between adjacent Sertoli cells were often incomplete, and necrotizing germ cells were more numerous in experimental animals compared to controls. Enhanced necrotizations of germ cells proved by a Fluoro Jade C method, and declined germ cells proliferation confirmed by proliferating cell nuclear antigen analysis, were detected in MR exposed animals. Our results revealed that the prenatal exposure to MR had an adverse effect on the postnatal testicular development in rats.

<https://pubmed.ncbi.nlm.nih.gov/34854072/>

Excerpts

The exact cellular pathways mediating the adverse effect of RF-EMR on testicular development and future spermatogenesis are still under investigation and further studies are still required. Many authors suggested that the most likely mechanism of RF-EMR associated testicular tissue injury is oxidative damage (Valko et al., 2006; Höytö et al., 2008; Desai et al., 2009; Tök et al. 2014; Yakymenko et al. 2016). This is in close agreement with our observation of the impaired testicular structure, ultrastructure, and regression of proliferative capacity of spermatogenic elements, accompanied by enhanced degeneration. Given the chosen methodology and the calculated SAR value, we have assumed that the primary source of testicular tissue damage was the non-thermal effect of RF-EMR. However, due to the impossibility of monitoring the hot spot phenomenon in the developing testes of rat foetuses, we cannot exclude the thermal effect.

In conclusion, our data attest to the harmful impact of 2.45 GHz microwave radiation on development of testes during the prenatal period which subsequently compromises the spermatogenic cycle and male reproductive function in the postnatal period of life. To preserve the steady reproductive health, it would be necessary to avoid or minimize exposure to RF-MR, especially in the most vulnerable period of intrauterine development of the individual.

THE EFFECTS OF GUM ARABIC, CURCUMIN (*CURCUMA LONGA*) AND GARCINIA KOLA ON THE RAT HIPPOCAMPUS AFTER ELECTROMAGNETIC FIELD EXPOSURE

Abdalla Ahmed Eldaw Elamin, Omur Gulsum Deniz, Suleyman Kaplan. The Effects of Gum arabic, Curcumin (*Curcuma longa*) and Garcinia kola on the Rat Hippocampus after Electromagnetic Field Exposure: A Stereological and Histological Study. *J Chem Neuroanat.* 2021 Dec 13;102060. doi: 10.1016/j.jchemneu.2021.102060.

Abstract

The present study was designed to focus on the potential effects of the electromagnetic field (EMF) emitted by mobile phones on hippocampal pyramidal neurons and to investigate the role of curcumin (Cur), Garcinia kola (GK) and Gum arabic (GA) in reducing these adverse effects. Fifty-four 12-week-old male Wistar albino rats were used. These were randomly divided into nine groups of six rats each. The control, Cur, GK and GA groups were not exposed to EMF, while the sham group was kept in the EMF exposure system without being exposed to EMF. The EMF+Cur, EMF+GK, EMF+GA and EMF groups were exposed to 900MHz EMF for one hour a day for 28 days. The number of the pyramidal neurons in the cornu ammonis (CA) of the hippocampus was estimated using the optical fractionator technique. Histopathological changes were evaluated under light and electron microscopes. The activities of the superoxide dismutase (SOD) and catalase (CAT) enzymes were also evaluated from serum samples. Significant levels of CAT and SOD activities were observed in the EMF group compared to the control group ($p=0.000$; $p=0.001$) respectively. Microscopic observations showed that dark-coloured nuclei with unclear neuron boundaries were frequently observed in the EMF group. Stereological data analysis revealed a significant decrease in the CA's total number of pyramidal neurons in the EMF group compared to the control and sham groups ($p=0.000$; $p=0.000$) respectively. Cur and GK were observed to provide significant protection in the EMF+Cur and EMF+GK groups compared to the EMF group ($p=0.000$; $p=0.000$) respectively. No significant difference was observed between the EMF+GA group and EMF

group ($p=0.989$). Exposure to 900MHz EMF causes severe alterations in the number and structure of hippocampal pyramidal neurons. Cur and GK exhibit a protective effect against these deleterious effects, but GA showed no protective effect.
<https://pubmed.ncbi.nlm.nih.gov/34915150/>

EXPOSURE OF THE SCROTUM TO 4G CELLPHONE RADIOFREQUENCY RADIATION AFFECTS SPERMATOGENESIS AND BLOOD-TESTIS BARRIER OF ADULT MALE RATS

Qing Cheng, Gang Yu, Gang Wang, Zhi-Ming Bai. [Zhonghua Nan Ke Xue. 2021 Aug;27(8):680-684.

Abstract

Objective: To investigate the direct impact of exposure of the scrotum to 4G cellphone radiofrequency electromagnetic radiation (RF-EMR) on the testis of adult rats.

Methods: We equally randomized 30 adult male SD rats into a control and an exposure group, the latter exposed to 4G cellphone RF-EMR 6 hours a day for 150 days. Then, we analyzed the changes in semen quality and seminiferous epithelia, and measured the levels antioxidant enzyme-peroxide and the expressions of blood- testis barrier proteins in the testis of the rats.

Results: Compared with the controls, the rats in the exposure group showed significantly decreased sperm concentration ($[6.39 \pm 0.82]$ vs $[4.74 \pm 0.87]$ $\times 10^7/\text{ml}$, $P < 0.05$), viability (62.11 ± 8.82 % vs $[41.44 \pm 7.33]$ %, $P < 0.05$), motility ($[55.71 \pm 7.39]$ vs $[36.22 \pm 6.36]$ %, $P < 0.05$), the percentage of morphologically normal sperm ($[84.89 \pm 5.11]$ % vs $[70.78 \pm 8.11]$ %, $P < 0.05$) and Johnsen's score (8.38 ± 0.98 vs 6.11 ± 1.56 , $P < 0.05$), increased Coentino's score (1.36 ± 0.21 vs 1.81 ± 0.34 , $P < 0.05$) and MDA level ($P < 0.05$), reduced levels of SOD, GSH and CAT ($P < 0.05$), and down-regulated expressions of occludin, ZO-1, CAR and N-Cadherin ($P < 0.05$).

Conclusions: Exposure of the scrotum to 4G cellphone RF-EMR directly causes injury of the testis, disorder of the blood-testis barrier, reduction of semen quality and consequently decline in the fertility of male rats.

<https://pubmed.ncbi.nlm.nih.gov/34914238/>

3.5-GHZ RADIOFREQUENCY ELECTROMAGNETIC RADIATION PROMOTES THE DEVELOPMENT OF DROSOPHILA MELANOGASTER

Yahong Wang, Zhihao Jiang, Lu Zhang, Ziyang Zhang, Yanyan Liao, Peng Cai. Environ Pollut. 2021 Dec 8;294:118646. doi: 10.1016/j.envpol.2021.118646.

Highlights

- Simulating the most representative 3.5 GHz RF-EMF band of 5G in China.
- We study the long-term effects of 3.5 GHz RF-EMF on flies' development.
- The microbial community may participate in the development under long-term RF-EMF.
- It provides a reference for the hidden health effects of RF-EMF.

Abstract

With the rapidly increasing popularity of 5G mobile technology, the effect of radiofrequency radiation on human health has caused public concern. This study explores the effects of a simulated 3.5 GHz radiofrequency electromagnetic radiation (RF-EMF) environment on the development and microbiome of flies under intensities of 0.1 W/m², 1 W/m² and 10 W/m². We found that the pupation percentages in the first 3 days and eclosion rate in the first 2 days were increased under exposure to RF-EMF, and the mean development time was shortened. In a study on third-instar larvae, the expression levels of the heat shock protein genes hsp22, hsp26 and hsp70 and humoral immune system genes AttC, TotC and TotA were all significantly increased. In the oxidative stress system, DuoX gene expression was decreased, sod2 and cat gene expression levels were increased, and SOD and CAT enzyme activity also showed a significant increase. According to the 16S rDNA results, the diversity and species abundance of the microbial community decreased significantly, and according to the functional prediction analysis, the genera Acetobacter and Lactobacillus were significantly increased. In conclusion, 3.5 GHz RF-EMF may enhance thermal stress, oxidative stress and humoral immunity, cause changes in the microbial community, and regulate the insulin/TOR and ecdysteroid signalling pathways to promote fly development.
<https://pubmed.ncbi.nlm.nih.gov/34896224/>

ABNORMAL EXPRESSION OF CONNEXIN43 IN CARDIAC INJURY INDUCED BY S-BAND AND X-BAND MICROWAVE EXPOSURE IN RATS

Yin Y, Xu X, Gao Y, Wang J, Yao B, Zhao L, Wang H, Wang H, Dong J, Zhang J, Peng R. Abnormal Expression of Connexin43 in Cardiac Injury Induced by S-Band and X-Band Microwave Exposure in Rats. *J Immunol Res*. 2021 Dec 17;2021:3985697. doi: 10.1155/2021/3985697.

Abstract

Although the effects of microwave exposure on the heart have gradually become the focus of domestic and foreign scholars, the biological effects caused by different doses and different frequency bands of exposure are still unclear. In this study, we will investigate the damaging effect of S-band and X-band microwave composite exposure on cardiac structure and function, as well as the pathophysiological significance of Cx43 in cardiac conduction dysfunction after exposure. We used S- and X-band radiation sources with the average power density of 5 and 10 mW/cm² to expose Wistar rats to single or composite exposure. At the 6th hour, on the 7th, 14th, and 28th days after exposure, ECG was used to detect the electrical conduction of the heart, and the myocardial enzyme was measured by the automatic biochemical analyzer. We selected the observation time points and groups with severe damage to observe the changes of myocardial structure and ultrastructure with an optical microscope and TEM; and to detect the expression and distribution of Cx43 by western blotting and immunohistochemistry. After exposure, the heart rate increased, the P wave amplitude decreased, and the R wave amplitude increased; the content of the myocardial enzyme in serum increased; the structure and ultrastructure of cardiac tissue were damaged. The damage was dose-dependent and frequency-dependent. The expression of Cx43 in myocardial tissue decreased, and distribution was abnormal. Taken together, these findings suggested that the mechanism of abnormal electrical conduction in the heart of rats by S- and X-band microwave exposure might be related to the decreased expression and disordered distribution of Cx43 after microwave exposure.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8709747/>

METHODOLOGY OF STUDYING EFFECTS OF MOBILE PHONE RADIATION ON ORGANISMS: TECHNICAL ASPECTS

Katerina Bartosova, Marek Neruda, Lukas Vojtech. Methodology of Studying Effects of Mobile Phone Radiation on Organisms: Technical Aspects. *Int J Environ Res Public Health*. 2021 Nov 30;18(23):12642. doi: 10.3390/ijerph182312642

Abstract

The negative influence of non-ionizing electromagnetic radiation on organisms, including humans, has been discussed widely in recent years. This paper deals with the methodology of examining possible harmful effects of mobile phone radiation, focusing on in vivo and in vitro laboratory methods of investigation and evaluation and their main problems and difficulties. Basic experimental parameters are summarized and discussed, and recent large studies are also mentioned. For the laboratory experiments, accurate setting and description of dosimetry are essential; therefore, we give recommendations for the technical parameters of the experiments, especially for a well-defined source of radiation by Software Defined Radio.

Excerpt

Moreover, we emphasize that the dosimetry should always be clearly set and observed in the laboratory tests. This includes using a radiofrequency radiation source where the frequency, time, and intensity of the radiation, including placement of the radiation source, can be precisely set and controlled. To comply with the above-mentioned criteria—precisely defined radiofrequency radiation on the one hand, and unpredictable changes of the signal on the other hand—we suggest using the Software Defined Radio (SDR) approach available for 2G, 3G [44,45], 4G, and 5G [46] instead of a signal vector generator or undefined cell phone. The SDR transceiver enables the generation of high-frequency multichannel/wideband power signals in repeatable scenarios (recorded signal or artificial signal), with respect to timing, modulation methods, waveform, transmission power, and its time changes, etc., of real mobile phone radio channel parameters and for different xG mobile generations, unlike a standard signal generator. In order to be able to repeat the experiments in a controlled manner, it is necessary to use a standardized antenna adapter for radiating high-frequency power, i.e., an antenna structure for near/far EM field. The SDR enables the creation of a new base transceiver station (2G), Node B (3G), eNode B (4G), or gNode B (5G), which directly communicates with mobile phones, i.e., it can generate a real unpredictable mobile phone signal (i.e., signal changing in an irregular, unpredictable way), as suggested by Panagopoulos [24]. In addition, the SDR also enables the creation of a mobile phone phantom, i.e., specific hardware solution, e.g., OsmocomBB [44], with a well-defined mobile phone phantom (antenna, controlled radiofrequency signal radiation setup, etc., in standardized measurement environment, i.e., measurement cell), which can facilitate the reproducibility of future experiments. All necessary parameters of individual mobile technologies, including time intervals, frequency, modulation, intensity, etc., can be precisely controlled. Using the SDR and standardized measurement cell could substantially help to avoid problems of the replication of studies. In addition, one of the advantages of this approach is the possibility of sharing a specific scenario between laboratories in order to ensure the repeatability of the radiation source.

Conclusions

The paper reviews laboratory methods studying the effects of radiofrequency radiation on organisms, focusing on technical aspects of the experiments. As can be concluded from many works, the effects of radiofrequency radiation can differ under different conditions and settings. Therefore, we emphasize that the methodology should be clearly and precisely set to ensure the results can be verified and reproduced. Very important is the choice of a suitable source of radiation, here we recommend the SDR, which can simulate a real mobile phone signal in a controlled and repeatable way. The influence of radiofrequency radiation is a multidisciplinary topic and includes many fields, such as medicine, biology, toxicology, physics, electrical and electronic engineering, telecommunication, and statistics. The future research requires close cooperation of scientists from all these fields. Following the given recommendations should increase the overall quality of the experiments and publications and give the possibility to compare, reproduce, and verify the results.

Open access paper: <https://www.mdpi.com/1660-4601/18/23/12642>

MULTI-HOP D2D COMMUNICATION IN CELLULAR NETWORKS TO MINIMIZE EMR

Avirup Das, Nabanita Das, Abhirup Das Barman. Multi-hop D2D Communication in Cellular Networks to Minimize EMR. IEEE Transactions on Green Communications and Networking. 30 December 2021. doi: 10.1109/TGCN.2021.3139286.

Abstract

With the advent of IoT, and the growing demand of infotainment, the number of data sources transmitting at high data rates for long duration is increasing rapidly. It results an ever increasing level of associated Electromagnetic Radiation (EMR) in our surroundings. The transmission powers from the near-field sources like mobile phones along with the duration of sessions are the crucial factors to accurately assess the harmful effects of EMR exposure from these devices. In order to limit and balance the EMR exposure from these sources, this paper, proposes short-range multi-hop packet forwarding technique with cognitive channel allocation for newer technologies of LTE. Theoretical analysis presented here, establishes bounds on the permissible channels and matching data rates to be used by the nodes to balance the duration of transmission maintaining the safe limit of Specific Absorption Rate (SAR). Based on this strategy, algorithm is developed for multi-hop packet routing with channel selection using cognitive radio. Extensive simulation studies on 5G New Radio (NR) band show significant improvement in energy efficiency with nearly balanced transmission duration and SAR reducing the near-field heating effect throughout the network.

<https://ieeexplore.ieee.org/document/9666462>

Effect of Magnetic Fields during Pregnancy on Fetal Growth and Preterm Labor: A Review Article

Zahra Khojastehfard, Somayeh Ghahremani, Maryam Sabaghian, Samaneh Ghalebizadeh, Zoleykha Asgarlou, Soheil Mohammadi Yazdi, Mahla Salarfard. Effect of Magnetic Fields during Pregnancy on Fetal Growth and Preterm Labor: A Review Article. Intl J Pediatrics. 10.22038/ijp.2020.48094.3874

Abstract

Background Low birth weight is one of the most important health indicators in evaluating pregnancy care worldwide. We aimed to evaluate the effect of cell phones used in pregnancy on fetal growth and preterm labor.

Materials and Methods The search process included screening the relevant articles on electronic databases of Scopus, EMBASE, Cochrane, Web of Science, and Medline to analyze the association of maternal exposure to magnetic fields with fetal growth, birth weight, small-for-gestational-age (SGA), and preterm labor.

Results In the first study, the duration of pregnancy was shorter for mothers using cell phones for more than one hour per day than those use shorter than one hour. In the second study, the intermediate use of cell phones by mothers indicated an increased risk of preterm birth. In the third study, the birth weight were lower in excessive cell phone users than normal status. In the fourth study, low, medium, and high cell phone exposure during the six months before conception had no effect on low birth weight, preterm birth, and SGA. In the fifth study, a negative correlation was observed between magnetic fields expose during pregnancy and the birth week, birth weight, and birth length of the newborn. In the sixth study, the risk of low birth weight, preterm birth, and SGA was close to null. In the seventh study, dental radiography during pregnancy was associated with low birth weight, specifically with term low birth weight.

Conclusion In general, there are contradictory findings on the impact of cell phone exposure during pregnancy. These results should be interpreted with caution because of the small sample size and recall bias.

Open access paper:

https://ijp.mums.ac.ir/article_15508_d1291b56392c9b107f6f008380ca99d1.pdf

MOTOR NEURON DISEASE RISK AND MAGNETIC FIELD EXPOSURES

Tom Sorahan, Linda Nichols. Motor neuron disease risk and magnetic field exposures. Occup Med (Lond). 2021 Dec 23; kqab180. doi: 10.1093/occmed/kqab180.

Abstract

Background: Many studies have investigated magnetic field exposure and the risks of motor neuron disease (MND). Meta-analyses have found positive associations but a causal relationship has not been established.

Aims: To investigate the risks of MND and occupational exposure to magnetic fields in a large UK cohort.

Methods: Mortality of 37 986 employees of the former Central Electricity Generating Board of England and Wales was investigated for the period 1987-2018. Employees were first employed in the period 1942-82 and were still in employment on the 1 November, 1987. Detailed calculations enabled estimates to be made of magnetic field exposures. Observed deaths were compared with expected numbers based on mortality rates for the general population of England and Wales and Poisson regression was used to calculate rate ratios (relative risks) for categories of lifetime, lagged (distant) and lagged (recent) magnetic field exposure.

Results: Mortality from MND in the total cohort was similar to national rates (observed 69, expected 71.3, SMR 97, 95% CI 76-122). There were no statistically significant trends of risks increasing with lifetime, recent or distant magnetic field exposure, although positive associations were observed for some categories of recent exposure.

Conclusions: The study did not find that the cohort had elevated risks of MND as a consequence of occupational lifetime exposure to magnetic fields, although a possible role for recent exposures could usefully be investigated in other datasets.

Funding: This work was supported by the Energy Networks Association (ENA). [The Energy Networks Association is the industry body funded by UK gas and electricity transmission and distribution licence holders.]

<https://pubmed.ncbi.nlm.nih.gov/34940878/>

INFLUENCE OF GEOMAGNETIC DISTURBANCES AT DIFFERENT TIMES OF DAY ON LOCOMOTOR ACTIVITY IN ZEBRAFISH (DANIO RERIO)

Viacheslav V Krylov. Influence of Geomagnetic Disturbances at Different Times of Day on Locomotor Activity in Zebrafish (Danio Rerio). *Clocks Sleep*. 2021 Nov 29;3(4):624-632. doi: 10.3390/clockssleep3040045.

Abstract

The influence of magnetic fields and natural geomagnetic storms on biological circadian rhythms are actively studied. This study reveals an impact of local natural perturbations in the geomagnetic field that occurred at different times of the day on circadian patterns of locomotor activity of zebrafish. A decrease in zebrafish swimming speed was observed during the geomagnetic disturbances before or after the fluctuations of diurnal geomagnetic variation. However, if the geomagnetic perturbations coincided with the fluctuations of diurnal geomagnetic variation, the decrease in zebrafish swimming speed was insignificant. This result suggests that the biological effects of geomagnetic disturbances may depend on synchronization with the diurnal geomagnetic variation. It implies that the previously published correlations between geomagnetic activity and medical or biological parameters could result from a disruption in circadian biorhythms.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8700499/>

EFFECT OF ELECTROMAGNETIC FIELD ON ABORTION: A SYSTEMATIC REVIEW AND META-ANALYSIS

Masumeh Ghazanfarpour, Zahra Atarodi Kashani, Reza Pakzad, Fatemeh Abdi, Fatemeh Alsadat Rahnemaei, Poursan Akhavan Akbari, Nasibeh Roozbeh. Effect of electromagnetic field on abortion: A systematic review and meta-analysis. Open Medicine. doi: 10.1515/med-2021-0384

Abstract

Background The increasing use of new technologies by pregnant women inevitably exposes them to the risks of the electromagnetic fields (EMFs). According to the World Health Organization, EMFs are the major sources of pollutants which harm human health. This study was aimed to evaluate the effects of EMF exposure on abortion.

Methods Web of Science, Cochrane Library, MEDLINE, PubMed, EMBASE, Scopus, and Google Scholar were searched until 2021. Pooled odds ratio (OR) with 95% confidence interval (CI) was estimated using a random-effects model. Heterogeneity was explored using Cochran's Q test and I² index. A meta-regression method was employed to investigate the factors affecting heterogeneity between the studies. The Newcastle-Ottawa scale was used to assess the credibility of the studies.

Results Eligible studies (N = 17) were analyzed with a total of 57,693 participants. The mean maternal age (95% CI) was 31.06 years (27.32–34.80). Based on meta-analysis results, the pooled estimate for OR of EMF with its effects was 1.27 (95% CI: 1.10–1.46). According to the results of meta-regression, sample size had a significant effect on heterogeneity between studies (p: 0.030), but mother's age and publication year had no significant effect on heterogeneity (p-value of both were >0.05). No publication bias was observed.

Conclusion Exposure to EMFs above 50 Hz or 16 mG is associated with 1.27× increased risk of abortion. It may be prudent to advise women against this potentially important environmental hazard. Indeed, pregnant women should receive tailored counselling.

Open access paper:

<https://www.degruyter.com/document/doi/10.1515/med-2021-0384/html>

ESTIMATED ALL-DAY AND EVENING WHOLE-BRAIN RADIOFREQUENCY ELECTROMAGNETIC FIELDS DOSES, AND SLEEP IN PREADOLESCENTS

Cabré-Riera A, van Wel L, Liorni I, Koopman-Verhoeff ME, Imaz L, Ibarluzea J, Huss A, Wiart J, Vermeulen R, Joseph W, Capstick M, Vrijheid M, Cardis E, Röösli M, Eeftens M, Thielens A, Tiemeier H, Guxens M. Environ Res. 2021 Oct 29;112291. doi: 10.1016/j.envres.2021.112291.

Highlights

- We estimated whole-brain RF-EMF doses (mJ/kg/day) for several RF-EMF sources.
- We estimated all-day and evening whole-brain RF-EMF doses.
- We assessed subjective and objective sleep measures in preadolescents.
- No association between all-day whole-brain RF-EMF doses and sleep.
- Evening whole-brain RF-EMF dose from phone calls related to shorter total sleep time.

Abstract

Objective: To investigate the association of estimated all-day and evening whole-brain radiofrequency electromagnetic field (RF-EMF) doses with sleep disturbances and objective sleep measures in preadolescents.

Methods: We included preadolescents aged 9-12 years from two population-based birth cohorts, the Dutch Generation R Study (n = 974) and the Spanish INfancia y Medio Ambiente Project (n = 868). All-day and evening overall whole-brain RF-EMF doses (mJ/kg/day) were estimated for several RF-EMF sources including mobile and Digital Enhanced Cordless Telecommunications (DECT) phone calls (named phone calls), other mobile phone uses, tablet use, laptop use (named screen activities), and far-field sources. We also estimated all-day and evening whole-brain RF-EMF doses in these three groups separately (i.e. phone calls, screen activities, and far-field). The Sleep Disturbance Scale for Children was completed by mothers to assess sleep disturbances. Wrist accelerometers together with sleep diaries were used to measure sleep characteristics objectively for 7 consecutive days.

Results: All-day whole-brain RF-EMF doses were not associated with self-reported sleep disturbances and objective sleep measures. Regarding evening doses, preadolescents with high evening whole-brain RF-EMF dose from phone calls had a shorter total sleep time compared to preadolescents with zero evening whole-brain RF-EMF dose from phone calls [-11.9 min (95%CI -21.2; -2.5)].

Conclusions: Our findings suggest the evening as a potentially relevant window of RF-EMF exposure for sleep. However, we cannot exclude that observed associations are due to the activities or reasons motivating the phone calls rather than the RF-EMF exposure itself or due to chance finding. <https://pubmed.ncbi.nlm.nih.gov/34757029/>

Excerpts

Preadolescents spend 48.9 min/day using mobile communication devices for screen activities and 2.5 min/day making phone calls (Supplementary Table S3). The median of the overall estimated all-day whole-brain RF-EMF dose was 60 (interquartile range (IQR) 20; 118) mJ/kg/day and the main contributor to the all-day whole-brain RF-EMF dose were phone calls (78%) (Table 2)

Preadolescents who spent more time with console/computer gaming or television watching were more likely to have higher evening whole-brain RF-EMF dose from screen activities.... Objective total sleep time was on average 7.5 h, sleep efficiency was 84%, and wake after sleep onset was 71.4 min in the study sample (Table 3)...

Preadolescents with high evening whole-brain RF-EMF dose from phone calls had a shorter total sleep time and longer sleep latency compared to preadolescents with zero evening whole-brain RF-EMF dose from phone calls [-11.9 min (95%CI -21.2; -2.5) and) 0.3 min (95%CI 0.0; 0.7), respectively] (Table 6). However, the latter association did not survive correction for multiple testing.

Overall evening whole-brain RF-EMF dose, and evening dose from screen activities were not associated with objective sleep measures (Table 6) and with sleep quality and restfulness (data not shown)....

This study investigated the association of overall and source-specific all-day and evening whole-brain RF-EMF doses with sleep parameters in preadolescents. We found that the all-day whole-brain RF-EMF doses were not associated with sleep disturbances and objective sleep measures. Regarding evening doses, preadolescents with a high evening whole-brain RF-EMF dose from phone calls had shorter total sleep time....

All-day and evening whole-brain RF-EMF dose from screen activities were not associated with subjective or objective sleep measures....

A more comprehensive study needs to be designed to disentangle between all the potential factors related to the use of mobile communication devices that might affect sleep, including RF-EMF exposure. Also, of note, preadolescents are reducing the use of phones for calling and those who make phone calls use hands-free devices or mobile phone applications that allow voice or video calls. This changing pattern might decrease the overall RF-EMF dose the brain receives but increase the amount of RF-EMF dose from screen activities with mobile communication devices....

The levels of RF-EMF exposure from far-field sources are low and do not produce peak and high intensity exposures to the brain such as those from personal use of mobile communication devices for phone calls or screen activities (Birks et al., 2018, 2021). In our study, RF-EMF dose from far-field sources was not related to subjective or objective sleep measures. Previous studies assessing the association between RF-EMF exposure from far-field sources and sleep showed mixed results....

Conclusion

Overall all-day whole-brain RF-EMF dose and all-day dose from phone calls were not associated with sleep, though evening whole-brain RF-EMF dose from phone calls were associated with less favourable sleep characteristics as objectively measured by actigraphy. These findings suggest the evening as a potentially relevant window of exposure. Since this is the first study investigating the association between RF-EMF dose and sleep and there is not known biological mechanism explaining the observed associations, our results should be interpreted with caution. Studies exploring the relationship of RF-EMF exposure to the brain and sleep should assess the amount of RF-EMF dose absorbed by the brain in the evening or at night which might be more relevant for adolescents' sleep.

ORIGINAL FINDINGS CONFIRMED IN REPLICATION STUDY: PROVOCATION WITH 2.4 GHZ CORDLESS PHONE AFFECTS AUTONOMIC NERVOUS SYSTEM AS MEASURED BY HEART RATE VARIABILITY

Magda Havas, Jeffrey Marrongelle. Original Findings Confirmed in Replication Study: Provocation with 2.4 GHz Cordless Phone affects the Autonomic Nervous System (ANS) as measured by Heart Rate Variability (HRV). Medical Research Archives. 9(11). Nov. 2021. doi: 10.18103/mra.v9i11.2605.

Abstract

This is a double-blind, placebo-controlled replication of a study that we previously conducted in Colorado with 25 subjects designed to test the effect of radio frequency radiation (RFR) generated by the base station of a cordless phone on heart rate variability (HRV). In this study, we analyzed the response of 69 subjects between the ages of 26 and 80 in both Canada and the USA. Subjects were exposed to radiation for 3-min intervals generated by a 2.4-GHz cordless phone base station (3–8 microW/cm²). Prior to provocation we conducted an orthostatic test to assess the state of adrenal exhaustion, which interferes with a person's ability to mount a response to a stressor. A few participants had a severe reaction to the RFR with an increase in heart rate and altered HRV indicative of an alarm response to stress. Based on the HRV analyses of the 69 subjects, 7% were classified as being "moderately to very sensitive", 29% were "little to moderately sensitive", 30% were "not to a little sensitive" and 6% were "unknown". These results are not psychosomatic and are not due to electromagnetic interference. Twenty-five percent of the subjects' self-proclaimed sensitivity corresponded to that based on the HRV analysis, while 32% overestimated their sensitivity and 42% did not know whether or not they were electrically sensitive. Of the 39 participants who claimed to experience some electrical hypersensitivity, 36% claimed they also reacted to a cordless phone and experienced heart symptoms and, of these, 64% were classified as having some degree of electrohypersensitivity (EHS) based on their HRV response. Novel findings include documentation of a delayed response to radiation. This protocol underestimates the reaction to electromagnetic radiation and may provide a false negative for those with a delayed reaction and/or with adrenal exhaustion. Orthostatic HRV testing combined with provocation testing may provide a useful diagnostic tool for some sufferers of EHS when they are exposed to electromagnetic radiation. It can be used to confirm EHS but not to reject EHS as a diagnosis since not everyone with EHS has an ANS reaction to electromagnetic radiation.

Conclusions

Our results show that 36% of the individuals tested reacted via altered heart rate variability (HRV) to the non- ionizing radiation generated by a cordless phone base station in this double-blind, placebo-controlled study. These reactions were not psychosomatic. In this study, we document an increased heart rate (HR), altered HRV and changes in the sympathetic and parasympathetic control of the autonomic nervous system (ANS) typical of a fight-or-flight stress response. These results are similar to our previous study. The results are not due to electromagnetic interference (EMI), since we have examples of a delayed response after the radiation was turned off and have tested EMI with much higher exposure using the same technology with no reactions noted. Our results demonstrate that the radiation from a 2.4-GHz cordless phone affects the ANS and may put some individuals with pre-existing heart conditions at risk when exposed to electromagnetic frequencies to which they

are sensitive. Individuals fell into 3 categories: those who had a healthy ANS and were able to tolerate the stress without reactions; those who reacted; and those who had a compromised ANS with adrenal exhaustion and were unable to mount a response leading, in some cases, to a false negative result. Although documenting a response is relatively simple, determining the degree of EHS is quite complex and requires further study especially for those with a compromised ANS.

Open access paper: <https://esmed.org/MRA/mra/article/view/2605>

MICRONUCLEUS ASSAY IN CELL PHONE USERS: IMPORTANCE OF ORAL MUCOSA SCREENING

Melika Ghandehari, Donia Sadri, Sareh Farhadi. Micronucleus Assay in Cell Phone Users: Importance of Oral Mucosa Screening. *Int J Prev Med.* 2021 Sep 29;12:125. doi: 10.4103/ijpvm.IJPVM_489_19.

Abstract

Background: One of the concerns of cell phone users is prolonged exposure to harmful and potentially carcinogenic waves. This study was aimed to investigate the correlation between amount of cell phone use and related factors with percentage of micronucleus containing cells.

Methods: This descriptive study was conducted on selected patients referring to Islamic Azad University Faculty of Dentistry using cell phones regarding related inclusion and exclusion criteria. Papanicolaou staining method was approached for mucosal smears of samples and frequency of micronucleus containing cells and also, frequency of micronucleus in each cell were recorded for each sample; then, correlation of these findings with amount of daily cell phone usage was statistically analyzed using the calculation of Pearson correlation coefficient and preparation of regression analysis (backward) with significant level of lower than 0.05.

Results: Of 100 samples, the frequency of micronucleus containing cells was $2.94\% \pm 1.89\%$ and the frequency of micronucleus in each cell was $1.02\% \pm 1.68\%$. The amount of cell phone usage was significantly correlated with the frequency of micronucleus containing cells ($r = 0.70$, $P = 0.0001$) and also with the frequency of micronucleus in each cell ($r = 0.57$, $P = 0.0001$). Also, age and sex were not significantly correlated with the frequency of micronucleus containing cells ($P = 0.47$ and 0.32) and also with the frequency of micronucleus in each cell, respectively ($P = 0.16$ and 0.27).

Conclusions: The present study showed that the increased amount of cell phone usage had a strong and significant correlation with the higher frequency of the micronucleus containing cells and the higher frequency of micronucleus in each cell in the buccal mucosa. Also, the related factors as age and sex were not significantly correlated with the frequency of micronucleus containing buccal mucosa cells.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8551792/>

THE BIOLOGICAL EFFECTS OF ELECTROMAGNETIC EXPOSURE ON IMMUNE CELLS AND POTENTIAL MECHANISMS

Chuanfu Yao, Li Zhao, Ruiyun Peng. Electromagn Biol Med. 2021 Nov 9;1-10. doi: 10.1080/15368378.2021.2001651.

Abstract

Anxiety about potential health hazards of electromagnetic exposure has been growing in the past decades, with their widely application in many fields. The immune system plays pivotal role in maintaining body's homeostasis. Importantly, immune system is also a sensitive target for electromagnetic fields. In recent years, the biological effects of electromagnetic fields on immune cells have been attracting more and more attentions. Accumulated data suggested that electromagnetic exposure could affect the number and function of immune cells to some extent, including cell proportion, cell cycle, apoptosis, killing activity, cytokines contents and so on. The research objects basically covered all types of immune cells, mainly on PBMC, T lymphocytes, B lymphocytes, NK cells and macrophages. Meanwhile, there also are negative reports of electromagnetic fields on immune cells. This article reviews the results of epidemiological investigation, the progresses in animal studies and in vitro experiments, and the current attempts to explore potential mechanisms. Knowledge of the biological effects on immune cells associated with electromagnetic fields is critical for proper health hazard evaluation, development of safety standards, and safe exploitation of new electromagnetic devices and applications. <https://pubmed.ncbi.nlm.nih.gov/34753364/>

Outlook

In an organism, the role of the defence system against external stressors is played by the immune system consisting of various cell types. The immune cells are engaged in many physiological processes and responsible for the proper function of the whole organism. Any factor with an ability to cause immunomodulatory effects may weaken or enhance the response of the immune system (Piszcze et al. 2021). The biological effects of electromagnetic fields on immune cells have been attracting more and more attention in recent years. Researchers from numerous groups focused on both epidemiological investigations and experimental studies. Accumulated data suggested that electromagnetic exposure could affect the number and function of immune cells to some extent, including cell proportion, cell cycle, apoptosis, killing activity, cytokines contents and so on. The research objects basically covered all types of immune cells, mainly on PBMC, T lymphocytes, B lymphocytes, NK cells and macrophages. Meanwhile, there also are negative reports of electromagnetic fields on immune cells. Electromagnetic fields-induced biological effects on immune cells are influenced by two general factors: the exposure parameters (i.e., frequency, power density, exposure duration, etc.) and the composition and/or properties of the cell target. Any deviation from the two general factors will lead to inconsistent results. This may be the reason why the results are inconsistent and no comparability between different laboratories. Therefore, it is extremely difficult to find out a mechanism which could play a dominant role in the effects of various types of immune cells under electromagnetic exposure in a wide range of parameters. However, ROS, NF- κ B signaling pathway, intracellular Ca²⁺ emerged as the potential mechanisms involved in electromagnetic field-induced activation and inhibition of immune cells; nevertheless, the specific mechanisms underlying the observed bio-effects in these cells remain unknown.

EVIDENCE FOR A CONNECTION BETWEEN CORONAVIRUS DISEASE-19 AND EXPOSURE TO RADIOFREQUENCY RADIATION FROM WIRELESS COMMUNICATIONS INCLUDING 5G

Beverly Rubik, Robert R Brown. Evidence for a connection between coronavirus disease-19 and exposure to radiofrequency radiation from wireless communications including 5G. J Clin Transl Res. 2021 Sep 29;7(5):666- 681.

My comments: If there is a connection between the Covid-19 pandemic and wireless communication radiation (WCR) due to impaired immune system response, it is likely attributable to the population's total EMF exposure including ELF EMF as well as WCR (e.g., 2G-5G, Wi-Fi, and Bluetooth). 5G is not universally available and is only partially deployed in locations that have this infrastructure. Moreover, since 5G operates a fraction of the time and depends heavily on 4G communication now, I suspect that densification of 4G via small cell deployment is more likely to be problematic than 5G.

Abstract

Background and aim: Coronavirus disease (COVID-19) public health policy has focused on the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus and its effects on human health while environmental factors have been largely ignored. In considering the epidemiological triad (agent-host-environment) applicable to all disease, we investigated a possible environmental factor in the COVID-19 pandemic: ambient radiofrequency radiation from wireless communication systems including microwaves and millimeter waves.

SARS-CoV-2, the virus that caused the COVID-19 pandemic, surfaced in Wuhan, China shortly after the implementation of city-wide (fifth generation [5G] of wireless communications radiation [WCR]), and rapidly spread globally, initially demonstrating a statistical correlation to international communities with recently established 5G networks. In this study, we examined the peer-reviewed scientific literature on the detrimental bioeffects of WCR and identified several mechanisms by which WCR may have contributed to the COVID-19 pandemic as a toxic environmental cofactor. By crossing boundaries between the disciplines of biophysics and pathophysiology, we present evidence that WCR may: (1) cause morphologic changes in erythrocytes including echinocyte and rouleaux formation that can contribute to hypercoagulation; (2) impair microcirculation and reduce erythrocyte and hemoglobin levels exacerbating hypoxia; (3) amplify immune system dysfunction, including immunosuppression, autoimmunity, and hyperinflammation; (4) increase cellular oxidative stress and the production of free radicals resulting in vascular injury and organ damage; (5) increase intracellular Ca²⁺ essential for viral entry, replication, and release, in addition to promoting pro-inflammatory pathways; and (6) worsen heart arrhythmias and cardiac disorders.

Relevance for patients: In short, WCR has become a ubiquitous environmental stressor that we propose may have contributed to adverse health outcomes of patients infected with SARS-CoV-2 and increased the severity of the COVID-19 pandemic. Therefore, we recommend that all people, particularly those suffering from SARS-CoV-2 infection, reduce their exposure to WCR as much as reasonably achievable until further research better clarifies the systemic health effects associated with chronic WCR exposure.

Excerpts

Epidemiologists, including those at the CDC, consider multiple causal factors when evaluating the virulence of an agent and understanding its ability to spread and cause disease. Most importantly, these variables include environmental cofactors and the health status of the host. Evidence from the literature summarized here suggests a possible connection between several adverse health effects of WCR exposure and the clinical course of COVID-19 in that WCR may have worsened the COVID-19 pandemic by weakening the host and exacerbating COVID-19 disease. However, none of the observations discussed here prove this linkage. Specifically, the evidence does not confirm causation. Clearly COVID-19 occurs in regions with little wireless communication. Furthermore, the relative morbidity caused by WCR exposure in COVID-19 is unknown....

Another shortcoming of this paper is that we do not have access to experimental data on 5G exposures. In fact, little is known about population exposure from real-world WCR, which includes exposure to WCR infrastructure and the plethora of WCR emitting devices. In relation to this, it is difficult to accurately quantify the average power density at a given location, which varies greatly, depending on the time, specific location, time-averaging interval, frequency, and modulation scheme. For a specific municipality it depends on the antenna density, which network protocols are used, as, for example, 2G, 3G, 4G, 5G, Wi-Fi, WiMAX (Worldwide Interoperability for Microwave Access), DECT (Digitally Enhanced Cordless Telecommunications), and RADAR (Radio Detection and Ranging). There is also WCR from ubiquitous radio wave transmitters, including antennas, base stations, smart meters, mobile phones, routers, satellites, and other wireless devices currently in use. All of these signals superimpose to yield the total average power density at a given location that typically fluctuates greatly over time. No experimental studies on adverse health effects or safety issues of 5G have been reported, and none are currently planned by the industry, although this is sorely needed....

This paper points to the need for further research on nonthermal WCR exposure and its potential role in COVID-19. Moreover, some of the WCR exposure bioeffects that we discuss here — oxidative stress, inflammation, and immune system disruption — are common to many chronic diseases, including autoimmune disease and diabetes. Thus, we hypothesize that WCR exposure may also be a potential contributing factor in many chronic diseases....

Conclusion

There is a substantial overlap in pathobiology between COVID-19 and WCR exposure. The evidence presented here indicates that mechanisms involved in the clinical progression of COVID-19 could also be generated, according to experimental data, by WCR exposure. Therefore, we propose a link between adverse bioeffects of WCR exposure from wireless devices and COVID-19.

Specifically, evidence presented here supports a premise that WCR and, in particular, 5G, which involves densification of 4G, may have exacerbated the COVID-19 pandemic by weakening host immunity and increasing SARS-CoV-2 virulence by (1) causing morphologic changes in erythrocytes including echinocyte and rouleaux formation that may be contributing to hypercoagulation; (2) impairing microcirculation and reducing erythrocyte and hemoglobin levels exacerbating hypoxia; (3) amplifying immune dysfunction, including immunosuppression, autoimmunity, and hyperinflammation; (4) increasing cellular oxidative

stress and the production of free radicals exacerbating vascular injury and organ damage; (5) increasing intracellular Ca²⁺ essential for viral entry, replication, and release, in addition to promoting pro-inflammatory pathways; and (6) worsening heart arrhythmias and cardiac disorders.

WCR exposure is a widespread, yet often neglected, environmental stressor that can produce a wide range of adverse bioeffects. For decades, independent research scientists worldwide have emphasized the health risks and cumulative damage caused by WCR [42,45]. The evidence presented here is consistent with a large body of established research. Healthcare workers and policymakers should consider WCR a potentially toxic environmental stressor. Methods for reducing WCR exposure should be provided to all patients and the general population.

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8580522/>

THE ASSOCIATION BETWEEN MOBILE PHONES AND THE RISK OF BRAIN CANCER MORTALITY: A 25-YEAR CROSS-COUNTRY ANALYSIS

Mialon HM, Nesson ET. The association between mobile phones and the risk of brain cancer mortality: A 25-year cross-country analysis. *Contemporary Economic Policy*. December 4, 2019. doi:10.1111/coep.12456.

Abstract

We investigate the relationship between cell phones and brain cancer using brain cancer death rates for 88 countries between 1990 and 2015 from the World Health Organization and country-level mobile phone subscription rates from the World Bank. We estimate difference-in-difference models including country and year fixed effects and time-varying country covariates. We find that mobile phone subscription rates are positively and statistically significantly associated with death rates from brain cancer 15–20 years later. In falsification tests, we find few positive associations between mobile phone subscription rates and deaths from rectal, pancreatic, stomach, breast or lung cancer or ischemic heart disease. Finally, differential effects models suggest that mobile phone subscription rates are associated with brain cancer deaths 15–19 years later relative to deaths from other causes.

Excerpts

The link between mobile phones and brain tumors has received much attention in the media (e.g., Hakim 2016; Knutson 2016; Storrs 2016), although previous evidence has not yielded conclusive results (Hardell and Carlberg 2014; Hardell, Carlberg, and Hansson Mild 2011; Hardell et al. 2013; M. Little et al. 2012; Myung et al. 2009).

...We find a positive association between mobile phone subscriptions and brain cancer death rates. Specifically, one more mobile phone subscription per 100 people is associated with 0.003 and 0.02 additional brain cancer deaths per 100,000 people 15 and 20 years later, respectively. This result is robust to a number of specification tests, although it is sensitive to the exclusion of country-specific linear time trends. We run a number of falsification tests where we examine the relationship between mobile phone subscription rates and mortality rates from five other common forms of cancer, rectal cancer, pancreatic cancer, stomach cancer, lung cancer, and breast cancer, as well as ischemic heart disease. We find much

smaller coefficients with much larger standard errors for these other mortality outcomes, although we do find a few positive and statistically significant coefficients. Given that we find some sensitivity to country-specific linear trends and some significant effects of mobile phone subscriptions on other cancers, we estimate a differential effects model where we compare the association between mobile phone subscription rates and brain cancer mortality relative to other disease mortality. In these models, we additionally find statistically significant, positive coefficients, suggesting that mobile phone prevalence is associated with brain cancer mortality about 15–20 years later relative to other disease mortality.

Our study is the first to analyze the potential link between brain cancer and mobile phone use on the aggregate, across multiple countries, and over a period spanning more than 20 years....

We collect information on annual mobile cellular subscriptions per 100 people from the World Bank (The World Bank 2016). The first mobile cellular phones were introduced in the Nordic countries in the early 1980s, and by 1990, Iceland, Norway, and Sweden had more than three mobile phone subscriptions per 100 people.

Additionally, the United States, Hong Kong, Singapore, Kuwait, and New Zealand were relatively early adopters, with more than one subscription per 100 people by 1990. By 1995, the European Union had an average of over four subscriptions per 100 people, growing to over 53 subscriptions per 100 people in 2000. North American and high income Asian and Middle Eastern countries followed similar patterns.

Table 1 shows summary statistics for all variables in our sample over the period from 1990 to 2015. On average, the annual brain cancer death rate is about 3.7 individuals per 100,000. There are about 62 mobile phone subscriptions per 100 people on average. However, there is a lot of temporal variation in mobile phone subscriptions, which we detail more below....

Results from Table 2:

... one more mobile phone subscription per 1,000 people is associated with 0.00012 more brain cancer deaths per 1,000 people 15 years later and 0.00073 more brain cancer deaths per 1,000 people 20 years later.

In the first column of Table 4, we find very similar effects to Table 2, and we again find that mobile phone subscriptions are related to increases in brain cancer mortality starting about 15–25 years later. The next four columns show results from differential effects models. In these models, we again find a positive and statistically significant association between mobile phone subscription rates and brain cancer mortality 15–19 years later relative to mortality from the other diseases studied. Our results here are robust to specification changes, with the exception that we lose statistical significance when not using country weights.³ However, we do note that the mobile phone coefficient is much larger for the 15–19 year lag compared to other lags.

We find a statistically significant relationship between mobile phone subscriptions and brain cancer death rates 15–20 years later. Our estimated effect sizes are small but statistically

significant. The implications of our findings are potentially very large. Almost 16,000 people died from brain cancer in the United States in 2014. If our findings are correct, the number of brain cancer deaths will continue to rise as the number of mobile phone users rose rapidly in the early part of the century. To err on the side of caution, individuals may want to make greater use of the speaker phone and/or texting options on their cell phones or investigate cell phone cases that may deflect radiation away from their ears.

Some caveats to our analysis are warranted. First, we cannot make an unambiguous claim of causality based on our results. Although we control for other characteristics likely to affect brain cancer death rates and find limited statistically significant relationships between mobile phone subscriptions and other mortality outcomes, there may be yet other factors that are correlated with mobile phone subscriptions and are also drivers of changes in brain cancer death rates. For example, ionizing radiation from computed tomography (CT) scans is an established risk factor for brain cancer (Pearce et al. 2012), and if the use of CT scans is correlated with mobile phone use, our estimates may be biased. We include controls for health spending, which should control for the number of X-rays to some degree, and moreover, if X-rays and mobile phone adoption were truly correlated, we would expect mobile phones to have an effect on cancer mortality rates in other sites. But, omitted variables do remain a concern in our analysis.

Second, our measure of mobile phone use, mobile phone subscriptions per 100 people, may not be perfectly accurate....

Third, long lags between mobile phone subscriptions and brain cancer death rates mean that, depending on the specified lag between mobile phone subscriptions and mortality, the large recent variations in mobile phone adoption may not be part of our analysis. Additionally, the recent innovations in mobile phone technology, for example switching from 2G to 3G and 4G have resulted in changes in the radiation emitted from mobile phones (Croft et al. 2010; Leung et al. 2011). Moreover, the transition of peoples' use of mobile phones as predominantly phones to the rise of texting and other forms of social media may also change the relationship between mobile phones and brain cancer in the future.

<https://onlinelibrary.wiley.com/doi/abs/10.1111/coep.12456>

EFFECT OF CELL PHONE USE ON SALIVARY COMPONENTS; A REVIEW OF LITERATURE

Fateme Arbabi Kalati, Tahereh Nosratzahi. Effect of cell phone use on salivary components; a review of literature. J Complement Integr Med. 2021 Nov 25. doi: 10.1515/jcim-2021-0397.

Abstract

Mobile phones have been increasingly used in the past decade and have become a cultural instrument. There is a great concern over the harmful effects of electromagnetic and radiofrequency waves as well as microwaves generated by mobile phones and their telecommunication stations on health. The saliva plays an important role in preserving oral homeostasis as the first defensive line against the microbial invasion which protects oral mucosa mechanically and immunologically. A search was run in Pub med, Goggle Scholar, Medline, and Web of Science databases using the following keywords: cell phone, mobile phone, antioxidant profile, saliva, oxidative stress, interleukin, and inflammation. Sixty-five

published articles were identified. Studies on the use of cell phones as educational aids, the use of immune histochemistry on salivary glands, or the evaluation of saliva in individuals with specific conditions, such as the use of orthodontic brackets, were excluded. In addition, duplicate articles are eliminated, and finally, 14 articles were included in the present study. Nowadays mobile phone is very popular, causing concern about the effect it has on people's health. Parotid salivary glands are in close contact with a cell phone while talking with the phone and the possibility of being affected by them; so this study was designed to investigate the effect of cell phone use on salivary components.

<https://pubmed.ncbi.nlm.nih.gov/34821111/>

Excerpt

The above studies show that using cell phones affects the antioxidative system, immune system, and inflammatory system of saliva. Although to date, no specific medical condition has been attributed to these changes, decreasing the frequency and duration of cellphone use is advisable.

Thus, based on these findings, it is recommended that a long-term/or excessive use of mobile phones, especially by young individuals, should be avoided. This goal can be accomplished by telehealth technology promotion activities targeting the more sensitive ages, children, and adolescents since their developing brains absorb more EMR from a mobile phone. Such activities include group discussions, public presentations, and mass communication through available electronic and print media sources.

OCCUPATIONAL EXPOSURE TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS (REVIEW)

Rianne Stam. 2021 Nov 16. doi: 10.2486/indhealth.2021-0129.

Abstract

High exposures to radiofrequency electromagnetic fields (RF EMF) are possible in workplaces involving sources used for broadcasting, telecommunication, security and identification, remote sensing and the heating and drying of goods. A systematic literature review of occupational RF EMF exposure measurements could help to clarify where more attention to occupational safety may be needed. This review identifies specific sources of occupational RF EMF exposure and compares the published maximum exposures to occupational exposure limits. A systematic search for peer-reviewed publications was conducted via PubMed and Scopus. Relevant grey literature was collected via web searches. For each publication, the highest measured electric field strength, magnetic flux density or power density was extracted. Maximum exposures exceeding the limits were reported for dielectric heating, scanners for security and radiofrequency identification, plasma devices and broadcasting and telecommunication transmitters. Occupational exposure exceeding the limits was rare for microwave heating and radar applications. Some publications concerned case studies of occupational accidents followed by a medical investigation of thermal health effects. These were found for broadcasting antennas, radar installations and a microwave oven and often involved maintenance personnel. New sources of occupational exposure such as those in fifth generation telecommunication systems or energy transition will require further assessment. Open access paper: https://www.jstage.jst.go.jp/article/indhealth/advpub/0/advpub_2021-0129/_article

MAGNETIC FIELD-INDUCED INTERACTIONS BETWEEN PHONES CONTAINING MAGNETS AND CARDIOVASCULAR IMPLANTABLE ELECTRONIC DEVICES - FLIP IT TO BE SAFE?

Lacour P, Dang PL, Heinzel FR, Parwani AS, Bähr F, Kucher A, Hohendanner F, Niendorf T, Rahimi F, Saha N, Han H, Rubarth K, Sherif M, Boldt LH, Pieske B, Blaschke F. *Magnetic Field-2021 Nov 9:S1547-5271(21)02334-1. doi: 10.1016/j.hrthm.2021.11.010.*

Abstract

Background: Recent case reports and small studies have reported activation of the magnet-sensitive switches in cardiovascular implantable electronic devices (CIED) by the new iPhone 12 series, initiating asynchronous pacing in pacemakers and suspension of anti-tachycardia therapies in ICDs.

Objective and methods: We performed a prospective single-center observational study to quantify the risk of magnetic field interactions of the iPhone 12 with CIEDs. A representative model of each CIED series from all manufacturers was tested ex vivo. Incidence and minimum distance necessary for magnet mode triggering were analyzed in 164 CIED patients with either the front or the back of the phone facing the device. The magnetic field of the iPhone 12 was analyzed using a 3-axis hall probe.

Results: Ex vivo, magnetic interferences occurred in 84.6% with the back compared to 46.2% with the front of the iPhone 12 facing the CIED. In vivo, activation of the magnet-sensitive switch occurred in 30 CIED patients (18.3%; 21 pacemaker, 9 ICDs) when the iPhone 12 was placed in close proximity over the CIED pocket and the back of the phone was facing the skin. Multiple binary logistic regression analysis identified the implantation depth (95% confidence interval [CI], 0.02 to 0.24) as independent predictor of magnet-sensitive switch activation.

Conclusion: Magnetic field interactions occur only in close proximity, and with precise alignment of the iPhone 12 and CIEDs. It is important to advise CIED patients to not put the iPhone 12 directly on the skin above the CIED. Further recommendations are not necessary.

Open access paper: [https://www.heartrhythmjournal.com/article/S1547-5271\(21\)02334-1/fulltext](https://www.heartrhythmjournal.com/article/S1547-5271(21)02334-1/fulltext)

COMPACT EXPOSIMETER DEVICE FOR THE CHARACTERIZATION AND RECORDING OF ELECTROMAGNETIC FIELDS FROM 78 MHZ TO 6 GHZ WITH SEVERAL NARROW BANDS (300 KHZ)

Marco Xavier Rivera González, Nazario Félix González, Isabel López, Juan Sebastián Ochoa Zambrano, Andrés Miranda Martínez, Ceferino Maestú Unturbe. *Sensors (Basel).* 2021 Nov 7;21(21):7395. doi: 10.3390/s21217395.

Abstract

A novel compact device with spectrum analyzer characteristics has been designed, which allows the measuring of the maximum power received in multiple narrow frequency bands of 300 kHz, recording the entire spectrum from 78 MHz to 6 GHz; the device is capable of measuring the entire communications spectrum and detecting multiple sources of electromagnetic fields using the same communications band. The proposed device permits the evaluation of the cross-talk effect that, in conventional exposimeters, generates a mistake

estimation of electromagnetic fields. The device was calibrated in an anechoic chamber for far-fields and was validated against a portable spectrum analyzer in a residential area. A strong correlation between the two devices with a confidence higher than 95% was obtained; indicating that the device could be considered as an important tool for electromagnetic field studies.

Conclusions

A compact exposimeter system has been proposed, which has similar characteristics to a spectrum analyzer, without losing the advantages of conventional exposimeters, and can be a useful tool for electromagnetic field studies analyzing a far-field. The proposed exposimeter measures the received power in the spectrum between 78 MHz and 6 GHz with a resolution bandwidth of 300 KHz, measuring approximately 19,500 narrow bands; the proposed exposimeter has passed the testing phase compared with spectrum analyzers with a significant factor of 0.948 for Student's t-test and a correlation factor of 0.9682 between the measurements of both devices.

The proposed exposimeter requires one second to measure the entire radioelectric spectrum; therefore, it has less sensitivity to variations in electromagnetic fields. This could be improved by restricting the measurements to the communication bands of interest; however, information from the rest of the radio spectrum is lost. It is important to define the relationship between the communication bands to be measured and the sensitivity to electromagnetic field variations with respect to time.

In future work, it is expected to use several of the proposed devices in electromagnetic field surveys to cover residential areas and provide information as an electromagnetic field sensor network to be used in epidemiological studies in the deployed area.

Open access paper: <https://www.mdpi.com/1424-8220/21/21/7395/htm>

COMPUTATIONAL MODELING INVESTIGATION OF PULSED HIGH PEAK POWER MICROWAVES AND THE POTENTIAL FOR TRAUMATIC BRAIN INJURY

Amy M Dagro, Justin W Wilkerson, Thaddeus P Thomas, Benjamin T Kalinosky, Jason A Payne. Sci Adv. 2021 Oct 29;7(44):eabd8405. doi: 10.1126/sciadv.abd8405. Amy Dagro and Thaddeus Thomas are with the U.S. Army Research Laboratory, Aberdeen Proving Ground, MD; Benjamin Kalinosky is with General Dynamics Information Technology, JBSA Fort Sam Houston, San Antonio, TX; and Jason Payne is with U.S. Air Force Research Laboratory, 711th Human Performance Wing, Airman Systems Directorate, Bioeffects Division, Radio Frequency Bioeffects Branch, JBSA Fort Sam Houston, San Antonio, TX.

Abstract

When considering safety standards for human exposure to radiofrequency (RF) and microwave energy, the dominant concerns pertain to a thermal effect. However, in the case of high-power pulsed RF/microwave energy, a rapid thermal expansion can lead to stress waves within the body. In this study, a computational model is used to estimate the temperature profile in the human brain resulting from exposure to various RF/microwave incident field parameters. The temperatures are subsequently used to simulate the resulting mechanical response of the brain. Our simulations show that, for certain extremely high-power microwave exposures (permissible by current safety standards), very high stresses may

occur within the brain that may have implications for neuropathological effects. Although the required power densities are orders of magnitude larger than most real-world exposure conditions, they can be achieved with devices meant to emit high-power electromagnetic pulses in military and research applications.

Excerpts

"The bulk of scientific literature uses continuous waves and moderate field strengths (typical of real-life scenarios), with less emphasis on pulsed fields of very high peak strength that may occur with ultrawideband pulse generators or EM pulse simulators (4). It is worth investigating whether extremely high peak power sources applied with a slow repetition frequency, or low duty cycle, can induce injurious effects without thermal buildup greater than a few degrees Celsius."

"With the exception of low intracranial absorption at 1400 MHz, the highest ratio of peak average intracranial SAR* to peak average skin SAR* occurs between 1 to 1.8 GHz."

"The MAE, also referred to as "microwave hearing" or the "Frey effect" due to its discovery by Allan Frey in 1961 (7, 8), was initially observed when subjects standing up to hundreds of feet away from a radar transponder could hear an audible tonal noise (e.g., chirping, buzzing, or clicking). The scientific underpinnings of the MAE were controversial for the first several years (9–11). After more than a decade of investigations, it became generally accepted that the perceived sound is due to the cochlea detecting stress waves that result from a rapid temperature rise in tissues within the head due to pulsed RF/microwave exposure (11, 12)."

"Typically, relatively low-average powers and small temperature changes (10–6°C) are required to elicit the MAE (12). Although adverse health effects from the MAE have not been previously established, one study on rodents suggests that very high-peak power pulsed microwaves can result in cognitive deficits (13)."

"This study uses a two-simulation approach to investigate whether an HPM source could theoretically induce adverse mechanical responses within the brain."

"This study has shown that, by applying a small temperature increase (<0.0005°C) in a very short amount of time (less than several microseconds), potentially injurious stress waves are created."

"For frequencies between 400 MHz to 2 GHz, the IEEE C95.1 RF exposure guidelines limit the exposure reference limit (ERL) to fmhz/200 (W/m²) over an averaging time of 30 min. For 1-GHz exposures, the IEEE C95.1 ERL of 5 W/m² over 30 min would equate to an average energy density of 9000 J/m². Our computational model shows that, for sufficiently high incident power densities, a single pulse could potentially result in biologically meaningful pressures. For example, large pressures may occur following 1-GHz frequency, a pulse duration of 5 μs, and incident power densities of at least 1.5×10^7 W/m². The energy density associated with such a pulse would be equal to $P_{IN} \times t_d$ or 75 J/m² (significantly less than the ERL standard)."

" Note that the proposed HPM power densities in this study are extremely large and several orders of magnitude larger than power densities typically experienced by the public. As an

illustrative example, at around 200 feet from a cell phone base station, a person will be exposed to a power density of only 0.001 mW/cm² or less (36). This study establishes a testable hypothesis between potential neurocognitive effects and the thermoelastic mechanism from HPM systems. To date, however, adverse effects from HPM systems have not been established in the scientific literature."

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8555891/>

ASSESSMENT OF HUMAN EXPOSURE IN IASI-CITY USING DATA PROVIDED BY THE (ROMANIAN) NATIONAL AUTONOMOUS RF- EMF MONITORING SYSTEM

M. Ursachianu, O. Bejenaru, C. Lazarescu, A. Salceanu and M. Paulet., 2021, pp. 225-230, doi: 10.1109/SIELMEN53755.2021.9600350.

Abstract

In the last decades, we are all witnessed to an explosive increase and development of applications that integrate the electromagnetic environment in our day-by-day life. The principal and delicate problem of the human exposure to a various types of electromagnetic field (EMF) sources is very challenging and up to date one, being considered to have a higher level of importance especially from the safety point of view. The population has always been reluctant and ambiguous about the introduction of new technologies, like the actual case of the implementation around the world of the new 5G NR mobile communications technology for example. For that, the responsible authorities from different countries must conduct studies to ensure and to track the fact that the exposure at new technology is safe or not. So, that being said, the process of monitoring the EMF produce by a diversity of different sources of radiofrequency electromagnetic fields (RF-EMFs) should be a priority in our days. In the study presented in this paper, using the National Autonomous Broadband EMF Monitoring System, a continuous 24/365(hours/days) monitoring system, we analyze real-time monitoring data for all months of the year 2020 for all EMF monitoring fixed station installed at that time in Iasi-city giving a special attention for the last month of the year, December 2020. It was made also an analysis and comparison off the measured data with those imposed by international organizations and national regulatory bodies. At the final a set of preliminary conclusions have been taken. <https://ieeexplore.ieee.org/document/9600350>

Conclusions

Evaluation the levels of the electromagnetic fields generated by a diversity of communication systems and other sources of RF-EMFs should be a priority in our days. An Autonomous Continuous and Intelligent Broadband EMF monitoring system with multiband frequency sensors represent an important step in this area. The network of tri-axials sensors from EMF monitoring system measures the total levels of electromagnetic field from all surrounding sources and can also do a real-time measure of the assessment results to the public.

The continuous monitoring of the electromagnetic field is very important to realize to reassure the population who is in an area of ambiguity regarding the exposure to electromagnetic radiation generated by various sources of radio frequency electromagnetic fields in our country. The exposure to RF-EMFs is an actual problem of concern for many people in nowadays and should be a priority area of investigation, especially regarding the emergence of new communication technologies, like 5G NR.

The present study offers an insight view into the EMF exposure levels recorded by the national EMF real-time monitoring system, for a full month of the year 2020. The results of our analysis indicate that the exposure levels to radio electromagnetic fields are much more below the reference levels values adopted by the national legislation for population.

The E-field values of different transmitters may vary over the time. In the case of base stations of cellular mobile services, the transmitted power depends highly on the actual traffic load and user behavior. During a monitoring of 24h the electric field values was well below the limits. Traffic load may also depend on the time of the day, day of the week, and even in season as we will see in a future study. Monitoring the E-Field values from mobile base station antenna may also depend on the distance from the source.

An EMF monitoring station with a tri-axial electric field probe who can measured the electric field in a broadband frequency range like one of those used in the national EMF monitoring system could be also use for providing a good assessment of human exposure to EMF generated by new types of mobile communication services. 5G NR (NR – New Radio).

Further research can be directed on the study of EMF monitoring values in case of extending the evaluation study over more EMF monitoring stations from different cities of the country, eventually for a long period of time, in order to obtain a better assess of the coverage and the temporal evolution over the problem of the exposure to different electromagnetic field sources.

5G BASE STATION DEPLOYMENT REVIEW FOR RF RADIATION

M. S. Elbasheir, R. A. Saeed, S. Edam. 5G Base Station Deployment Review for RF Radiation. 2021 International Symposium on Networks, Computers and Communications (ISNCC), 2021, pp. 1-5, doi: 10.1109/ISNCC52172.2021.9615689.

Abstract

The 5G is expected to make great change for Mobile network and technology in the coming years. there is extensive discussion about the electromagnetic radiation that the 5G will contribute and its impact to the human and other technologies. The aggressive deployment of the technology associated with the new massive.

Internet of Things (IoT) devices, all are indicator to the great electromagnetic radiation and exposure that 5G may cause. This problem considered as a challenge constrain for deployment of massive 5G base stations especially in residential areas. This paper reviews the recent works on the Electromagnetic Fields (EMF) radiation assessment for 5G base stations (BS) on human evaluation and analysis from different perspectives. The review covers the international standard exposure limits adopted by some of the regulatory bodies. The reviewed cases are classified into two main categories, assessment based on model simulation, and on field measurement, where both categories have the same objective to assess and evaluate the EMF radiation exposure from BSs.

Conclusions

EMF exposure has increasingly earned attention in the community and media. Recently the 5G has been proposed to be deployed under the 3GPP released 18. One of the main

challenges for 5G deployment is the EMF exposure, especially when accumulated with the other deployed technologies i.e., 4G, 3G and low-power wide- area network (LPWAN) technologies. This paper reviewed the EMF radiation assessment from 5G BSs, where we reviewed some recent related work recently done through measurement and simulations. Various results were discussed, analyzed and compared with the regulation bodies like ICNIRP and FCC. there are many EMF assessment methodologies and techniques were presented and discussed. Most of the recommendations were to conduct more investigations with different perspectives, parameters, assumptions and scenarios.
<https://ieeexplore.ieee.org/document/9615689>

MAGNETITE PARTICLE PRESENCE IN THE HUMAN BRAIN: A COMPUTATIONAL DOSIMETRIC STUDY TO EMPHASIZE THE NEED OF A COMPLETE ASSESSMENT OF THE ELECTROMAGNETIC POWER DEPOSITION AT 3.5 GHZ

D. Vatamanu, S. Miclaus. Magnetite Particle Presence in the Human Brain: A Computational Dosimetric Study to Emphasize the Need of a Complete Assessment of the Electromagnetic Power Deposition at 3.5 GHz. Eng. Technol. Appl. Sci. Res., 11(5): 7720–7729, Oct. 2021.

Abstract

The growing evidence of increased magnetite nanoparticles (both endo- and exo-genic) in the human brain raises the importance of assessing the entire power deposition when electromagnetic waves at GHz frequencies propagate in such tissues. This frequency range corresponds to many popular portable communication devices that emit radiation close to a human's head. At these frequencies, the current dosimetric numerical codes can not accurately compute the magnetic losses part. This is due to the lack of an implemented computational algorithm based on solving the coupled Maxwell and Landau-Lifshitz-Gilbert equations, in the case of magneto- dielectrics, considering eddy currents losses and specific properties of magnetic sub-millimetric particles. This paper focuses on analyzing the limits and the inconsistencies when using commercial dosimetric numerical software to analyze the total absorbed power in brain models having ferrimagnetic content and being exposed to 3.5GHz electromagnetic waves. Magnetic losses computed using Polder's permeability tensor as constitutive relation lead to unreliable results. However, using such software can provide a preliminary view of the electromagnetic impact of ultra- and super-high frequencies on magnetic-dielectric tissues.

Open access paper: <https://www.etasr.com/index.php/ETASR/article/view/4466>

1,800 MHZ RF ELECTROMAGNETIC IRRADIATION IMPAIRS NEURITE OUTGROWTH WITH A DECREASE IN RAP1-GTP IN PRIMARY MOUSE HIPPOCAMPAL NEURONS AND NEURO2A CELLS

Li Yanqi, Deng Ping, Chen Chunhai, Ma Qinlong, Pi Huifeng, He Mindi, Lu Yonghui, Gao Peng, Zhou Chao, He Zhixin, Zhang Yanwen, Yu Zhengping, Zhang Lei. Vol. 9:1854. 2021. doi: 10.3389/fpubh.2021.771508

Abstract

Background: With the global popularity of communication devices such as mobile phones, there are increasing concerns regarding the effect of radiofrequency electromagnetic radiation (RF-EMR) on the brain, one of the most important organs sensitive to RF-EMR exposure at 1,800 MHz. However, the effects of RF-EMR exposure on neuronal cells are

unclear. Neurite outgrowth plays a critical role in brain development, therefore, determining the effects of 1,800 MHz RF-EMR exposure on neurite outgrowth is important for exploring its effects on brain development.

Objectives: We aimed to investigate the effects of 1,800 MHz RF-EMR exposure for 48 h on neurite outgrowth in neuronal cells and to explore the associated role of the Rap1 signaling pathway.

Material and Methods: Primary hippocampal neurons from C57BL/6 mice and Neuro2a cells were exposed to 1,800 MHz RF-EMR at a specific absorption rate (SAR) value of 4 W/kg for 48 h. CCK-8 assays were used to determine the cell viability after 24, 48, and 72 h of irradiation. Neurite outgrowth of primary hippocampal neurons (DIV 2) and Neuro2a cells was observed with a 20 × optical microscope and recognized by ImageJ software. Rap1a and Rap1b gene expressions were detected by real-time quantitative PCR. Rap1, Rap1a, Rap1b, Rap1GAP, and p-MEK1/2 protein expressions were detected by western blot. Rap1-GTP expression was detected by immunoprecipitation. The role of Rap1-GTP was assessed by transfecting a constitutively active mutant plasmid (Rap1-Gly_Val-GFP) into Neuro2a cells.

Results: Exposure to 1,800 MHz RF-EMR for 24, 48, and 72 h at 4 W/kg did not influence cell viability. The neurite length, primary and secondary neurite numbers, and branch points of primary mouse hippocampal neurons were significantly impaired by 48-h RF-EMR exposure. The neurite-bearing cell percentage and neurite length of Neuro2a cells were also inhibited by 48-h RF-EMR exposure. Rap1 activity was inhibited by 48-h RF-EMR with no detectable alteration in either gene or protein expression of Rap1. The protein expression of Rap1GAP increased after 48-h RF-EMR exposure, while the expression of p-MEK1/2 protein decreased. Overexpression of constitutively active Rap1 reversed the decrease in Rap1-GTP and the neurite outgrowth impairment in Neuro2a cells induced by 1,800 MHz RF-EMR exposure for 48 h.

Conclusion: Rap1 activity and related signaling pathways are involved in the disturbance of neurite outgrowth induced by 48-h 1,800 MHz RF-EMR exposure. The effects of RF-EMR exposure on neuronal development in infants and children deserve greater focus.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.771508>

SOLAR AND GEOMAGNETIC ACTIVITY ENHANCE THE EFFECTS OF AIR POLLUTANTS ON ATRIAL FIBRILLATION

Zilli Vieira CL, Link MS, Garshick E, Peralta AA, Luttmann-Gibson H, Laden F, Liu M, Gold DR, Koutrakis P. Solar and geomagnetic activity enhance the effects of air pollutants on atrial fibrillation. *Europace*. 2021 Nov 13;euab269. doi: 10.1093/europace/euab269.

Abstract

Aims: Cardiac arrhythmias have been associated with intense solar and geomagnetic activity (SGA) and exposures to air pollution.

Methods: We examined whether oscillations of SGA can modify the effect of hourly exposures to air pollutants on atrial fibrillation ≥30 s (AF) risk in patients with dual-chamber

implantable cardioverter-defibrillators. The effects of SGA on ambient particulate matter <2.5 µm (PM_{2.5}), black carbon (BC), ultrafine particles (PN), and associations with AF were assessed. Measures of SGA included solar wind proton density (SW), total interplanetary magnetic field strength (IMF), and Kp index, a measure of global geomagnetic activity.

Results: Overall time lags between 0 and 24 h, periods of increased SGA (>50th percentile in IMF, SW, and Kp index) enhanced the effects of all three air pollutants on AF, while during periods of reduced SGA the associations were considerably weaker or absent. During periods of intense SW 6 h prior to an AF event, the odds ratio (OR) for PM_{2.5} exposure per interquartile range (IQR) of 5.6 µg/m³ was 1.7 [95% confidence interval (CI) 1.3-2.3, P = 0.0001]. For periods of reduced SW, the OR for PM_{2.5} exposure per IQR was 1.2 (95% CI 0.9-1.5; P = 0.27). There were similar effects for PN and BC exposures. In patients with multiple AF events per hour, the associations with air pollutants during intense SGA were even greater.

Conclusion: The effects of air pollutants up to 24 h before AF events were enhanced during periods of increased SGA. Our results suggest that these effects may account for variation in AF risk. <https://pubmed.ncbi.nlm.nih.gov/34791174/>

HUMAN RESTING-STATE EEG AND RADIOFREQUENCY GSM MOBILE PHONE EXPOSURE: THE IMPACT OF THE INDIVIDUAL ALPHA FREQUENCY

Jasmina Wallace, Lydia Yahia-Cherif, Christophe Gitton, Laurent Hugueville, Jean-Didier Lemaréchal, Brahim Selmaoui. Human resting-state EEG and radiofrequency GSM mobile phone exposure: The impact of the individual alpha frequency. *Int J Radiat Biol.* 2021 Nov 19;1-29. doi: 10.1080/09553002.2021.2009146.

Abstract

Purpose- With the extensive use of mobile phone (MP) several studies have been realized to investigate the effects of radiofrequency electromagnetic fields (RF-EMF) exposure on brain activity at rest via electroencephalography (EEG), and the most consistent effect has been seen on the alpha band power spectral density (PSD). However, some studies reported an increase or a decrease of the PSD, while others showed no effect. It has been suggested that these differences might partly be due to a variability of the physiological state of the brain between subjects. So, the aim of this study was to investigate the alpha band modulation, exploring the impact of the alpha band frequency ranges applied in the PSD analysis.

Materials & Methods- Twenty-one healthy volunteers took part in the study with a double-blind, randomized and counterbalanced crossover design, during which eyes-open (EO) and eyes-closed (EC) resting-state EEG was recorded. The exposure system was a sham or a real GSM (global system for mobile) 900 MHz MP (pulse modulated at 217 Hz, mean power of 250 mW and 2 W peak, with a maximum specific absorption rate of 0.70 W/kg on 1 g tissue). The experimental protocol presented a baseline recording phase without MP exposure, an exposure phase during which the exposure system was placed against the left ear, and the post-exposure phase without MP. EEG data from baseline and exposure phases were analyzed and PSD was computed for the alpha band in the fixed range of 8-12 Hz and for the individual alpha band frequency range (IAF).

Results- Results showed a trend in decrease or increase of EEG power of both alpha oscillations during exposure in relation to EC and EO recording conditions, respectively, but not reaching statistical significance. Findings did not provide evidence for a different sensitivity to RF-EMF MP related to individual variability in the frequency of the alpha band.

Conclusion- In conclusion, these results did not show alpha band activity modulation during resting-state under RF-EMF. It might be argued the need of a delay after the exposure in order to appreciate an EEG spectral power modulation related to RF-EMF exposure.

<https://pubmed.ncbi.nlm.nih.gov/34797205/>

DOES EXPOSURE TO RADIATION EMITTED FROM MOBILE JAMMERS INFLUENCE SPATIAL MEMORY?

A. Shahi, F. Shahnazar, S. Nematolahi, A. Dehghan, M.B. Shojaeifard. Does exposure to radiation emitted from mobile jammers influence the spatial memory? *Int J Radiat Res* 2021, 19(4): 993-1000.

Abstract

Background: The central nervous system is sensitive to radiation exposure as environmental pollution. This project aimed to evaluate the influence of jammer exposure radiation on spatial learning and memory.

Materials and Methods: 50 adult male Sprague-Dawley rats were divided into five groups. The experimental groups were exposed to jammer radiation for 2h/day once or 2h/day for two weeks. Sham groups were similar to the experimental groups which were exposed to switch-off. The distance from the jammer router to the animals' cages was 30 cm. In the second phase, using Morris Water Maze, the effect of jammer exposure radiation on spatial learning and memory was studied.

Results: Data showed that radiation exposure once a day for 2h caused a significant increase in the learning procedure in the experimental group1, but we observed an increase in the parameters of distance and escape latency in finding the platform for two weeks' exposure in the experimental group 2.

Conclusion: Results indicated that probably the motivation of escape and use of the different cues led to learning in the animals, while disrupting mobile performance through jammer router from animal environment caused a better brain performance in the spatial learning and memory in short term jammer radiation-treated animals group. However, with the same test conditions, in long term jammer the radiation-treated animals' motivation reduced, which affected the responses and performance and reduced the learning. Moreover, environmental conflicts such as radiofrequency waves lead to behavioral alternations.

Exposure set up

The animals were placed in Plexiglas cages individually and exposed to radiation. The electromagnetic field measurements were done using an Aaronia Spectran HF-4060 device (Euscheid, Germany) with the frequency range 100 MHz -6 GHz. The frequency band, power output, and power density measured from electromagnetic fields of Jammer router were 1052-1979(MHz), -32 ± 3 (dB mw), 86.18000 ($\mu\text{w}/\text{m}^2$), and ICNIRP 0.31(%) respectively.

Jammer router (MB06 Mobile Blocker) works in four different frequency ranges (Global System for Mobile Communications [GSM], digital cellular service, code division multiple access, third generation). Shielding radii for this jammer are up to 40m.
Open access paper: <http://ijrr.com/article-1-3981-en.html>

ELECTROMAGNETIC FIELD EXPOSURE-INDUCED DEPRESSION FEATURES COULD BE ALLEVIATED BY HEAT ACCLIMATION BASED ON REMODELING THE GUT MICROBIOTA

Luo X, Huang X, Luo Z, Wang Z, He G, Tan Y, Zhang B, Zhou H, Li P, Shen T, Yu X, Yang X. *Ecotoxicol Environ Saf*. 2021 Nov 15;228:112980. doi: 10.1016/j.ecoenv.2021.112980.

Highlights

- EMF exposure induced depression features.
- EMF cause gut microbiota and metabolites disturbance similar to depression model.
- HA alleviated depression caused by EMF.
- HA alleviated depression by gut microbiota, might be mechanism of cross-tolerance.

Abstract

Background: Electromagnetic pollution cannot be ignored. Long-term low-dose electromagnetic field (EMF) exposure can cause central nervous system dysfunction without effective prevention.

Materials/methods: Male C57BL/6J mice (6-8 weeks, 17-20 g) were used in this study. Depression-like and anxiety-like behaviors detected by behavioral experiments were compared among different treatments. 16S rRNA gene sequencing and non-targeted liquid chromatography-mass spectrometry (LC-MS) metabolomics were used to explore the relationship between EMF exposure and heat acclimation (HA) effects on gut microbes and serum metabolites.

Results: Both EMF and HA regulated the proportions of p_Firmicutes and p_Bacteroidota. EMF exposure caused the proportions of 6 kinds of bacteria, such as g_Butyricoccus and g_Anaerotruncus, to change significantly ($p < 0.05$). HA restored the balance of gut microbes that was affected by EMF exposure and the proportion of probiotics (g_Lactobacillus) increased significantly ($p < 0.01$). Serum metabolite analysis suggested that HA alleviated the disturbance of serum metabolites (such as cholesterol and D-mannose) induced by EMF exposure. Both the metabolic KEGG pathways and PICRUSt functional analysis demonstrated that tryptophan metabolism, pyrimidine metabolism and amino acid biosynthesis were involved.

Conclusions: EMF exposure not only led to depression-like neurobehavioral disorders, but also to gut microbiota imbalance. HA alleviated the depression features caused by EMF exposure. Based on the analysis of gut microbiota associated with serum metabolites, we speculated that gut microbiota might play a vital role in the cross-tolerance provided by HA.
<https://pubmed.ncbi.nlm.nih.gov/34794024/>

THE EFFECTS OF DIFFERENT HERBALS ON THE RAT HIPPOCAMPUS EXPOSED TO ELECTROMAGNETIC FIELD FOR ONE HOUR DURING THE PRENATAL PERIOD

Ömür Gülsüm Deniz, Süleyman Kaplan. J Chem Neuroanat. 2021 Nov 19;102043. doi: 10.1016/j.jchemneu.2021.102043.

Highlights

- 900 MHz electromagnetic field exposure can cause adverse effects on the hippocampus.
- Garcinia kola, Momordica charantia, thymoquinone may reduce deleterious effects of electromagnetic field.
- It was revealed that the protective effect of Momordica charantia is stronger than other antioxidant substances used in the present study.

Note: Momordica charantia is commonly called bitter melon.

Abstract

The purpose of this study was to highlight the possible effects on the hippocampus of the electromagnetic field (EMF) emitted by mobile phones, and to investigate whether these potential effects can be reduced using various antioxidant substances. Twenty-seven female Wistar albino rats were divided into nine equal groups, each containing three pregnant rats aged 8-10 weeks and weighing 200-250 gr. The EMF groups were exposed to 900 Megahertz (MHz) EMF for 1hour (hr) a day for 21 days. No EMF exposure was applied to the Cont and also the groups given only Garcinia kola (GK), Momordica charantia (MC), and thymoquinone (TQ). The Sham group was kept in the polycarbonate EMF exposure system, but was not exposed to EMF. Four weeks after birth, rat pups were subjected to behavioural tests. Brain tissue samples were evaluated using histological, stereological, functional, and immunohistochemical methods. The numbers of pyramidal neurons in the rat cornu ammonis (CA) were determined using the optical fractionator method. Superoxide dismutase (SOD) and catalase (CAT) enzyme activities in the blood samples were also evaluated. The analysis data indicated that total pyramidal neuron numbers were decreased significantly in the CA of the EMF (1hr) group ($p<0.01$). Our results also showed that the protective effect of MC was more potent than that of the other antioxidant substances ($p<0.01$). A 900MHz EMF can cause deleterious changes in the brain. It can also be suggested that GK, MC and TQ are capable of reducing these adverse effects. <https://pubmed.ncbi.nlm.nih.gov/34808256/>

ENVIRONMENTAL FACTORS AND RISKS OF COGNITIVE IMPAIRMENT AND DEMENTIA: A SYSTEMATIC REVIEW AND META- ANALYSIS

Yong-Li Zhao, Yi Qu, Ya-Nan Ou, Ya-Ru Zhang, Lan Tan, Jin-Tai Yu. Ageing Res Rev. 2021 Oct 29;72:101504. doi: 10.1016/j.arr.2021.101504.

Highlights

- Air pollution, tobacco smoke and pesticides could elevate the risks of dementia.
- Exposure to electromagnetic fields was found positive associated with dementia.
- Solvents and aluminum were found hazard for dementia, while silicon might be benefit.
- Living in rural area and more deprived neighborhood were adverse for dementia.
- Residential greenness and social contacts were proposed as beneficial factors.

Abstract

Background: Dementia is a challenging neurodegenerative disease. This systematic review aimed to summarize natural, physical, and social environmental factors that are associated with age-related cognitive impairment and dementia.

Methods: We systematically searched PubMed, EMBASE, Web of Science, and PsychINFO till January 11, 2021 for observational studies. The hazard ratio (HR), relative risk (RR), and odds ratio (OR) with 95% confidence interval (CI) were aggregated using random-effects methods. The quality of evidence for each association was evaluated.

Results: Of the 48,399 publications identified, there were 185 suitable for review across 44 environmental factors. Meta-analyses were performed for 22 factors. With high-to-moderate quality of evidence, risks were suggested in exposure to PM_{2.5} (HR=1.24, 95%CI: 1.17-1.31), NO₂ (HR=1.07, 95%CI: 1.02-1.12), aluminum (OR=1.35, 95%CI: 1.14-1.59), solvents (OR=1.14, 95%CI: 1.07-1.22), road proximity (OR=1.08, 95%CI: 1.04-1.12) and other air pollution, yet more frequent social contact (HR=0.82, 95%CI: 0.76-0.90) and more greenness (OR=0.97, 95%CI: 0.95-0.995) were protective. With low-to-very low quality, electromagnetic fields, pesticides, SO₂, neighborhood socioeconomic status, and rural living were suggested risks, but more community cultural engagement might be protective. No significant associations were observed in exposure to PM₁₀, NO_x, noise, silicon, community group, and temperature. For the remaining 22 factors, only a descriptive analysis was undertaken as too few studies or lack of information.

Conclusions: This review highlights that air pollution, especially PM_{2.5} and NO₂ play important role in the risk for age-related cognitive impairment and dementia. <https://pubmed.ncbi.nlm.nih.gov/34755643/>

Excerpts

3.2.2. Electromagnetic fields (EMFs)

From 27 studies, we found electromagnetic fields exposure was associated with dementia (OR=1.27, 95% CI: 1.13–1.43, I² =76.7%, S- grade, Fig. 3) with low-quality evidence, and among them, 11 were cohort studies and 16 were case-control studies (Appendix Figure 1.7). The result was robust via sensitivity analysis (Appendix Figure 3.8). Moreover, 23 studies focused on the outcome of AD, and 4 focused on VaD. Subgroup analysis displayed a significant association between EMFs with AD, but not with VaD (Fig. 3). In addition, the correlation with AD was observed in both females (OR=2.33, 95% CI: 1.19–4.57, I² =49.8%) and male (OR=1.51, 95% CI: 1.21–1.89, I² =46.4%, Appendix Figure 1.7.4). EMFs Exposure was mostly from occupation according to job- exposure information, and living next to the power line in life was not linked to dementia or AD (Fig. 3).

THE ECOLOGY OF ELECTRICITY AND ELECTRORECEPTION

Sam J. England, Daniel Robert. The ecology of electricity and electroreception. *Biological Reviews*. 12 October 2021. <https://doi.org/10.1111/brv.12804>

Abstract

Electricity, the interaction between electrically charged objects, is widely known to be fundamental to the functioning of living systems. However, this appreciation has largely been restricted to the scale of atoms, molecules, and cells. By contrast, the role of electricity at the ecological scale has historically been largely neglected, characterised by punctuated islands of research infrequently connected to one another. Recently, however, an understanding of the ubiquity of electrical forces within the natural environment has begun to grow, along with a realisation of the multitude of ecological interactions that these forces may influence. Herein, we provide the first comprehensive collation and synthesis of research in this emerging field of electric ecology. This includes assessments of the role electricity plays in the natural ecology of predator–prey interactions, pollination, and animal dispersal, among many others, as well as the impact of anthropogenic activity on these systems. A detailed introduction to the ecology and physiology of electroreception – the biological detection of ecologically relevant electric fields – is also provided. Further to this, we suggest avenues for future research that show particular promise, most notably those investigating the recently discovered sense of aerial electroreception.

Open access paper: <https://onlinelibrary.wiley.com/doi/full/10.1111/brv.12804>

ASSESSMENT OF HUMAN EXPOSURE (INCLUDING INTERFERENCE TO IMPLANTABLE DEVICES) TO LOW-FREQUENCY ELECTROMAGNETIC FIELD IN MODERN MICROGRIDS, POWER SYSTEMS AND ELECTRIC TRANSPORTS

Andrea Mariscotti. *Energies* 2021. 14(20): 6789. doi:10.3390/en14206789.

Abstract

Electromagnetic field emissions of modern power systems have increased in complexity if the many power conversion forms by means of power electronics and static converters are considered. In addition, the installed electric power has grown in many everyday applications such as wireless charging of vehicles, home integrated photovoltaic systems, high-performance electrified transportation systems, and so on. Attention must then be shifted to include harmonics and commutation components on one side, as well as closer interaction with humans, that concretizes in impact on physiological functions and interference to implantable medical devices and hearing aids. The panorama is complex in that standards and regulations have also increased significantly or underwent extensive revisions in the last 10 years or so. For assessment, the straightforward application of the limits of exposure is hindered by measurement problems (time or frequency domain methods, positioning errors, impact of uncertainty) and complex scenarios of exposure (multiple sources, large field gradient, time- varying emissions). This work considers thus both the clarification of the principles of interaction for each affected system (including humans) and the discussion of the large set of related normative and technical documents, deriving a picture of requirements and constraints. The methods of assessment are discussed in a metrological perspective using a range of examples. Open access paper: <https://www.mdpi.com/1996-1073/14/20/6789/htm>

ON THE POWER LINES—ELECTROMAGNETIC SHIELDING USING MAGNETIC STEEL LAMINATES

Tatiana Damatopoulou, Spyros Angelopoulos, Christos Christodoulou, Ioannis Gonos, Evangelos Hristoforou and Antonios Kladas. On the Power Lines—Electromagnetic Shielding Using Magnetic Steel Laminates. *Energies* 2021, 14(21): 7215. 2 November 2021. doi:10.3390/en14217215 (This article belongs to the Special Issue Advanced Electrical Measurements Technologies)

Abstract

Protection against the electromagnetic fields around high-voltage transmission lines is an issue of great importance, especially in the case of buildings near power lines. Indeed, the developed fields can be harmful for the habitants and electrical/electronic devices, so the implementation of appropriate measures to address the above electromagnetic interference issue is necessary in order to ensure the safety of both human beings and equipment. Several practices have been proposed to reduce the electric and the magnetic fields around overhead and underground transmission lines (minimum distance, shielded cables, anechoic chamber etc.). In this context, the scope of the current paper is the use of highly permeable magnetic sheets for shielding purposes, along with the development of an appropriate procedure, based on finite element analysis (FEA) for the efficient design of passive shielding. The simulation results are compared with laboratory measurements in order to confirm the adequacy of the proposed methodology. The good agreement between the FEA outcomes and the experimental results confirms that the developed FEA tool can be trustfully used for the design of the shielding means in the case of overhead or underground power lines.

Open access paper: <https://www.mdpi.com/1996-1073/14/21/7215/htm>

INFLUENCE OF MATERIALS, WINDOWS & SHIELDING LAYERS ON LOW-FREQUENCY ELECTROMAGNETIC ENVIRONMENT OF SUBWAY VEHICLE & HUMAN EXPOSURE RESEARCH

Zhiyuan Wang, Weinan Liu. Influence of Materials, Windows and Shielding Layers on Low-Frequency Electromagnetic Environment of Subway Vehicle and Human Exposure Research. *Progress In Electromagnetics Research M*, 105:151-160, 2021. doi:10.2528/PIERM21080403.

Abstract

The numerous high-power devices and cables gathered around the subway vehicle will aggravate the deterioration of the electromagnetic environment, which may cause the train to fail to operate normally or threaten the health of passengers with a pacemaker or defibrillator. In order to study the distribution characteristics of low-frequency magnetic field of the subway in complex electromagnetic environment and the influence of various factors on human electromagnetic exposure, the magnetic flux density nephograms of the subway train with different vehicle body materials, with or without windows and with the shielding layer are calculated and analyzed. Specific energy absorption rate (SAR) values have been calculated in a standing voxel model from exposure to electromagnetic fields at 2.4 GHz, frequencies commonly used by Wi-Fi devices. The numerical results show that the average value of magnetic flux density in the stainless-steel carriage is less than that in the aluminum alloy carriage and the carbon fiber reinforce plastic (CFRP) carriage. Compared with the vehicle with windows, the average value of magnetic flux density in the vehicle without windows is less. The added shielding layer decreases the average value of magnetic flux

density from 10.5 uT to 3 uT. The maximum value of magnetic flux density in the carriage under different factors is about 10 uT, which is far less than the magnetic flux density reference limit of 0.1 mT of the International Commission of Non-Ionizing Radiation Protection (ICNIRP) standard. When the Wi-Fi device is closest to the human body, the highest Specific Absorption Ratio (SAR) value of human tissue is 0.00749 W/kg, which is far less than the electromagnetic exposure limit of 1.6 W/kg of IEEE standard.

<http://www.jpier.org/PIERM/pier.php?paper=21080403>

EFFECTIVENESS OF OMEGA-3 FATTY ACID IN PROTECTION FROM MAGNETIC FIELD EXPOSURE DURING PREGNANCY: AN EXPERIMENTAL RAT STUDY

Ugurkan Erkayiran, Tufan Arslanca. Effectiveness of Omega-3 Fatty Acid in Protection from Magnetic Field Exposure During Pregnancy: An Experimental Rat Study. Journal of Reproductive Medicine. Nov 2021.

OBJECTIVE: Magnetic fields (MFs) can have several negative effects on the human body. This study aimed to reveal the relationship of adenosine deaminase enzyme (ADA), thyroid-stimulating hormone (TSH), and fetal neural degeneration (FND) with the curative effect of omega-3 fatty acids.

STUDY DESIGN: Pregnant Wistar Albino rats (n=30) were separated into 3 groups: Group 1 (sham group), Group 2 (rats exposed to 20 microteslas [μ T] for 1 h/day for 10 days), and Group 3 (rats exposed to 20 μ T for 4 h/day for 10 days). Blood samples were taken on days 1, 10, and 17 to examine ADA and TSH. Neurodegeneration scores were analyzed with microscopic examination.

RESULTS: FND was determined to be greater in the rats exposed to the long-term MFs as compared to short-term MFs ($p < 0.05$). There was a significant difference in terms of ADA and TSH in the rats that received short- and long-term MF as compared to the sham group ($p < 0.05$). The mean scores were higher in the short- and long-term MF groups than in the sham group ($p < 0.05$). ADA activity was found to be higher and TSH was lower in the short- and long-term MF groups ($p < 0.05$). With the use of omega-3, positive alterations in myelin degeneration were observed in the groups exposed to short- and long-term MF ($p < 0.05$).

CONCLUSION: While both ADA and TSH increased in both short- and long-term MF exposure, long-term MFs affected FND more than short-term exposure. Although using omega-3 showed a protective effect on the MF-induced FND, it was only protective against short-term exposure.

https://www.reproductivemedicine.com/toc/auto_abstract.php?id=24943

Exposure to 10 Hz Pulsed Magnetic Fields Do Not Induce Cellular Senescence in Human Fetal Lung Fibroblasts

Sun Chuan, Huang Zheng, Qin Houbing, Zhang Jing, Wang Sanying, Xu Xiaogang, Ying Shibo, Mao Genxiang. Exposure to 10 Hz Pulsed Magnetic Fields Do Not Induce Cellular Senescence in Human Fetal Lung Fibroblasts. *Frontiers in Public Health*. 9:1657. 2021. doi: 10.3389/fpubh.2021.761069.

Abstract

Rapid population aging has led to a global burden of late-life diseases. As the largest risk factor for a multitude of age-related diseases, aging is not only the result of genotype but also closely related to external factors. With the rapid expansion in the usage of electromagnetic fields (EMFs), the effect of EMFs on aging has also attracted attention. Cells are the basic unit of organs and body tissues, and cellular senescence plays an important role in the aging process. The effect of EMFs on cellular senescence has been investigated in a few studies, but the information is limited, and the results are inconsistent; thus, further investigation is required. In this study, we investigated the effect of 10 Hz pulsed magnetic fields (MFs) on cellular senescence in a 2BS cell line, isolated from human fetal lung fibroblasts, and found that intermittent (1 d on/1 d off) exposure to 10 Hz pulsed MFs at 1.0 mT for 2 weeks induced DNA damage, but no other significant phenotype of cellular senescence in 2BS cells.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.761069>

THE IMPACT OF INTRASPECIES VARIABILITY ON GROWTH RATE AND CELLULAR METABOLIC ACTIVITY OF BACTERIA EXPOSED TO ROTATING MAGNETIC FIELD

Woroszyło M, Ciecholewska-Juśko D, Junka A, Pruss A, Kwiatkowski P, Wardach M, Fijałkowski K. The Impact of Intraspecies Variability on Growth Rate and Cellular Metabolic Activity of Bacteria Exposed to Rotating Magnetic Field. *Pathogens*. 2021 Nov 4;10(11):1427. doi: 10.3390/pathogens10111427.

Abstract

Majority of research on the influence of magnetic fields on microorganisms has been carried out with the use of different species or different groups of microorganisms, but not with the use of different strains belonging to one species. The purpose of the present study was to assess the effect of rotating magnetic fields (RMF) of 5 and 50 Hz on the growth and cellular metabolic activity of eight species of bacteria: *Staphylococcus aureus*, *Pseudomonas aeruginosa*, *Proteus mirabilis*, *Klebsiella pneumoniae*, *Enterococcus faecalis*, *Enterobacter cloacae*, *Moraxella catarrhalis*, and *Bacillus cereus*. However, contrary to the research conducted so far, each species was represented by at least four different strains. Moreover, an additional group of *S. aureus* belonging to a single clonal type but representing different biotypes was also included in the experiment. The results showed a varied influence of RMF on growth dynamics and cellular metabolic activity, diversified to the greatest extent in dependence on the bacterial strain exposed to the RMF and to a lesser extent in dependence on the frequency of the generated magnetic field. It was found that, with regard to the exposed strain of the same species, the effect exerted by the RMF may be positive (i.e., manifests as the increase in the growth rate or/and cellular metabolic activity) or negative (i.e., manifests as a reduction of both aforementioned features) or none. Even when one clonal type of *S. aureus* was used, the results of RMF exposure also varied (although the degree of differentiation was lower than for strains representing different clones). Therefore,

the research has proven that, apart from the previously described factors related primarily to the physical parameters of the magnetic field, one of the key parameters affecting the final result of its influence is the bacterial intraspecies variability.

Open access paper: <https://www.mdpi.com/2076-0817/10/11/1427>

THE CONSPIRACY OF COVID-19 AND 5G: SPATIAL ANALYSIS FALLACIES IN THE AGE OF DATA DEMOCRATIZATION

Eoin Flaherty, Tristan Sturm, Elizabeth Farries. The conspiracy of Covid-19 and 5G: Spatial analysis fallacies in the age of data democratization. *Social Science & Medicine*. 2021, 114546. 2021. doi: 10.1016/j.socscimed.2021.114546.

Abstract

In a context of mistrust in public health institutions and practices, anti-COVID/vaccination protests and the storming of Congress have illustrated that conspiracy theories are real and immanent threat to health and wellbeing, democracy, and public understanding of science. One manifestation of this is the suggested correlation of COVID-19 with 5G mobile technology. Throughout 2020, this alleged correlation was promoted and distributed widely on social media, often in the form of maps overlaying the distribution of COVID-19 cases with the instillation of 5G towers. These conspiracy theories are not fringe phenomena, and they form part of a growing repertoire for conspiracist activist groups with capacities for organised violence. In this paper, we outline how spatial data have been co-opted, and spatial correlations asserted by conspiracy theorists. We consider the basis of their claims of causal association with reference to three key areas of geographical explanation: (1) how social properties are constituted and how they exert complex causal forces, (2) the pitfalls of correlation with spatial and ecological data, and (3) the challenges of specifying and interpreting causal effects with spatial data. For each, we consider the unique theoretical and technical challenges involved in specifying meaningful correlation, and how their discarding facilitates conspiracist attribution. In doing so, we offer a basis both to interrogate conspiracists' uses and interpretation of data from elementary principles and offer some cautionary notes on the potential for their future misuse in an age of data democratization. Finally, this paper contributes to work on the basis of conspiracy theories in general, by asserting how – absent an appreciation of these key methodological principles – spatial health data may be especially prone to co-option by conspiracist groups.

<https://www.sciencedirect.com/science/article/pii/S0277953621008789>

SOLAR ACTIVITY IS ASSOCIATED WITH DIASTOLIC AND SYSTOLIC BLOOD PRESSURE IN ELDERLY ADULTS

Veronica A Wang, Carolina L Zilli Vieira, Eric Garshick, Joel D Schwartz, Michael S Garshick, Pantel Vokonas, Petros Koutrakis. Solar Activity Is Associated With Diastolic and Systolic Blood Pressure in Elderly Adults. *J Am Heart Assoc*. 2021 Nov 2;10(21):e021006. doi: 10.1161/JAHA.120.021006.

Abstract

Background Since solar activity and related geomagnetic disturbances modulate autonomic nervous system activity, we hypothesized that these events would be associated with blood pressure (BP).

Methods We studied 675 elderly men from the Normative Aging Study (Boston, MA) with 1949 BP measurements between 2000 and 2017. Mixed-effects regression models were used to investigate the association of average 1-day (ie, day of BP measurement) to 28-day interplanetary magnetic field intensity, sunspot number, and a dichotomized measure of global geomagnetic activity (Kp index) in 4-day increments with diastolic and systolic BP. We adjusted for meteorological conditions and other covariates associated with BP, and in additional models adjusted for ambient air pollutants (particulate matter with an aerodynamic diameter ≤ 2.5 μm , black carbon, and particle number) and ambient particle radioactivity.

Results There were positive associations between interplanetary magnetic field, sunspot number, and Kp index and BP that were greatest with these exposures averaged over 16 through 28 days before BP measurement. An interquartile range increase of 16-day interplanetary magnetic field and sunspot number and higher Kp index were associated with a 2.5 (95% CI, 1.7–3.2), 2.8 (95% CI, 2.1–3.4), and 1.7 (95% CI, 0.8–2.5) mm Hg increase, respectively, for diastolic BP as well as a 2.1 (95% CI, 0.7–3.6), 2.7 (95% CI, 1.5–4.0), and 0.4 (95% CI, -1.2 to 2.1) mm Hg increase, respectively, for systolic BP. Associations remained after adjustment for ambient air pollutants and ambient particle radioactivity.

Conclusions Solar activity and solar-driven geomagnetic disturbances were positively associated with BP, suggesting that these natural phenomena influence BP in elderly men.

Clinical Perspective

What Is New?

- Solar and geomagnetic activity were associated with increases in blood pressure in a large cohort of predominantly White, elderly men.
- The association with blood pressure was similar to or greater than that of particulate pollution and of radioactivity associated with ambient particles.
- The association of solar and geomagnetic activity with blood pressure was independent of these pollutants.

What Are the Clinical Implications?

- These findings suggest that natural phenomena linked to the solar cycle contributes to increases in blood pressure and, therefore, may influence hypertension management.

The Earth's magnetic field protects living organisms from long-term, harmful extra-terrestrial radiation. Despite this protective shield, solar activity can cause geomagnetic disturbances (GMD), disruptions to the Earth's natural magnetic field oscillations, and can impact autonomic nervous system activities,^{1, 2} which can, in turn, directly and indirectly play a role in initiating and sustaining high blood pressure (BP).³ Numerous pathogenic risk factors such as genetic predisposition,⁴ physical activity,⁵ and diet⁶ have been identified to play key roles in the development of hypertension. A recent review⁷ highlighted the role of environmental factors, such as temperature, altitude, latitude, and air pollutants, in elevating BP, but few studies considered solar activity and GMD as risk factors for the development of hypertension

or transient increases in BP. In those that have,8, 9 the findings suggest that individuals have elevated BP several days before and after magnetic storms.

To gain insight and provide awareness into the association between solar activity and BP in elderly men, a vulnerable population at high risk for cardiovascular disease,10 we conducted a repeated measures analysis to examine the association of average 1-day (ie, day of BP measurement) to 28-day solar activity and GMD with BP among elderly men in Boston, MA who had between 1 and 8 health assessments over 17 years. We hypothesized that solar and geomagnetic activity is positively associated with systolic blood pressure (SBP) and diastolic blood pressure (DBP).

Open access paper: <https://www.ahajournals.org/doi/10.1161/JAHA.120.021006>

ASSESSMENT OF HUMAN EXPOSURE TO ELECTROMAGNETIC FIELDS: REVIEW AND FUTURE DIRECTIONS

Akimasa Hirata, Yinliang Diao, Teruo Onishi, Kensuke Sasaki, Seunyoung Ahn, Davide Colombi, Valerio De Santis, Ilkka Laakso, Luca Giaccone, Wout Joseph, Essam A. Rashed, Wolfgang Kainz, Ji Chen. Assessment of Human Exposure to Electromagnetic Fields: Review and Future Directions. IEEE Transactions on Electromagnetic Compatibility, doi: 10.1109/TEMPC.2021.3109249. Wolfgang Kainz, Center for Devices and Radiological Health, Food and Drug Administration, Silver Spring, MD 20852 USA (e-mail: wolfgang.kainz@fda.hhs.gov).

Abstract

This article reviews recent standardization activities and scientific studies related to the assessment of human exposure to electromagnetic fields (EMF). The differences of human exposure standards and assessment of consumer products and medical applications are summarized. First, we reviewed human body modeling and tissue dielectric properties. Then, we explain the rationale of current exposure standards from the viewpoint of EMF and the standardization process for product compliance based on these exposure standards. The assessment of wireless power transfer, as an example of emerging wireless devices, and environmental EMFs in our daily lives are reviewed. Safety in magnetic resonance systems, where the EMF exposure is much larger than from typical consumer devices, is also reviewed. Finally, we summarize future research directions and research needs for EMF safety.

Excerpts

The overall improvement in human model accuracy is an important topic as mentioned in Section II-A and B. This may include an intersubject variability study using different models or an analysis of the population level. In addition, the accurate measurement of dielectric human tissue properties is essential for evaluating the induced field strength. Recent studies suggest that the conductivity of skin and brain tissues may be significantly higher than those commonly used in LF dosimetry (e.g., [31]). Recent publications reported a novel approach for dielectric measurements that allows the estimation of dielectric human tissue properties in vivo using MRI [137]. These and other novel medical imaging and measurement technologies will help to further improve human body modeling.

Research needs in the areas of human EMF safety (Section III) are summarized by various international standardization bodies [4], [138], [139]. The topics listed therein relate to

human body modeling, as well as the necessity of more accurate dosimetry, which includes reductions of uncertainties, and multiphysics or multiscale methods for correlating the field quantities with substantiated biological effects (e.g., [140]). The assessments of exposure to emerging LF and RF technologies and simultaneous exposure (local and whole-body) to multiple sources will also prove useful for identifying potential changes in the dominant factors affecting EM safety.

Additional research on human EM safety will provide more scientific data and improve the justifications for exposure limits, especially at frequencies higher than 6 GHz, and in the intermediate frequency range.

New mobile wireless technologies, such as 5G/6G and beyond, are being developed continuously. Their safe use demands not only conservative short- and long-term exposure limits for the new frequencies, but also adequately standardized compliance methods. Therefore, additional relevant exposure evaluation methods are needed to support frequencies from 100 to 300 GHz within the next 10 years. As the complexity of new wireless technologies has substantially increased over the years, further development and standardization of novel and more efficient EMF compliance testing methodologies are needed. International standardization will continue to play a key role in harmonizing global test procedures to establish compliance with current and future wireless technologies. One challenge faced by standardization will be to define test conditions that provide conservative, but realistic exposure conditions based on real-life situations, rather than relying on theoretical or unreasonable assumptions and simplifications. The assessment methods by the computation may become more common for emerging wireless systems, and thus further research is needed for verification of its effectiveness in different scenarios.

Future work for the environmental spectral measurement (see Section V) will focus on measurement procedures to assess worst-cases and realistic exposure to EMF from 5G New Radio (NR)—Ma-MIMO with a focus on sub-6 GHz and millimeter wave bands. The literature scarcely documents in situ assessment of 5G Ma- MIMO signals at millimeter waves and the proper adjustment of measurement settings. For personal exposimeter studies, protocols for 5G personal exposure assessment will need to include active and inactive users. In contrast to legacy technologies, Ma-MIMO base station exposure strongly depends on the activity of the individual user (e.g., streaming data, downloading files, and web browsing). Involving millimeter waves for personal exposure assessment is a significant challenge.

Also, in RF-MR safety, it is necessary to develop clinically relevant human body models for measurements in addition to developing new computational methods for multiscale and multiphysics applications. Most computational human body models use over 30 different types of tissue. However, below 500 MHz dielectric human tissues properties are not significantly different from each other. Some preliminary studies provided evidence that simplified human body models can still achieve accurate EM simulation results. This leads to the possibility of developing simplified computational and experimental models for different RF-MR safety measurement needs for PIMDs, AIMDs, and high-field MR systems (e.g., [141]). Furthermore, simple and equivalent test methods/equipment those are based on the physics principles should be developed (e.g., [142]).

<https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9557785&isnumber=4358749>

ACCURATELY ASSESSING EXPOSURE TO RADIO FREQUENCY ELECTROMAGNETIC FIELDS FROM 5G NETWORKS

Accurately assessing exposure to radio frequency electromagnetic fields from 5G networks. Ericsson white paper. GFTL-21:001105 Uen. October 2021.

Excerpts

"Measurements were recently taken in a large number of street-level locations around base stations in a commercial 5G network with massive MIMO base stations operating in the 3.5GHz band [9]. It was found that the contribution from the 5G network to the total environmental RF EMF exposure was less than 10 percent even in the case of 100 percent induced traffic and that the maximum exposure levels from the 5G base stations were 150 to 200 times below the international limits set by the ICNIRP."

Conclusion

"The high spectrum efficiency and the advanced antenna technologies used by 5G NR lead to lower levels of RF EMF exposure than from earlier generations of mobile networks for comparable services. The base stations in 5G NR networks need to comply with the same RF EMF safety regulations as other radio equipment, and the limits cover all frequency bands used by 5G, including those in the millimeter-wave range. International RF EMF exposure guidelines have recently been published based on the latest available scientific research, with conservative limits that have been or will be adopted in national regulations. Assessing compliance with RF EMF limits may be a challenge for massive MIMO 5G base stations due to dynamic beam steering, but solutions such as envelope beam pattern files together with recommended power reduction factors are available to enable accurate evaluations. The typical overall environmental RF EMF exposure will remain at a small fraction of international limits even with 5G being deployed since the contribution from 5G is relatively small."

Open access report: <https://www.ericsson.com/en/reports-and-papers/white-papers/accurately-assessing-exposure-to-radio-frequency-electromagnetic-fields-from-5g-networks>

PHYSICAL UNITS TO REPORT INTENSITY OF ELECTROMAGNETIC WAVE

R Ramirez-Vazquez, I Escobar, T Franco, E Arribas. Environ Res. 2021 Nov 2;112341. doi: 10.1016/j.envres.2021.112341.

Abstract

The aim of this work is to propose a consensus to scientific community that handles personal exposimeters, which measure intensity of an electromagnetic wave (W/m^2). To express the intensity of an electromagnetic wave there is a duality in the way of expressing it. Some scientists prefer to use W/m^2 while others use V/m , which is a unit of the electric field. There is also a duality in the name, sometimes it is called it power flux density and some other times, wave intensity. We believe that this second name is more appropriate from the point of view of physics. We suggest expressing intensity of an electromagnetic wave in W/m^2 instead of giving the value of their electric field which is measured in V/m . There is a quadratic relation between electric field and intensity of the wave, and it is necessary to do a mathematical operation, so in our opinion, it is preferable to use W/m^2 which directly gives us the value of the measured intensity. Furthermore, if the intensity is very low, it may be expressed in

μW/m² and with only three significant figures, due to sensitivity of the current exposimeters used. <https://pubmed.ncbi.nlm.nih.gov/34740620/>

THE EFFECTS OF RADIOFREQUENCY EXPOSURE ON MALE FERTILITY AND ADVERSE REPRODUCTIVE OUTCOMES: A PROTOCOL FOR TWO SYSTEMATIC REVIEWS OF HUMAN OBSERVATIONAL STUDIES WITH META-ANALYSIS

Kenny RPW, Millar EB, Adesanya A, Richmond C, Beyer F, Calderon C, Rankin J, Toledano M, Feychting M, Pearce MS, Craig D, Pearson F. The effects of radiofrequency exposure on male fertility and adverse reproductive outcomes: A protocol for two systematic reviews of human observational studies with meta-analysis. *Environ Int.* 2021 Nov 1;158:106968. doi: 10.1016/j.envint.2021.106968.

Abstract

Background: The World Health Organization (WHO) is bringing together evidence on radiofrequency electromagnetic field (RF-EMF) exposure in relation to health outcomes, previously identified as priorities for evaluation by experts in the field, to inform exposure guidelines. A suite of systematic reviews are being undertaken by a network of topic experts and methodologists in order to collect, assess and synthesise data relevant to these guidelines. Here, we present the protocol for the systematic review on the effect of exposure to RF on adverse reproductive outcomes (human observational studies), also referred to as Systematic Review (SR) 3 within the series of systematic reviews currently being commissioned.

Objectives: Following the WHO handbook for guideline development and the COSTER conduct guidelines, we will systematically review the effect of RF-EMF exposure on both male fertility (SR3A) and adverse pregnancy outcomes (SR3B) in human observational studies. Herein we adhere to the PRISMA-P reporting guidelines.

Data sources: We will conduct a broad search for potentially relevant records relevant for both reviews within the following bibliographic databases: MEDLINE; Embase; and EMF Portal. We will also conduct searches of grey literature through relevant databases and organisational websites. RF-EMF experts will also be consulted. We will hand search citation and reference lists of included study records.

Study eligibility criteria: We will include quantitative human observational studies on the effect of RF-EMF exposure: (in SR3A) in adult male participants on infertility, sperm morphology, concentration or total sperm count or motility; and (in SR3B) in preconception adults or pregnant women on preterm birth, small for gestational age (associated with intrauterine growth restriction), miscarriage, stillbirth and congenital anomalies.

Study appraisal and synthesis methods: Titles, abstracts and then full texts will be screened in blinded duplicate against eligibility criteria with input from a third reviewer as required. Data extraction from included studies will be completed by two reviewers as will risk of bias assessment using the Office of Health Assessment and Translation (OHAT) tool. If appropriate we will undertake meta-analysis to pool effect measures and explore heterogeneity using sub-group analyses or meta-regression as feasible. We will conduct sensitivity analysis to assess the impact of any assumptions made throughout the review process. The OHAT

methodology, based on the GRADE guidelines for evidence assessment, will be used to evaluate the certainty of evidence per outcome and to conclude the level of evidence of a health effect.

Conclusion: This manuscript details the protocols for two systematic reviews. The aims of publishing details of both protocols are to: pre-specify their scope and methods; reduce the impact of reviewer bias; promote transparency and replicability; and improve the review process.

Prospero registration: CRD42021265401 (SR3A), CRD42021266268 (SR3B).

<https://pubmed.ncbi.nlm.nih.gov/34735951/>

THE EFFECTS OF RADIOFREQUENCY ELECTROMAGNETIC FIELDS EXPOSURE ON HUMAN SELF-REPORTED SYMPTOMS: A PROTOCOL FOR A SYSTEMATIC REVIEW OF HUMAN EXPERIMENTAL STUDIES

Bosch-Capblanch X, Esu E, Dongus S, Oringanje CM, Jalilian H, Evers J, Oftedal G, Meremikwu M, Rösli M. The effects of radiofrequency electromagnetic fields exposure on human self-reported symptoms: A protocol for a systematic review of human experimental studies. *Environ Int.* 2021 Nov 1;158:106953. doi: 10.1016/j.envint.2021.106953.

Abstract

Background: The technological applications of radiofrequency electromagnetic fields (RF-EMF) have been steadily increasing since the 1950s across multiple sectors exposing large proportions of the population. This fact has raised concerns related to the potential consequences to people's health. The World Health Organization (WHO) is assessing the potential health effects of exposure to RF-EMF and has carried out an international survey amongst experts, who have identified six priority topics to be further addressed through systematic reviews, whereof the effects on symptoms is one of them. We report here the systematic review protocol of experimental studies in humans assessing the effects of RF-EMF on symptoms.

Objective: Our objectives are to assess the effects of exposure to electromagnetic fields (compared to no or lower exposure levels) on symptoms in human subjects. We will also assess the accuracy of perception of presence of exposure in volunteers with and without idiopathic environmental intolerance attributed to electromagnetic fields (IEI-EMF).

Eligibility criteria: We will search relevant literature sources (e.g. the Web of Science, Medline, Embase, Epistemonikos) for randomized trials (comparing at least two arms) and randomised crossover trials of RF-EMF exposure that have assessed the effects on symptoms. We will also include studies that have measured the accuracy of the perception of the presence or absence of exposure. We will include studies in any language.

Study appraisal and synthesis: Studies will be assessed against inclusion criteria by two independent reviewers. Data on study characteristics, participants, exposure, comparators and effects will be extracted using a specific template for this review, by two independent reviewers. Discrepancies will be solved by consensus. Risk of bias

(ROB) will be assessed using the ROB Rating Tool for Human and Animal Studies and the level of confidence in the evidence of the exposure-outcome relations will be assessed using the GRADE approach. For the perception studies, we will use adapted versions of the ROB tool and GRADE assessment. Where appropriate, data will be combined using meta-analytical techniques. <https://pubmed.ncbi.nlm.nih.gov/34735955/>

RADIO-FREQUENCY EXPOSURE OF THE YELLOW FEVER MOSQUITO (A. AEGYPTI) FROM 2 TO 240 GHZ

De Borre E, Joseph W, Aminzadeh R, Müller P, Boone MN, Josipovic I, et al. (2021) Radio-frequency exposure of the yellow fever mosquito (*A. aegypti*) from 2 to 240 GHz. *PLoS Comput Biol* 17(10): e1009460. doi: 10.1371/journal.pcbi.1009460.

Abstract

Fifth generation networks (5G) will be associated with a partial shift to higher carrier frequencies, including wavelengths of insects. This may lead to higher absorption of radio frequency (RF) electromagnetic fields (EMF) by insects and could cause dielectric heating. The yellow fever mosquito (*Aedes aegypti*), a vector for diseases such as yellow and dengue fever, favors warm climates. Being exposed to higher frequency RF EMFs causing possible dielectric heating, could have an influence on behavior, physiology and morphology, and could be a possible factor for introduction of the species in regions where the yellow fever mosquito normally does not appear. In this study, the influence of far field RF exposure on *A. aegypti* was examined between 2 and 240 GHz. Using Finite Difference Time Domain (FDTD) simulations, the distribution of the electric field in and around the insect and the absorbed RF power were found for six different mosquito models (three male, three female). The 3D models were created from micro-CT scans of real mosquitoes. The dielectric properties used in the simulation were measured from a mixture of homogenized *A. aegypti*. For a given incident RF power, the absorption increases with increasing frequency between 2 and 90 GHz with a maximum between 90 and 240 GHz. The absorption was maximal in the region where the wavelength matches the size of the mosquito. For a same incident field strength, the power absorption by the mosquito is 16 times higher at 60 GHz than at 6 GHz. The higher absorption of RF power by future technologies can result in dielectric heating and potentially influence the biology of this mosquito.

Author summary

Radio Frequency (RF) exposure of the *A. aegypti* mosquito can lead to absorption and dielectric heating. We used Finite Difference Time Domain (FDTD) simulations between 2 and 240 GHz to study the RF power absorbed by the insect and the distribution of the electric field (EF) in and around it. For this, three male and three female mosquito 3D models were constructed from micro-CT scans. We used high resolution models and dielectric properties, both retrieved from real insects, to gain realistic outputs. For increasing frequency up to 90 GHz, the absorbed power increases for all models. At 90–120 GHz, the wavelength is comparable to the body size, and the increase in absorbed powers reaches a maximum. Therefore, moving to higher frequencies in 5G, implies higher absorbed power and possibly higher dielectric heating of the insect.

Excerpt

The insect of interest in this paper is the yellow fever mosquito, *Aedes aegypti*, it is known as a vector for diseases such as yellow fever, dengue fever and zika virus infections [14, 15]. According to the Centers of Disease Control and Prevention in the U.S., yellow fever cases and deaths worldwide are estimated at 200,000 and 30,000 each year [16], respectively. The yellow fever mosquito is a tropical species favouring a hot and humid environment. Temperature affects the life cycle and feeding behaviour of the mosquito and the reproduction of the viruses [14, 15]. RF power absorption and dielectric heating can cause disturbance in for example the behaviour or development of the mosquito. Another interesting consequence of dielectric heating and higher body temperature, may be the spread of the mosquito to areas that are normally unfavorable for them. Be that as it may, the focus in this paper is on the RF power absorption, the dielectric heating or other consequences are not considered.

Open access paper:

<https://journals.plos.org/ploscompbiol/article?id=10.1371/journal.pcbi.1009460>

THE DETRIMENTAL EFFECT OF CELL PHONE RADIATION ON SPERM BIOLOGICAL CHARACTERISTICS IN NORMOZOOSPERMIC

Mohammadmehdi Hassanzadeh-Taheri, Mohammad Ali Khalili, Ali Hosseini Mohebbati, Mahmood Zardast, Mehran Hosseini, Maria Grazia Palmerini, Mohammad Reza Doostabadi. The detrimental effect of cell phone radiation on sperm biological characteristics in normozoospermic. *Andrologia*. 10 October 2021. doi: 10.1111/and.14257.

Abstract

Radiofrequency electromagnetic radiation emitted from cell phone has harmful effects on some organs of the body, such as the brain, heart, and testes. This study aimed to assess the effects of cell phones on sperm parameters, DNA fragmentation, and apoptosis in normozoospermic. Normal sperm samples were divided into two groups of control and case. The samples from the case were placed for 60 min at a distance of approximately 2.5 cm from the cell phone set in the active antenna position. Control samples were exposed to cell phones without active antennas. All specimens were analysed by World Health Organization criteria. Sperm viability, sperm with chromatin abnormality and maturity, DNA fragmentation, and apoptosis were examined. Viability and motility in the case were significantly lower than the control ($p < .001$, $p = .004$ respectively). The percentage of apoptotic sperms and DNA fragmentation were significantly higher in the case when compared with the control ($p = .031$, $p < .001$ respectively). The other parameters studied such as morphology, chromatin abnormality, and maturity showed no significant difference between the case and control groups. Cell phone waves had a detrimental effect on human sperm's biological features. Therefore, it is recommended to keep the cell phone away from the pelvis as much as possible.

Conclusion

The cell phone waves can reduce the sperm's biological characteristics, such as morphology, motility, viability, DNA integrity, and an increase in apoptosis in normozoospermic men. Therefore, it is recommended for men to keep the cell phone away from their pelvic.

<https://pubmed.ncbi.nlm.nih.gov/34628682/>

PRENATAL CHRONIC EXPOSURE TO ELECTROMAGNETIC FIELDS MODULATED ADENOSINE DEAMINASE ACTIVITY IN SERUM AND BRAIN OF WISTAR RATS' OFFSPRING

Behrooz Mohammadi, Mahdi Sadegh & Homa Soleimani. Prenatal chronic exposure to electromagnetic fields modulated adenosine deaminase activity in serum and brain of Wistar rats' offspring, *Journal of Microwave Power and Electromagnetic Energy*. 2021. DOI: 10.1080/08327823.2021.1993045

Abstract

Considering the importance of the embryonic stage, we investigated the brain and serum ADA of offspring chronically exposed to low and high frequency electromagnetic field (EMF-LF/HF) during their embryonic period. Male and female rats were randomly selected for mating. After mating, the females were divided into four groups. The first group was exposed to extremely EMF-LF (50 Hz) through a solenoid for 30 minutes per day, the second group was constantly exposed to EMF-HF (900 MHz) via a cell phone, the third group (sham) was placed inside solenoid without any exposure, and the last group (control) was placed in their cage. The brain and serum samples of offspring (6 females and 6 males in each group) were collected four weeks after birth. ADA was measured by a specific enzymatic kit and a spectrophotometer. Exposure to EMF-LF (1.5 mT) was significantly increased the brain ADA activity in both male and female offspring. In addition, long-term HF exposure significantly enhanced serum ADA activity in males in compared with control group but, the brain ADA level was increased significantly in both genders ($p < 0.01$). The present study revealed alterations in the serum and brain ADA following embryonic exposure to HF and ELF. These alterations depended on gender, frequency and wave intensity. <https://www.tandfonline.com/doi/abs/10.1080/08327823.2021.1993045?journalCode=tpe>
[e20](#)

THE EFFECT OF MOBILE RADIATION ON THE OXIDATIVE STRESS BIOMARKERS IN PREGNANT MICE

Nargess Moghadasi, Iraj Alimohammadi, Ali Safari Variani, Azadeh Ashtarinezhad. The Effect of Mobile Radiation on the Oxidative Stress Biomarkers in Pregnant Mice. *J Family Reprod Health*. 2021 Sep;15(3):172-178. doi: 10.18502/jfrh.v15i3.7134.

Abstract

Objective: Due to the growing use of communication instruments such as cell phones and wireless devices, there is growing public concern about possible harmful effects, especially in sensitive groups such as pregnant women. This study aimed to investigate the oxidative stress induced by exposure to 900 MHz mobile phone radiation and the effect of vitamin C intake on reducing possible changes in pregnant mice.

Materials and methods: Twenty-one pregnant mice were divided into three groups (control, mobile radiation- exposed, and mobile radiation plus with vitamin C intake co-exposed (200 mg /kg)). The mice in exposure groups were exposed to 900 MHz, 2 watts, and a power density of 0.045 $\mu\text{W}/\text{cm}^2$ mobile radiation for eight hours/day for ten consecutive days. After five days of rest, MDA (Malondialdehyde), 8-OHdG (8-hydroxy-2' - deoxyguanosine), and TAC (Total Antioxidant Capacity) levels were measured in the blood of animals. The results were analyzed by SPSS.22.0 software.

Results: The results showed that exposure to mobile radiation increased MDA ($P=0.002$), and 8-OHdG ($P=0.001$) significantly and decreased Total Antioxidant Capacity in the exposed groups ($P=0.001$). Taking vitamin C inhibited the significant increase in MDA and 8-OHdG levels in exposed groups.

Conclusion: Although exposure to mobile radiation can cause oxidative stress in the blood of pregnant mice, vitamin C as an antioxidant can prevent it.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8536820/>

NO SIGNIFICANT EFFECTS OF CELLPHONE ELECTROMAGNETIC RADIATION ON MICE MEMORY OR ANXIETY: SOME MIXED EFFECTS ON TRAUMATIC BRAIN INJURED MICE

Doaa Qubty, Shaul Schreiber, Vardit Rubovitch, Amir Boag, Chaim G Pick. No Significant Effects of Cellphone Electromagnetic Radiation on Mice Memory or Anxiety: Some Mixed Effects on Traumatic Brain Injured Mice. *Neurotrauma Rep.* 2021 Aug 17;2(1):381-390. doi: 10.1089/neur.2021.0009.

Abstract

Current literature details an array of contradictory results regarding the effect of radiofrequency electromagnetic radiation (RF-EMR) on health, both in humans and in animal models. The present study was designed to ascertain the conflicting data published regarding the possible impact of cellular exposure (radiation) on male and female mice as far as spatial memory, anxiety, and general well-being is concerned. To increase the likelihood of identifying possible "subtle" effects, we chose to test it in already cognitively impaired (following mild traumatic brain injury; mTBI) mice. Exposure to cellular radiation by itself had no significant impact on anxiety levels or spatial/visual memory in mice. When examining the dual impact of mTBI and cellular radiation on anxiety, no differences were found in the anxiety-like behavior as seen at the elevated plus maze (EPM). When exposed to both mTBI and cellular radiation, our results show improvement of visual memory impairment in both female and male mice, but worsening of the spatial memory of female mice. These results do not allow for a decisive conclusion regarding the possible hazards of cellular radiation on brain function in mice, and the mTBI did not facilitate identification of subtle effects by augmenting them. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8550818/>

CELLULAR STRESS RESPONSE TO EXTREMELY LOW-FREQUENCY ELECTROMAGNETIC FIELDS (ELF-EMF): AN EXPLANATION FOR CONTROVERSIAL EFFECTS OF ELF-EMF ON APOPTOSIS (REVIEW)

Barati M, Darvishi B, Javidi MA, Mohammadian A, Shariatpanahi SP, Eisavand MR, Madjid Ansari A. Cellular stress response to extremely low-frequency electromagnetic fields (ELF-EMF): An explanation for controversial effects of ELF-EMF on apoptosis. *Cell Prolif.* 2021 Nov 6:e13154. doi: 10.1111/cpr.13154.

Abstract

Impaired apoptosis is one of the hallmarks of cancer, and almost all of the non-surgical approaches of eradicating tumour cells somehow promote induction of apoptosis. Indeed, numerous studies have stated that non-ionizing non-thermal extremely low-frequency magnetic fields (ELF-MF) can modulate the induction of apoptosis in exposed cells; however,

much controversy exists in observations. When cells are exposed to ELF- EMF alone, very low or no statistically significant changes in apoptosis are observed. Contrarily, exposure to ELF-EMF in the presence of a co-stressor, including a chemotherapeutic agent or ionizing radiation, can either potentiate or inhibit apoptotic effects of the co-stressor. In our idea, the main point neglected in interpreting these discrepancies is "the cellular stress responses" of cells following ELF-EMF exposure and its interplay with apoptosis. The main purpose of the current review was to outline the triangle of ELF-EMF, the cellular stress response of cells and apoptosis and to interpret and unify discrepancies in results based on it. Therefore, initially, we will describe studies performed on identifying the effect of ELF-EMF on induction/inhibition of apoptosis and enumerate proposed pathways through which ELF-EMF exposure may affect apoptosis; then, we will explain cellular stress response and cues for its induction in response to ELF-EMF exposure; and finally, we will explain why such controversies have been observed by different investigators.

Open access paper: <https://onlinelibrary.wiley.com/doi/10.1111/cpr.13154>

MELATONIN: A POTENTIAL SHIELD AGAINST ELECTROMAGNETIC WAVES

Maya Jammoul, Nada Lawand. Melatonin: a Potential Shield against Electromagnetic Waves. *Curr Neuropharmacol*. 2021 Jun 9. doi: 10.2174/1570159X19666210609163946.

Abstract

Melatonin, a vital hormone synthesized by the pineal gland, has been implicated in various physiological functions and in circadian rhythm regulation. Its role in the protection against the non-ionizing electromagnetic field (EMF), known to disrupt the body's oxidative/anti-oxidative balance, has been called into question due to inconsistent results observed across studies. This review provides the current state of knowledge on the interwoven relationship between melatonin, EMF, and oxidative stress. Based on synthesized evidence, we present a model that best describes the mechanisms underlying the protective effects of melatonin against RF/ELF-EMF induced oxidative stress. We show that the free radical scavenger activity of melatonin is enabled through reduction of the radical pair singlet-triplet conversion rate and the concentration of the triplet products.

Moreover, this review aims to highlight the potential therapeutic benefits of melatonin against the detrimental effects of EMF, in general, and electromagnetic hypersensitivity (EHS), in particular. <https://pubmed.ncbi.nlm.nih.gov/34635042/>

STATIC MAGNETIC FIELD MEASUREMENTS OF SMART PHONES AND WATCHES AND APPLICABILITY TO TRIGGERING MAGNET MODES IN IMPLANTABLE PACEMAKERS AND IMPLANTABLE CARDIOVERTER-DEFIBRILLATORS

Seth J Seidman, Joshua Guag, Brian Beard, Zane Arp. *Heart Rhythm*. 2021 Oct;18(10):1741-1744. doi: 10.1016/j.hrthm.2021.06.1203.

Abstract

Background: Implantable pacemakers and implantable cardioverter-defibrillators (ICDs) are designed to include a "magnet mode" feature that can be activated from magnets stronger than 10 G. This feature is designed to be used when a patient is undergoing a procedure where electromagnetic interference is possible, or anytime suspension of tachycardia detection and

therapy is needed. A publication in Heart Rhythm demonstrates an iPhone 12 triggering the magnet mode of a Medtronic ICD.

Objective: The purpose of this study is to determine the separation distance between consumer electronic devices that may create magnetic interference, including cell phones and smart watches, and implantable pacemakers and ICDs where magnet mode can be triggered.

Methods: The static magnetic fields of the iPhone 12 models and Apple Watch were measured at several planes in 1 cm resolution using an FW Bell 5180 Gauss Meter with STD18-0404 Transverse probe (unidirectional probe).

Results: All iPhone 12 and Apple Watch 6 models tested have static magnetic fields significantly greater than 10 G in close proximity (1-11 mm), which attenuates to below 10 G between 11 and 20 mm.

Conclusion: The findings of this study support the US Food and Drug Administration recommendation that patients keep any consumer electronic devices that may create magnetic interference, including cell phones and smart watches, at least 6 inches away from implanted medical devices, in particular pacemakers and cardiac defibrillators.

Open access paper: <https://www.frontiersin.org/articles/10.3389/fpubh.2021.721166/full>

INTERPRETATION OF TIMETRENDS (1996-2017) OF THE INCIDENCE OF SELECTED CANCERS IN ENGLAND IN RELATION TO MOBILE PHONE USE AS A POSSIBLE RISK FACTOR

Frank de Vocht. Interpretation of Timetrends (1996-2017) of the Incidence of Selected Cancers in England in Relation to Mobile Phone Use as a Possible Risk Factor. Bioelectromagnetics. 2021 Oct 11. doi: 10.1002/bem.22375.

Abstract

Radiofrequency (RF) radiation from mobile phones has been classified as possibly carcinogenic to humans (2b) by IARC. However, to date, the discussion on whether mobile phone use is a cancer risk factor has not been solved. In this context of continuing uncertainty, it is important to continue to monitor cancer incidence trends. Annual incidence rates and directly age-standardized rates of selected cancers by sex and 5-year age groups for 1996 to 2017 for England were obtained from the UK Office for National Statistics. Interpretation in light of mobile phone use as a contributing risk factor was conducted for cancers of the brain, parotid gland, thyroid, and colorectal cancer, which have all been hypothesized to be associated with RF exposure. Brain and parotid gland cancers were updated by an additional 10 years following a previous publication, and continue to provide little evidence of an association with mobile phone use. Although mobile phone use as a potential risk factor contributing to increased incidence of colorectal or thyroid cancer could not be excluded based on these ecological data, it is implausible that it is an important risk factor for either. In the absence of clarity from epidemiological studies, it remains important to continue to monitor trends. However, for the time being, and in agreement with data from other countries, there is little evidence of an association between mobile phone use and brain or parotid gland cancer, while the hypotheses of associations with thyroid or colorectal cancer are similarly weak. <https://pubmed.ncbi.nlm.nih.gov/34633685/>

My comments:

This paper provides a biased interpretation of data from a weak observational study that employs ecological, time series data. Given the limitations of cancer registry data, interpretation of cancer incidence trends is fraught with problems. Although the results of this observational study are not clear-cut, the overall results suggest that standardized cancer incidence rates in England for all four cancers under investigation increased over time:

“Although the DAS rates suggest an increase of 30% in the incidence rates of brain cancers in women and 34% in men from 1996 to 2017....”

“A steady increase ($p < 0.001$) from 0.4 to 0.6 incident cases per 100,000 women and 0.7 to 0.8 for men, respectively, over the 1999-2017 time period was observed for incidence of cancer of the parotid gland; mainly in DAS rates following the use of the revised ESP (Figure 2).”

“Cancer of the thyroid gland has increased steadily over time ($p < 0.001$), especially in women where the DAS rate has more than tripled from 1996 to 2017 from 2.7 incident cases to 8.5 per 100,000 women but also in men (from 1.2 to 3.6 incident cases per 100,000 men), with little differences between population rates and DAS rates (Figure 3).”

“The incidence rates of colorectal cancer in England have been stable for women (+3% from 1999 to 2017; $p < 0.001$) and slightly increasing for men (+15% from 1999 to 2017; $p < 0.01$) (Figure 4)....In younger age groups clear increase in incidence can be observed, especially in the age groups 25-34 year olds describing annual increases of 7% in both sexes ($p < 0.001$), with some indication of plateauing of this pattern in those of 29 years or younger.”

The above results could have been interpreted as supportive of the association with increases in mobile phone use over time in England due to the considerable lag between exposure to a risk factor and diagnosis of solid tumors. Furthermore, the case-control research suggests the increased cancer risk is a function of heavy mobile phone use (e.g., 17 minutes per day over a ten-year period for risk of glioma [Choi et al., 2020]) and perhaps also genetic susceptibility (for thyroid cancer risk [Luo et al., 2018, 2020]) so not everyone would be affected.

Analyzing time trend data may be a fruitless endeavor when there are multiple risk factors for the tumors under investigation because some carcinogens may be increasing over time while others are decreasing. It is impossible to make causal attributions with time series designs when there is a considerable lag between an exposure (e.g., mobile phone use) and an outcome (e.g., cancer). Similarly, it is problematic to rule out a risk factor with such a weak research design. The results clearly do not warrant the following assertions made by this paper:

“Although mobile phone use as a potential risk factor contributing to increased incidence of colorectal or thyroid cancer could not be excluded based on these ecological data, it is implausible that it is an important risk factor for either.”

“...there is little evidence of a causal relation of mobile phone use with brain or parotid gland cancer, while the hypotheses of associations with thyroid or colorectal cancer is similarly weak.”

Although the strongest epidemiologic evidence for increased cancer risk associated with mobile phone use is for glioma, a subtype of brain cancer (IARC Working Group, 2013), the paper did not cite Philips et al. (2018). Yet the Philips study examined brain cancer incidence data for England from 1995-2015 to calculate incidence rates (ASR) per 100,000 person-years, age-standardized to the European Standard Population (ESP–2013). Philips et al. found “a sustained and highly statistically significant ASR rise in glioblastoma multiforme (GBM) [a subtype of glioma] across all ages. The ASR for GBM more than doubled from 2.4 to 5.0, with annual case numbers rising from 983 to 2531. Overall, this rise is mostly hidden in the overall data by a reduced incidence of lower grade tumours.” Thus, it should come as no surprise that the current paper failed to find much of an increase in overall brain cancer incidence in England for almost the same time period.

PSYCHOLOGICAL MODELS OF DEVELOPMENT OF IDIOPATHIC ENVIRONMENTAL INTOLERANCES: EVIDENCE FROM LONGITUDINAL POPULATION-BASED DATA

Luc Watrin, Steven Nordin, Renáta Szemerszky, Oliver Wilhelm, Michael Witthöft, Ferenc Köteles. *Environ Res.* 2021 Sep 8;204(Pt A):111774. doi: 10.1016/j.envres.2021.111774.

Abstract

The origin of idiopathic environmental intolerances (IEIs) is an open question. According to the psychological approaches, various top-down factors play a dominant role in the development of IEIs. The general psychopathology model assumes a propensity towards mental ill-health (negative affectivity) increases the probability of developing IEIs. The attribution model emphasizes the importance of mistaken attribution of experienced somatic symptoms; thus, more symptoms should lead to more IEIs. Finally, the nocebo model highlights the role of expectations in the development of IEIs. In this case, worries about the harmful effects of environmental factors are assumed to evoke IEIs. We estimated cross-lagged panel models with latent variables based on longitudinal data obtained at two time points (six years apart) from a large near-representative community sample to test the hypothesized associations. Indicators of chemical intolerance, electromagnetic hypersensitivity, and sound sensitivity fit well under a common latent factor of IEIs. This factor, in turn, showed considerable temporal stability. However, whereas a positive association was found between IEIs and increased somatic symptoms and modern health worries six years later, the changes therein could not be predicted as hypothesized by the three psychological models. We discuss the implications of these results, as well as methodological aspects in the measurement and prediction of change in IEIs.

Highlights

- Compares and integrates theories of Idiopathic Environmental Intolerances (IEI).
- Tests competing theories in a large longitudinal study (N = 1837).
- Correlations, but no cross-lagged associations of IEIs with neighboring constructs.
- Latent variable models indicate a strong commonality of different IEIs.
- We advocate for more attention on the nomothetic span of IEIs.

Conclusion

In the current study, cross-lagged panel models were applied to investigate reciprocal association between IELs and general psychopathology, somatic symptoms, and modern health worries in longitudinal data of a large near-representative community sample. Contrary to theoretical predictions, higher initial levels in either of the three constructs were not associated with increases in IELs over a period of 6 years. The proneness to idiopathic environmental intolerances (IEIs), such as chemical intolerance, noise sensitivity, and electromagnetic hypersensitivity, appeared very stable. Somatic symptoms, indicators of negative affectivity, and modern health worries showed reliable associations with IELs. However, they did not predict IELs. Therefore, popular theories of the development of IELs could not be confirmed empirically. <https://pubmed.ncbi.nlm.nih.gov/34506786/>

RF MEASUREMENTS OF EMFS FOR HUMAN EXPOSURE ASSESSMENT DUE TO MODERN CELLULAR BASE STATIONS (5G)

D. Capriglione. In-Situ RF Measurements of EMFs for Human Exposure Assessment Due to Modern Cellular Base Stations. IEEE Instrumentation & Measurement Magazine, vol. 24, no. 8, pp. 31-36, November 2021, doi: 10.1109/MIM.2021.9580794.

Abstract

TC-37-Measurements and Networking (current Chair Prof. Domenico Capriglione, University of Cassino and Southern Lazio, Italy) was built some years ago as an idea of Prof. Leopoldo Angrisani (University of Napoli Federico II, Italy) and Prof. Claudio Narduzzi (University of Padova, Italy), having in mind that given the complexity of modern networks and related systems, a multidisciplinary approach has to be followed for correctly addressing the technological challenges and issues that are arising and will arise in the future in the field of telecommunication systems and networking. So, the TC-37 is actively trying to promote the international cooperation and integration of researchers belonging to the Instrumentation and Measurement Society with ones coming from other areas of telecommunication and information technologies.

Conclusions and Open Issues

The measurement of human exposure to RF EMFs is a topic of great interest today because of the growing diffusion and fast evolution of communication technologies. Due to the importance of the topic and the technical difficulties arising from the ever-increasing level of complexity of the communication technologies and experimental scenarios, worldwide researchers are involved in designing and fine-tuning measurement methods, standard procedures, and instruments for achieving reliable results of human exposure.

The practical examples in this paper highlight how the research in this field needs continuous updates, and it should involve long-term experimental campaigns for assessing the reliability of measurement techniques and procedures in several experimental conditions and scenarios. These statements are strongly supported by current trends that push to employ DSS (Dynamic Spectrum Sharing) and 5G cellular technologies which will offer new challenges for the measurement of human exposure to EMFs generated by these kinds of sources. In particular, the antenna beamforming and the complexity of 5G technology will require the design of new and effective measurement techniques and protocols able to warrant an adequate tradeoff between accuracy and time needed to completely characterize human

exposure in high-density urban scenarios, where several base stations simultaneously operate in the same area. These aspects shall be investigated for both narrowband and broadband approaches.

Furthermore, the estimation of the measurement uncertainty will be an important topic to be addressed, taking into account several quantities of influence among which the response of the antennas, probes, and instruments to the signals generated by modern communication systems, as well as the long-term variability of the power emitted by the related Base Stations in given points of analysis should be examined.

<https://ieeexplore.ieee.org/document/9580794>

SPATIAL VARIABILITY OF OUTDOOR EXPOSURE TO RADIOFREQUENCY RADIATION FROM MOBILE PHONE BASE STATIONS, IN KHARTOUM, SUDAN

Mohammed O A Mohammed, Ahmed A Elzaki, Babiker A Babiker, Omer I Eid. *Environ Sci Pollut Res Int.* 2021 Oct 8. doi: 10.1007/s11356-021-16555-x.

Abstract

The wide-spread exposure to constantly evolving wireless technologies believed to pose a serious health threat. Human beings are persistently exposed to RF radiation from mobile phones and their base stations. The current study aimed at classifying and characterizing the exposure to RF radiation from the mobile phone base stations. Spatial distribution measurements were carried out in Khartoum city during two time periods, first in 2012 (pilot survey) and again during Sept. 2019-Jan. 2020, to cover a total of 282 antennas operating with GSM900, GSM1800, and UMTS2100. The tested antennas belong to three mobile communication companies namely Sudani, Zain, and MTN companies, that randomly coded into company A, company B, and company C for security purposes. Measurements were performed using frequency-selective RF analyzer at fixed distances from the antennas/towers. Data were subjected to advanced repeated measures ANOVA, linear discriminant analysis (LDA), and spatial interpolation with ArcGIS. The averages of GSM900, GSM1800, and UMTS measurements were 0.01933 W/m², 0.0067 W/m², and 0.0046 W/m². The high levels of power densities for each single antenna were recorded at 90 m, 110 m, 130 m, and at 150 m distances, for the majority (70%) of the measured antennas and the peak/highest values reported mainly at 110 m distance. Conversely, the discriminant loadings as part of LDA, suggested that, much of variance among measurements is attributed to measurements at 150 m, 170 m, and 190 m distances, while visual illustration of group centroids implied that, the RF signals of the different companies were measured separately which support accuracy of frequency-selective measurements. The LDA has confirmed the ANOVA results that, the overall difference between the three companies was statistically significant for UMTS, and GSM900 measurements but not significant for GSM1800 measurements. Kriging interpolation using ArcGIS provided a strong evidence of great spatial distribution of exposure across the study area, with market places and typical urban residential quarters showing highest levels of RF. Few extreme values exceeding ICNIRP limits are reported but excluded from the calculations because of an issue of normality of data that is considered a prerequisite for parametric data analysis. Existence of extreme levels of RF indicates a need for further investigation and some antennas of Company B are mounted on towers belongs to Company C, implying multi exposure. Unexpected pattern of RF levels

continued to increase up to 190 m distance and possibly beyond 190 m is reported for UMTS measurements of Company C. <https://pubmed.ncbi.nlm.nih.gov/34622411/>

NEAR FIELD EXPOSURE CONDITIONS BY UHF-RFID SYSTEMS IN SMART HEALTHCARE ENVIRONMENTS

S. de Miguel-Bilbao et al. Near Field Exposure Conditions by UHF-RFID Systems in Smart Healthcare Environments. 2021 IEEE International Joint EMC/SI/PI and EMC Europe Symposium. 2021, pp. 13-18, doi: 10.1109/EMC/SI/PI/EMCEurope52599.2021.9559189.

Abstract

Short range wireless RFID technology has many applications in socialcare and healthcare environments, having to coexist with other sources of electromagnetic (EM) radiation and even with patients' implanted devices. This work provides an overview of exposure conditions in near EM field conditions and evaluates this exposure. The near EM field conditions by RFID reader is discussed based on the results of measurements inside an anechoic chamber under strict experimental conditions, and numerical modelling with simulation software. The obtained results were considered with respect to the human EM exposure evaluation principles and exposure limitations provided by the relevant international guidelines and regulations. In areas close to the RFID reader, the local exposure to EM radiation has the near field nature, i.e. the impedance of EM field significantly differs from the far field (free space). Evaluating human exposure requires measurements of the electric and magnetic field strength, or even the numerical modelling of Specific Energy Absorption Rate (SAR). It was found that a near field nature of EM radiation near an RFID reader ranges several times longer when the operator is present nearby, compared to the same emitting device considered alone in the empty space. This is of significance when evaluating EM exposure of humans (patients or health care personnel, especially users of medical implants) if, for any reason, they are less than 50 cm away from an RFID reader, especially when emission from it exceed 2 W.

<https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9559189&isnumber=9559134>

HUMAN-MADE ELECTROMAGNETIC FIELDS: ION FORCED-OSCILLATION AND VOLTAGE-GATED ION CHANNEL DYSFUNCTION, OXIDATIVE STRESS AND DNA DAMAGE (REVIEW)

Dimitris J Panagopoulos, Andreas Karabarbounis, Igor Yakymenko, George P Chrousos. Human-made electromagnetic fields: Ion forced-oscillation and voltage-gated ion channel dysfunction, oxidative stress and DNA damage (Review). Int J Oncol. 2021 Nov;59(5):92. doi: 10.3892/ijo.2021.5272.

Abstract

Exposure of animals/biological samples to human-made electromagnetic fields (EMFs), especially in the extremely low frequency (ELF) band, and the microwave/radio frequency (RF) band which is always combined with ELF, may lead to DNA damage. DNA damage is connected with cell death, infertility and other pathologies, including cancer. ELF exposure from high-voltage power lines and complex RF exposure from wireless communication antennas/devices are linked to increased cancer risk. Almost all human-made RF EMFs include ELF components in the form of modulation, pulsing and random variability. Thus, in addition to polarization and coherence, the existence of ELFs is a common feature of almost all human-made EMFs. The present study reviews the DNA damage and related effects induced by human-made EMFs. The ion forced-oscillation mechanism for irregular gating of voltage-

gated ion channels on cell membranes by polarized/coherent EMFs is extensively described. Dysfunction of ion channels disrupts intracellular ionic concentrations, which determine the cell's electrochemical balance and homeostasis. The present study shows how this can result in DNA damage through reactive oxygen species/free radical overproduction. Thus, a complete picture is provided of how human-made EMF exposure may indeed lead to DNA damage and related pathologies, including cancer.

Moreover, it is suggested that the non-thermal biological effects attributed to RF EMFs are actually due to their ELF components.

Open access paper: <https://www.spandidos-publications.com/ijo/59/5/92>

HIPPOCAMPAL OXIDATIVE STRESS INDUCED BY RADIOFREQUENCY ELECTROMAGNETIC RADIATION AND THE NEUROPROTECTIVE EFFECTS OF AEROBIC EXERCISE IN RATS: A RANDOMIZED CONTROL TRIAL

Mina Rasouli Mojez, Abbas Ali Gaeini, Siroos Choobineh, Mohsen Sheykhloovand. Hippocampal Oxidative Stress Induced by Radiofrequency Electromagnetic Radiation and the Neuroprotective Effects of Aerobic Exercise in Rats: A Randomized Control Trial. J Phys Act Health. 2021 Oct 25;1-7. doi: 10.1123/jpah.2021-0213.

Abstract

Background: The present study determined whether 4 weeks of moderate aerobic exercise improves antioxidant capacity on the brain of rats against oxidative stress caused by radiofrequency electromagnetic radiation emitted from cell phones.

Methods: Responses of malondialdehyde, catalase, glutathione peroxidase, and superoxide dismutase, as well as the number of hippocampal dead cells, were examined. Male Wistar rats (10-12 wk old) were randomly assigned to 1 of 4 groups (N = 8): (1) moderate aerobic exercise (EXE) (2 × 15-30 min at 1215 m/min speed with 5 min of active recovery between sets), (2) exposure to 900/1800 MHz radiofrequency electromagnetic waves 3 hours per day (RAD), (3) EXE + RAD, and (4) exposure to an experimental phone without battery.

Results: Following the exposure, the number of the hippocampal dead cells was significantly higher in group RAD compared with groups EXE, EXE + RAD, and control group. Malondialdehyde concentration in group RAD was significantly higher than that of groups EXE, EXE + RAD, and control group. Also, the activity of catalase, glutathione peroxidase, and superoxide dismutase in groups EXE, EXE + RAD, and control group was significantly higher compared with those of the exposure group.

Conclusion: This study demonstrated that moderate aerobic exercise enhances hippocampal antioxidant capacity against oxidative challenge in the form of radiofrequency electromagnetic waves.

<https://pubmed.ncbi.nlm.nih.gov/34697252/>

VM-RELATED EXTRACELLULAR POTENTIALS OBSERVED IN RED BLOOD CELLS

Hughes MP, Kruchek EJ, Beale AD, Kitcatt SJ, Qureshi S, Trott ZP, Charbonnel O, Agbaje PA, Henslee EA, Dorey RA, Lewis R, Labeed FH. Vm-related extracellular potentials observed in red blood cells. *Sci Rep.* 2021 Sep 30;11(1):19446. doi: 10.1038/s41598-021-98102-9.

Abstract

Even in nonexcitable cells, the membrane potential Vm is fundamental to cell function, with roles from ion channel regulation, development, to cancer metastasis. Vm arises from transmembrane ion concentration gradients; standard models assume homogeneous extracellular and intracellular ion concentrations, and that Vm only exists across the cell membrane and has no significance beyond it. Using red blood cells, we show that this is incorrect, or at least incomplete; Vm is detectable beyond the cell surface, and modulating Vm produces quantifiable and consistent changes in extracellular potential. Evidence strongly suggests this is due to capacitive coupling between Vm and the electrical double layer, rather than molecular transporters. We show that modulating Vm changes the extracellular ion composition, mimicking the behaviour of voltage-gated ion channels in non-excitable channels. We also observed Vm-synchronised circadian rhythms in extracellular potential, with significant implications for cell-cell interactions and cardiovascular disease.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8484267/>

A SURVEY ON ELECTROMAGNETIC HYPERSENSITIVITY: THE EXAMPLE FROM POLAND

Grzegorz Tatoń, Artur Kacprzyk, Tomasz Rok, Monika Pytlarz, Rafał Pawlak, Eugeniusz Rokita. A survey on electromagnetic hypersensitivity: the example from Poland. *Electromagn Biol Med.* 2021 Oct 22;1-8. doi: 10.1080/15368378.2021.1995873.

Abstract

Idiopathic environmental intolerance attributed to electromagnetic field (IEI-EMF) called electromagnetic hypersensitivity or electrosensitivity appeared in Polish society awareness due to a considerable change made at the end of 2019 in Polish telecommunication laws. The aims of the project were to assess the prevalence of IEI-EMF in Poland and to define a reliable methodology to study this phenomenon. The first step was the internet survey performed at the end of 2018. The IEI-EMF prevalence estimated at the level of 39.7% suggested considerable bias affecting the results. The faults of the first approach were analysed and then a second study stage was performed as a telephone survey at the end of 2020. The latter survey allowed estimating the prevalence of IEI-EMF as less than 1.8%. These discrepancies in the results of both surveys were connected to the medium used in the first survey (Internet) indirectly causing that the group pooled was not representative. The second pitfall was the definition of the criteria used for an electrosensitive person classification. This is why the IEI-EMF prevalence was investigated in the second stage with the use of numerous criteria. The application of different criteria allowed for essential conclusions concerning the appropriate methodology for such kinds of studies. Corrections of the methodology before the second survey allowed reliable results consistent with the results obtained in similar studies performed in other countries. Our findings also show that the IEI-EMF frequency reports presented in the literature have to be treated carefully and with some dose of scepticism.

<https://pubmed.ncbi.nlm.nih.gov/34686066/> The worldwide prevalence of IEI-EMF is estimated at 1–13%. The results vary according to the data source and methodology of the studies used (Huang et al. 2018; Szemerszky et al. 2019).

Conclusions

The first study overestimated the incidence of IEI-EMF in Polish society at 39.7% due the limitations accompanying the study. The more realistic estimation based on the second research stage is 1.8%.

The proposed criterion of IEI-EMF qualification assumes that a hypersensitive person is one whose symptoms differ from the rest of society and are so severe that she or he seeks medical help.

Our results support the thesis that the criterion used in such studies for assigning subjects to the IEI-EMF group should be accurately defined and harmonized.

Even slight differences in the survey questions and methodologies can result in considerably different IEI-EMF prevalence estimations.

THE HAVANA SYNDROME AND MICROWAVE WEAPONS [HEALTH MATTERS]

James C. Lin. The Havana Syndrome and Microwave Weapons [Health Matters]. IEEE Microwave Magazine. 22(11):13-14. Nov. 2021. doi: 10.1109/MMM.2021.3102201.

Abstract

Every few months, if not weeks, another mysterious attack on U.S. diplomatic and intelligence personnel is reported. Some of the attacks occurred years ago, while others were recounted as recently as July 2021 [1] – [3]. Over the past four or five years, nearly 200 U.S. personnel have reported similar attacks while working in places like Havana, Guangzhou, London, Moscow, Vienna, and Washington, D.C. The acute symptoms include headache and nausea immediately following the sounds of loud buzzing or bursts. The illness and symptoms have been called the “Havana Syndrome” after the place where cases were first reported. It refers to the range of symptoms first experienced by U.S. State Department personnel overseas.

Excerpt

Assuming reported accounts are reliable, the microwave auditory effect provides a scientific explanation for the Havana Syndrome [4], [5]. Pulsed microwaves can create an acoustic wave inside the head [6], [7], [14]. It is plausible that the loud buzzing, burst of sound, or pressure waves could have been covertly delivered using a beam of high power pulsed microwave radiation rather than blasting the subjects with conventional sonic sources. Microwave hearing doesn’t go through the ear; it goes directly from the brain tissue to the cochlea.

Absorption of short pulses of microwave energy by brain tissues creates a rapid expansion of brain matter and launches an acoustic wave of pressure (sound wave) that travels inside the head to the inner ear cochlea [7], [14]. The short high-power microwave pulses do not generate noticeable amounts of heat in the brain tissues.

<https://ieeexplore.ieee.org/document/9557212>

ELECTROMAGNETIC EXPOSURE DOSIMETRY STUDY ON TWO FREE RATS AT 1.8 GHZ VIA NUMERICAL SIMULATION

Wang Xianghui, Xia Chengjie, Lu Lu, Qi Hongxin, Zhang Jie. Electromagnetic Exposure Dosimetry Study on Two Free Rats at 1.8 GHz via Numerical Simulation. *Frontiers in Public Health*. 9:1404. 2021. doi: 10.3389/fpubh.2021.721166

Abstract

Normally, the impact of electromagnetic exposure on human health is evaluated by animal study. The biological effect caused by electromagnetic exposure on such experimental animals as rats has been proven to be dose- dependent. However, though the dose of radio frequency (RF) electromagnetic exposure described by the specific absorbing rate (SAR) on fixed rats has been relatively well-studied utilizing the numerical simulations, the dosimetry study of exposure on free rat is insufficient, especially in the cases of two or more free rats. Therefore, the present work focuses on the variation of SAR caused by the existence of neighboring free rat in the same cage. Here, infrared thermography was used to record the activity of the two free rats who lived in the same cage that mounted at the far-field region in the microwave darkroom for a duration of 48 h. Then, using image processing techniques, the relative positions and orientations of the two rats are identified, which are defined by three parameters, such as the relative distance (d), relative direction angle (α), and relative orientation angle (β). Using the simulation software XFDTD 7.3, the influence of d , α , and β on the whole-body average SAR (WB-avgSAR) of the rats exposed to 1.8 GHz electromagnetic wave was calculated and analyzed. Then, the average variation of WB-avgSAR of the two rats compared with that of a single rat within 48 h was calculated. The numerical simulation results showed that the relative posture position described by (d , α , and β) of the two rats affects their WB-avgSAR and leads to fluctuations at different positions. However, the variation rate of the 48-h-average WB-avgSAR was only 10.3%, which implied that the over-time average SAR of two or more rats can be roughly described by the WB-avgSAR of a single free rat, except when a real-time precise control of exposure dose is necessary.

<https://www.frontiersin.org/article/10.3389/fpubh.2021.721166>

DESIGN AND DOSIMETRIC CHARACTERIZATION OF A BROADBAND EXPOSURE FACILITY FOR IN VITRO EXPERIMENTS IN THE FREQUENCY RANGE 18-40.5 GHZ

Gernot Schmid, Rene Hirtl, Isabel Gronau, Vivian Meyer, Karen Drees, Alexander Lerchl. Design and Dosimetric Characterization of a Broadband Exposure Facility for In Vitro Experiments in the Frequency Range 18-40.5 GHz. *Bioelectromagnetics*. 2021 Oct 7. doi: 10.1002/bem.22376.

Abstract

A novel exposure facility for exposing cell monolayers to centimeter and millimeter waves (18-40.5 GHz) used by future 5G mobile communication technology and similar applications has been developed. A detailed dosimetric characterization of the apparatus for frequencies of 27 and 40.5 GHz and 60 mm petri dishes, used in a presently ongoing study on human dermal fibroblasts and keratinocytes, was carried out. The exposure facility enables a well-defined, randomized, and blinded application of sham exposure and exposure with selectable values of incident power flux density, and additionally provides the possibility of continuous monitoring of the sample temperature during exposure while it does not require significant deviations from routine in vitro handling procedures, i.e. petri dishes are not required to be

placed inside waveguides or TEM cells. Mean specific absorption rate (SAR) values inside the cell monolayer of 115 W/kg (27 GHz) and 160 W/kg (40.5 GHz) per watt antenna input power and corresponding transmitted power density (S_t) values at the bottom of the cell monolayer of 65 W/m² (27 GHz) and 70 W/m² (40.5 GHz) per watt antenna input power can be achieved, respectively. For reasonable amounts of harvested cells (80% of petri dish bottom area), the variation (max/min) of SAR and S_t over the cell monolayer remains below 3.7 dB (27 GHz) and 3.0 dB (40.5 GHz), respectively. <https://pubmed.ncbi.nlm.nih.gov/34622469/>

THE ECOLOGY OF ELECTRICITY AND ELECTRORECEPTION

Sam J England, Daniel Robert. The ecology of electricity and electroreception. *Biol Rev Camb Philos Soc.* 2021 Oct 12. doi: 10.1111/brv.12804.

Abstract

Electricity, the interaction between electrically charged objects, is widely known to be fundamental to the functioning of living systems. However, this appreciation has largely been restricted to the scale of atoms, molecules, and cells. By contrast, the role of electricity at the ecological scale has historically been largely neglected, characterised by punctuated islands of research infrequently connected to one another. Recently, however, an understanding of the ubiquity of electrical forces within the natural environment has begun to grow, along with a realisation of the multitude of ecological interactions that these forces may influence. Herein, we provide the first comprehensive collation and synthesis of research in this emerging field of electric ecology. This includes assessments of the role electricity plays in the natural ecology of predator-prey interactions, pollination, and animal dispersal, among many others, as well as the impact of anthropogenic activity on these systems. A detailed introduction to the ecology and physiology of electroreception - the biological detection of ecologically relevant electric fields - is also provided. Further to this, we suggest avenues for future research that show particular promise, most notably those investigating the recently discovered sense of aerial electroreception. <https://pubmed.ncbi.nlm.nih.gov/34643022/>

EFFECTS OF AN ELECTRIC FIELD ON SLEEP QUALITY AND LIFE SPAN MEDIATED BY ULTRAVIOLET (UV)-A/BLEU LIGHT PHOTORECEPTOR CRYPTOCHROME IN DROSOPHILA

Haruhisa Kawasaki, Hideyuki Okano, Takaki Nedachi, Yuzo Nakagawa-Yagi, Akikuni Hara, Norio Ishida. Effects of an electric field on sleep quality and life span mediated by ultraviolet (UV)-A/blue light photoreceptor CRYPTOCHROME in *Drosophila*. *Sci Rep.* 2021 Oct 15;11(1):20543. doi: 10.1038/s41598-021-99753-4.

Abstract

Although electric fields (EF) exert beneficial effects on animal wound healing, differentiation, cancers and rheumatoid arthritis, the molecular mechanisms of these effects have remained unclear about a half century. Therefore, we aimed to elucidate the molecular mechanisms underlying EF effects in *Drosophila melanogaster* as a genetic animal model. Here we show that the sleep quality of wild type (WT) flies was improved by exposure to a 50-Hz (35 kV/m) constant electric field during the day time, but not during the night time. The effect was undetectable in cryptochrome mutant (*cryb*) flies. Exposure to a 50-Hz electric field under low nutrient conditions elongated the lifespan of male and female WT flies by ~ 18%, but not of several *cry* mutants and *cry* RNAi strains. Metabolome analysis indicated that the adenosine triphosphate (ATP) content was higher in intact WT than *cry* gene mutant strains exposed to

an electric field. A putative magnetoreceptor protein and UV-A/blue light photoreceptor, CRYPTOCHROME (CRY) is involved in electric field (EF) receptors in animals. The present findings constitute hitherto unknown genetic evidence of a CRY-based system that is electric field sensitive in animals. Open access paper: <https://www.nature.com/articles/s41598-021-99753-4>

TRACKING DEVICES FOR PETS: HEALTH RISK ASSESSMENT FOR EXPOSURE TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS

Judith Klune, Christine Arhant, Ines Windschnurer, Veronika Heizmann, Günther Schaubberger. Tracking Devices for Pets: Health Risk Assessment for Exposure to Radiofrequency Electromagnetic Fields. *Animals (Basel)*. 2021 Sep 17;11(9):2721. doi: 10.3390/ani11092721.

Abstract

Every year, approximately 3% of cats and dogs are lost. In addition to passive methods for identifying pets, radiofrequency tracking devices (TDs) are available. These TDs can track a pet's geographic position, which is transmitted by radio frequencies. The health risk to the animals from continuous exposure to radiofrequency electromagnetic fields (RF-EMFs) was reviewed. Fourteen out of twenty-one commercially available TDs use 2G, 3G, or 4G mobile networks, and the others work with public frequencies, WLAN, Bluetooth, etc. The exposure of pets to RF-EMFs was assessed, including ambient exposure (radios, TVs, and base stations of mobile networks), exposure from indoor devices (DECT, WLAN, Bluetooth, etc.), and the exposure from TDs. The exposure levels of the three areas were found to be distinctly below the International Commission on Non-Ionising Radiation Protection (ICNIRP) reference levels, which assure far-reaching protection from adverse health effects. The highest uncertainty regarding the exposure of pets was related to that caused by indoor RF-emitting devices using WLAN and DECT. This exposure can be limited considerably through a reduction in the exposure time and an increase in the distance between the animal and the RF-emitting device. Even though the total RF-EMF exposure level experienced by pets was found to be below the reference limits, recommendations were derived to reduce potential risks from exposure to TDs and indoor devices.

Open access paper: <https://www.mdpi.com/2076-2615/11/9/2721>

EFFECT OF RADIATION EMITTED BY WIRELESS DEVICES ON MALE REPRODUCTIVE HORMONES: A SYSTEMATIC REVIEW

Maluin Sofwatul Mokhtarah, Osman Khairul, Jaffar Farah Hanan Fathihah, Ibrahim Siti Fatimah. Effect of Radiation Emitted by Wireless Devices on Male Reproductive Hormones: A Systematic Review. *Frontiers in Physiology*. 12:1568. 2021. doi:10.3389/fphys.2021.732420.

Abstract

Exposure to radiofrequency electromagnetic radiation (RF-EMR) from various wireless devices has increased dramatically with the advancement of technology. One of the most vulnerable organs to the RF-EMR is the testes. This is due to the fact that testicular tissues are more susceptible to oxidative stress due to a high rate of cell division and mitochondrial oxygen consumption. As a result of extensive cell proliferation, replication errors occur, resulting in DNA fragmentation in the sperm. While high oxygen consumption increases the level of oxidative phosphorylation by-products (free radicals) in the mitochondria. Furthermore, due to its inability to effectively dissipate excess heat, testes are also

susceptible to thermal effects from RF-EMR exposure. As a result, people are concerned about its impact on male reproductive function. The aim of this article was to conduct a review of literature on the effects of RF-EMR emitted by wireless devices on male reproductive hormones in experimental animals and humans. According to the findings of the studies, RF-EMR emitted by mobile phones and Wi-Fi devices can cause testosterone reduction. However, the effect on gonadotrophic hormones (follicle-stimulating hormone and luteinizing hormone) is inconclusive. These findings were influenced by several factors, which can influence energy absorption and the biological effect of RF-EMR. The effect of RF-EMR in the majority of animal and human studies appeared to be related to the duration of mobile phone use. Thus, limiting the use of wireless devices is recommended. Conclusion Existing animal and human data on the effect of RF-EMR emitted from wireless devices on male reproductive hormones are inconsistent and difficult to evaluate due to the heterogeneity of study design. However, most studies are consistent with the assertion that long-term exposure to RF-EMR from mobile phones and Wi-Fi devices can disrupt male reproductive hormones, particularly testosterone. Thus, avoiding long-term and excessive use of mobile phone is advisable to reduce the detrimental effect of RF-EMR.

Open access paper: <https://www.frontiersin.org/article/10.3389/fphys.2021.732420>

CHALLENGES ON THE EFFECT OF CELL PHONE RADIATION ON MAMMALIAN EMBRYOS AND FETUSES: A REVIEW OF THE LITERATURE

Maryam Mahaldashtian, Mohammad Ali Khalili, Fatemeh Anbari, Mohammad Seify, Manuel Belli. Challenges on the effect of cell phone radiation on mammalian embryos and fetuses: a review of the literature. *Zygote*. 2021 Sep 29;1-7. doi: 10.1017/S0967199421000691.

Abstract

Cell phones operate with a wide range of frequency bands and emit radiofrequency-electromagnetic radiation (RF-EMR). Concern on the possible health hazards of RF-EMR has been growing in many countries because these RF-EMR pulses may be absorbed into the body cells, directly affecting them. There are some in vitro and in vivo animal studies related to the consequences of RF-EMR exposure from cell phones on embryo development and offspring. In addition, some studies have revealed that RF-EMR from cellular phone may lead to decrease in the rates of fertilization and embryo development, as well as the risk of the developmental anomalies, other studies have reported that it does not interfere with in vitro fertilization or intracytoplasmic sperm injection success rates, or the chromosomal aberration rate. Of course, it is unethical to study the effect of waves generated from cell phones on the forming human embryos. Conversely, other mammals have many similarities to humans in terms of anatomy, physiology and genetics. Therefore, in this review we focused on the existing literature evaluating the potential effects of RF-EMR on mammalian embryonic and fetal development. <https://pubmed.ncbi.nlm.nih.gov/34583799/>

Excerpts

Mobile phones are commonly placed on the lap or in pockets (Swerdlow et al., 2011), therefore exposing the genital area to RF-EMF. Germ cells are more susceptible to RF-EMF, as they are rapidly dividing through meiosis and mitosis. EMF induces modifications in cellular levels such as activation of voltage-gated calcium channels, formation of free radicals, protein misfolding and DNA damage (Altun et al., 2018). In the general population, pregnant women

are at specific risk of exposure to environmental RF-EMF because of their higher oxygen consumption and amniotic fluid-induced ROS production (Çiğ and Nazıroğlu, 2015) (Fig. 1). Individuals should reduce their rate of exposure to RF-EMF-emitting devices including mobile phones (Okechukwu, 2020) as electromagnetic waves enhanced the amount of oxygen free radicals in the body that led to disturbance of spermatogenesis process and variations in spermatozoa membrane, which resulted in changes in capacitation, acrosome reaction and therefore disorder in the process of spermatozoa binding to the oocyte and failure of normal fertilization both in vivo and in vitro (Fatehi et al., 2018). In addition, exposure to RF may lead to a decline in ovarian follicle reservoirs at the start of the prepubertal time (Türedi et al., 2016).

To reduce the contact with RF-EMR radiated by cell phones, people should avoid keeping their phones inside their pockets, also use of hand-free materials and gadgets and mobile covers limit direct expose of cell phones to the body (Okechukwu, 2020). Also, people do not appreciate that a cell phone is a small base station and usually underrate the importance of distance between the cell phone and the body (Cousin and Siegrist, 2010). In analyzing the relationship between the distance of cell phone antenna from the body and SAR, Hossain and colleagues reported that increasing the distance of cell phone from the body led to diminishing of both electric and magnetic field strength to the body, therefore reducing SAR values (Hossain et al., 2015). Furthermore, it is known that the EMF decreases with distance and magnetic induction...

Conclusion

At this time, it is difficult from the available animal studies to document confidently the role of RF-EMR exposure on human embryo development, both in vivo and in vitro. Further investigations with complementary techniques will be necessary to understand the mechanism of action of RF-EMR emitted by cell phones and the consequences on mammals, particularly human beings.

AN EXPLORATION OF THE EFFECTS OF RADIOFREQUENCY RADIATION EMITTED BY MOBILE PHONES AND EXTREMELY LOW FREQUENCY RADIATION ON THYROID HORMONES AND THYROID GLAND HISTOPATHOLOGY

Alkayyali T, Ochuba O, Srivastava K, Sandhu JK, Joseph C, Ruo SW, Jain A, Waqar A, Poudel S. An Exploration of the Effects of Radiofrequency Radiation Emitted by Mobile Phones and Extremely Low Frequency Radiation on Thyroid Hormones and Thyroid Gland Histopathology. *Cureus*. 2021 Aug 20;13(8):e17329. doi: 10.7759/cureus.17329.

Abstract

The use of mobile phones has widely increased over the last two decades. Mobile phones produce a radiofrequency electromagnetic field (RF-EMF), a form of non-ionizing radiation. In contrast to the ionizing radiation proven to cause DNA damage, the harmful effects of non-ionizing radiation on the human body have not been discovered yet. The thyroid gland is among the most susceptible organs to mobile phone radiation due to its location in the anterior neck. Our purpose in this literature review is to explore the effects of the electromagnetic field (EMF), especially radiofrequency emitted from mobile phones, on thyroid hormones and thyroid gland histopathology. We searched PubMed and Google Scholar databases for relevant studies published after the year 2000, using the following

keywords: 'cell phones', 'mobile phones', 'telephones', 'electromagnetic fields', 'radiofrequency radiation', 'microwaves', 'thyroid gland', 'thyroid hormones', and 'thyroid cancer'. Our review revealed that mobile phone radiofrequency radiation (RFR) might be associated with thyroid gland insufficiency and alterations in serum thyroid hormone levels, with a possible disruption in the hypothalamic-pituitary-thyroid axis. The review also showed histopathological changes in the thyroid gland follicles after exposure of rats to non-ionizing radiation. The results were directly related to the amount and duration of exposure to EMF radiation. Further human studies exploring thyroid gland hormones, microscopic morphology, and thyroid cancer are highly recommended for future researches.

Conclusions

This article aimed to explore the effects of RF-EMF and ELF-EMF on the thyroid gland hormones and histopathology. Studies collected in this review showed that GSM mobile phone RFR could be associated with alterations in T3, T4, and TSH serum hormone levels. EMF emitted from mobile phones could disrupt the function of the HPT axis and lead to thyroid insufficiency. In addition, EMF could lead to hyperstimulation of thyroid gland follicles, causing oxidative stress and apoptosis of follicular cells. Most studies revealed a proportional correlation between thyroid gland dysfunction and the exposure duration, intensity, and SAR value of radiation. Moreover, non-ionizing radiation was seen to be significantly associated with histopathological changes in the thyroid gland follicles. The exposure duration and intensity also determined the degree of morphological damage occurring in the thyroid gland tissue. Non-ionizing EMF radiation might be responsible for the recent increase in the incidence of thyroid insufficiency and cancer in the general population. However, not enough data was found related to thyroid cancer risk with non-ionizing radiation exposure. Keeping in mind the ethical considerations, we recommend future observational studies be conducted on human beings to further explore the association of non-ionizing radiation emitted from mobile phones on the thyroid gland's hormones, histopathology, and cancers over the long term. Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8451508/>

MOBILE PHONE USE AND TIME TREND OF BRAIN CANCER INCIDENCE RATE IN KOREA

Kyung-Hwa Choi, Johyun Ha, Sanghyuk Bae, Ae-Kyoung Lee, Hyung-Do Choi, Young Hwan Ahn, Mina Ha, Hyunjoo Joo, Ho-Jang Kwon, Kyu-Won Jung, Mobile Phone Use and Time Trend of Brain Cancer Incidence Rate in Korea. 20 September 2021. /doi: 10.1002/bem.22373.

Abstract

This study evaluated the time trends in mobile phone subscriber number by mobile network generation (G) and brain cancer incidence by type in Korea. We obtained data from the Information Technology Statistics of Korea (1984–2017) and Korea Central Cancer Registry (1999–2017). The average annual percent change was estimated using Joinpoint regression analysis. We evaluated 29,721 brain cancer cases with an age-standardized incidence rate (ASR) of 2.89/100,000 persons. The glioma and glioblastoma annual ASR significantly increased in 2.6% and 3.9% of males and 3.0% and 3.8% of females, respectively. The ASR for frontal lobe involvement was the highest. The ASR of gliomas of unspecified grade annually increased by 7.8%; those for unspecified topology and histology decreased. The incidence of glioma, glioblastoma, frontal, temporal, and high-grade glioma increased among those aged ≥60 years. No association was observed between the mobile phone subscriber number and

brain cancer incidence in Korea. Furthermore, long-term research is warranted because of the latency period of brain cancer.

Grant sponsor: ICT R&D program of MSIT/IITP [2019-0-00102, A study on public health and safety in a complex EMF environment].

My summary of results:

- Glioma incidence increased over time (1999-2017) for adults aged 60 years and over.
- Glioblastoma incidence increased over time for adults aged 60 years and over.
- Unspecified brain tumor incidence decreased over time for adults aged 60 years and over.
- High-grade glioma incidence increased over time for male and female adults aged 60 years and over.
- The trend over time in standardized incidence of glioma exceeded the predicted relative risk of 1.5 based on mobile phone subscriber number.
- The trend over time in standardized incidence of glioblastoma exceeded the predicted relative risk of 1.5 based on mobile phone subscriber number.
- The trend over time in standardized incidence of brain cancer in the frontal lobe exceeded a predicted relative risk of 1.5 based on mobile phone subscriber number.

Excerpts

... According to the type of histology, the incidence of glioma was higher than that of other types, and it was significantly increased in both sexes (AAPC [95% CI], male: 2.6% [2.1, 3.2], female: 3.0% [2.1, 3.8]), whereas those of total brain cancers were not (AAPC [95% CI], male: 0.2% [-0.3, 0.6], female: 0.1% [-0.3, 0.5]). In particular, the incidence of glioblastoma increased more than that of other gliomas. The incidence of cancers with unspecified histology statistically decreased in both sexes (AAPC [95% CI], male: 7.9% [-9.6, -6.1], female: 7.8% [-8.9, -6.7]). According to the type of topology, the ASR of frontal lobe involvement (male: 0.53, female: 0.45) was higher than that of the temporal and parietal lobes. The ASR of cancers with unspecified topology decreased by 4.6% per year in males and 3.8% per year in females, whereas that of cancers of the other lobes increased. According to the type of grade, the incidence and AAPC of high-grade gliomas were higher than those of low-grade gliomas. The ASR of gliomas of unspecified grade increased by 7.8% per year, whereas those of unspecified topology (3.8% per year) and unspecified histology (7.8% per year) decreased.

... The incidence of glioma, glioblastoma, frontal, temporal, and high-grade glioma increased among those aged ≥ 60 years, and increased significantly among those aged ≥ 70 years (AAPC [95% CI]: glioma (male: 6.9 [5.6, 8.2], female: 8.5 [6.5, 10.6]); glioblastoma (male: 8.1 [6.1, 10.2], female: 9.9 [7.3, 12.5]); frontal (male: 6.2 [4.5, 8.0], female: 6.6 [4.0, 9.4]); temporal (male: 5.8 [2.9, 8.7], female: 7.1 [4.6, 9.7]); high-grade glioma (male: 7.5 [5.8, 9.3], female: 9.2 [7.0, 11.4]). There were no differences in the trend for unspecified brain cancer by histology, topology, or grade of glioma according to age group...

This study has several limitations. First, it is difficult to prove whether exposure to RF-EMF from mobile phones increases the incidence of brain cancers because this is a descriptive study. Further studies are required to evaluate the association between brain cancer

incidence and mobile phone subscribers by age group, because this study shows different results for each age group, such as an increase in the incidence of brain cancer among older adults. Therefore, we could not determine whether the increase in brain cancer incidence, especially glioma and glioblastoma, was associated with an increase of mobile phone subscribers in older age groups.

Second, it was not possible to calculate the predicted incidence according to the mobile phone subscription by sex and age. For this, it is better to acquire detailed information such as sex, age, and residential address of mobile phone subscribers in future studies and analyze them further. Studies have revealed a latency period of more than 30 years for cancer [Sadetzki et al., 2005]. However, data on cancer incidence are only available for approximately 20 years in Korea. Hence, further analysis is required to verify our results.

This study found no evidence of an increase in brain cancer incidence associated with the increasing number of mobile phone subscribers in Korea. Further research is warranted using detailed information on mobile phone subscriptions, with research carried out for longer durations considering the latency period for brain cancer. <https://onlinelibrary.wiley.com/doi/10.1002/bem.22373>

5G NEW RADIO REQUIRES THE BEST POSSIBLE RISK ASSESSMENT STUDIES: PERSPECTIVE AND RECOMMENDED GUIDELINES

Mats-Olof Mattsson, Myrtill Simkó, Kenneth R. Foster. 5G New Radio Requires the Best Possible Risk Assessment Studies: Perspective and Recommended Guidelines. *Frontiers in Communications and Networks*. Vol 2:49. 2021. doi: 10.3389/frcmn.2021.724772.

Abstract

The development and establishment of mobile communication technologies has necessitated assessments of possible risks to human health from exposures to radio-frequency electromagnetic fields (RF EMF). A number of expert committees have concluded that there is no evidence for such risks as long as exposures are at or below levels that do not allow tissue heating. These assessments have been based primarily on studies investigating frequencies up to 6 GHz including frequencies similar to those used by two of three major bands of fifth generation (more accurately 5G New Radio or 5G NR) of mobile communication. Bioeffects studies in so-called high-band at 25–39 GHz are particularly sparse. Future assessments relevant for these frequencies will need to rely on still unperformed studies. Due to few available studies at 5G NR “high band” frequencies, and questions raised by some existing studies, a recent review recommended a wide range of RF biostudies be done at 5G NR “high band” frequencies. It is of importance that such studies be done using the best possible science. Here we suggest factors to consider when performing future studies in this area. The present focus is on laboratory studies to clarify biological effects of radiofrequency (RF) energy at 5G “high band” frequencies and, more generally at millimeter wave (mm-wave) frequencies (30–300 GHz) which will be increasingly used by communications technologies in the future. Similar comments would apply to epidemiology and exposure assessment studies, but those are not the focus of the present Perspective. Open access paper: <https://www.frontiersin.org/article/10.3389/frcmn.2021.724772>

MAPPING OF STATIC MAGNETIC FIELDS NEAR THE SURFACE OF MOBILE PHONES

L. Zastko, L. Makinistian, A. Tvarožná, F. L. Ferreyra & I. Belyaev. Mapping of static magnetic fields near the surface of mobile phones. *Scientific Reports*. 11:19002. 2021. doi: 10.1038/s41598-021-98083-9.

Abstract

Whether the use of mobile phones (MP) represents a health hazard is still under debate. As part of the attempts to resolve this uncertainty, there has been an extensive characterization of the electromagnetic fields MP emit and receive. While the radiofrequencies (RF) have been studied exhaustively, the static magnetic fields (SMF) have received much less attention, regardless of the fact there is a wealth of evidence demonstrating their biological effects. We performed 2D maps of the SMF at several distances from the screen of 5 MP (models between 2013 and 2018) using a tri-axis magnetometer. We built a mathematical model to fit our measurements, extrapolated them down to the phones' screen, and calculated the SMF on the skin of a 3D head model, showing that exposure is in the μT to mT range. Our literature survey prompts the need of further research not only on the biological effects of SMF and their gradients, but also on their combination with extremely low frequency (ELF) and RF fields. The study of combined fields (SMF, ELF, and RF) as similar as possible to the ones that occur in reality should provide a more sensible assessment of potential risks.

Excerpts

The maximum SMF estimated in this work (6.6 mT) and others' (20 mT) 14 are far below the 400 mT exposure limit for the general public recommended by the International Commission on Non-Ionizing Radiation Protection (ICNIRP)26. This might lead to the temptation of completely disregarding SMF as a variable of interest when assessing MP fields. In fact, a recent fact sheet from the Swiss Federal Office of Public Health 13 completely omitted any reference to SMF. Nevertheless, the sole capacity of these fields of inducing potentially harmful biological effects such as promotion of the production of reactive oxygen species (ROS) 21 should be enough to consider them seriously. As for the possible underlying mechanisms of interaction, the one based on magnetite nanoparticles 27 is of particular interest because these nanoparticles have been found in the human brain 28,29. Besides the brain, the ear is of an obvious interest for its proximity to the upper HS of MP. Interestingly, magnetic material was reported on the lagenal otoliths of fish and birds 30 (although this structure is not conserved in mammals). To the best of our knowledge, there have not been studies reporting the search of magnetic materials as constituents of the human inner ear. In this regard, it is to be mentioned that some epidemiological studies found an association between MP use and tinnitus (the perception of a sound in the absence of an external source), while a meta-analysis found no association 31...

Our measurements and calculations, along with the preceding discussion of the literature, prompt the need of further studies of the biological effects of the SMF and RF EMF generated by MP. We suggest special emphasis should be put into (1) inhomogeneous SMF (with a detailed characterization of both, intensities and gradients), and (2) combination of SMF (homogeneous and inhomogeneous) with RF EMF, with a fine sweeping of the parameters within the range of typical exposures (in opposition to testing single combinations). In even more complicated experiments, ELF EMF should also be considered 12 since their relationship with SMF have been long known 53. Lastly it is worth mentioning that substantial reduction

of the SMF from MP could be accomplished by replacing the standard electrodynamic loudspeakers and microphones (which include a permanent magnet) by their capacitive counterparts (which do not). Open access paper: <https://www.nature.com/articles/s41598-021-98083-9>

RADIO-FREQUENCY ELECTROMAGNETIC FIELDS: SIMULTANEOUS EXPOSURE TO INFINITE SOURCES FROM TYPICAL BASE STATIONS

A. Linhares, A. P. de Azevedo and L. C. Fernandes. Radio-Frequency Electromagnetic Fields: Simultaneous Exposure to Infinite Sources From Typical Base Stations. IEEE Antennas and Propagation Magazine. doi: 10.1109/MAP.2021.3106829.

Introduction

This article shows that simultaneous exposure to infinite electromagnetic field (EMF) sources from typical base stations transmitting at multiple frequency bands is lower than international limits. Considering a real base station density and conservative path loss models, we show that the power density at a given location can be modeled by a convergent infinite series. To validate this model, we apply it to Brasília, Brazil, and compare the results with measurements. While the proposed model shows that the exposure ratio (ER) will not exceed 2.62%, the highest measured value reaches 85% of this estimation.

Conclusions

In this article, we proposed a method to estimate an upper limit for the TER due to infinite sources of typical base stations. When assessing human exposure to RF EMFs due to specific base stations, the exposure caused by all other base stations can be estimated using the model. The model was applied using base station density data from Brasília and shows that simultaneous exposure from infinite cellular macro base stations distributed as shown in Figure 2 will not be more than 2.62% of the ICNIRP limits. It was validated using simulations and more than 3,000 measurements, and our results agree with them regardless of the probe location. However, for a more conservative approach, when assessing individual base stations, the model can be used to estimate an upper limit for all other base stations (i.e., it provides an exposure addition to the results of the base station under test).

The model is not bounded to a specific frequency band and generic enough to be used by superimposing different categories of base stations, such as micro and macro cells. It may also be employed for 5G, but due to its antenna beamforming, the approach would be more conservative since RF EMF exposure may decrease due to its dynamic directional characteristics. This topic will be the subject of future studies. To ensure the reproducibility of the results in this article, the equations were implemented in numeric code and are available in [20]. The code and data to generate the results in Tables 1 and 2 and Figures 4, 5, 7, 8, and 9 are also available. <https://ieeexplore.ieee.org/document/9532004>

ALTERATION OF INTRAPANCREATIC SEROTONIN, HOMOCYSTEINE, TNF- α , AND NGF LEVELS AS PREDISPOSING FACTORS FOR DIABETES FOLLOWING EXPOSURE TO 900-MHZ WAVES

Gholamali Jelodar, Mansour Azimzadeh, Fatemeh Radmard, Narges Darvishhoo. Alteration of intrapancreatic serotonin, homocysteine, TNF- α , and NGF levels as predisposing factors for diabetes following exposure to 900- MHz waves. *Toxicol Ind Health*. 2021 Aug;37(8):496-503. doi: 10.1177/07482337211022634.

Abstract

Exposure to mobile phone radiation causes deleterious health effects on biological systems. The objects of this study were to investigate the effect of 900-MHz radiofrequency waves (RFW) emitted from base transceiver station antenna on intrapancreatic homocysteine (Hcy), tumor necrosis factor- α (TNF- α), and nerve growth factor (NGF) as predisposing factors involved in pancreatic beta cell damage. Thirty male rats (Sprague-Dawley, 200 $\pm$ 10 g) were randomly divided into the control (without any exposure) and exposed groups: short time (2 h/day), long time (4 h/day), and exposed to 900-MHz RFW for 30 consecutive days. On the last days of the experiment, animals were killed and pancreas tissue was dissected out for evaluation of serotonin, Hcy, TNF- α , and NGF. There was a significant decrease in the serotonin and NGF levels in the pancreatic tissue of exposed groups compared to the control group ($p < 0.05$). Also, the levels of serotonin and NGF in the long-time exposure were significantly lower than the short-time exposure ($p < 0.05$). However, levels of Hcy and TNF- α were significantly increased in the pancreas of exposed groups compared to the control groups ($p < 0.05$). Exposure to 900-MHz RFW decreased pancreatic NGF and serotonin levels and increased the proinflammatory markers (Hcy and TNF- α), which can be a predisposing factor for type 2 diabetes. <https://pubmed.ncbi.nlm.nih.gov/34151670/>

THE RESEARCH AND SAFETY DISCUSSION OF ELECTROMAGNETIC RADIATION IN ELECTRIC VEHICLES

X. Wang et al. The Research and Safety Discussion of Electromagnetic Radiation in Electric Vehicles. 2021 International Applied Computational Electromagnetics Society Symposium (ACES), 2021, pp. 1-3, doi: 10.1109/ACES53325.2021.00038.

Abstract

To study the electromagnetic radiation in electric vehicles for human exposure, the electric field (6 Hz to 3 GHz) and magnetic field (6 Hz to 30 MHz) strengths in electric vehicles of driving mode, charging mode and stationary mode are measured, and the characteristics of the electric and magnetic field in electric vehicles are analysed and studied here. The studies show that the electromagnetic radiation of electric vehicles is mainly consisted of electromagnetic wave from 6 Hz to 500 Hz, and it might be higher in driving mode than that of charging mode and stationary mode. The abnormal values of electromagnetic radiation in electric vehicles often occur in driving mode, and usually appear in the front of the electric vehicles. <https://ieeexplore.ieee.org/document/9528788>

OXIDATIVE STRESS AND NADPH OXIDASE: CONNECTING ELECTROMAGNETIC FIELDS, CATION CHANNELS AND BIOLOGICAL EFFECTS

Christos D Georgiou, Lukas H Margaritis. Oxidative Stress and NADPH Oxidase: Connecting Electromagnetic Fields, Cation Channels and Biological Effects. *Int J Mol Sci.* 2021 Sep 17;22(18):10041. doi: 10.3390/ijms221810041.

Abstract

Electromagnetic fields (EMFs) disrupt the electrochemical balance of biological membranes, thereby causing abnormal cation movement and deterioration of the function of membrane voltage-gated ion channels. These can trigger an increase of oxidative stress (OS) and the impairment of all cellular functions, including DNA damage and subsequent carcinogenesis. In this review we focus on the main mechanisms of OS generation by EMF-sensitized NADPH oxidase (NOX), the involved OS biochemistry, and the associated key biological effects.

Conclusions

On the basis of the above findings, an EMF mechanism can involve ROS formation due to membrane and voltage-gated cation channel function deterioration [2,3,7,8] followed by stress activation and heat-shock protein overexpression [56], which may be associated with behavioural and physiological effects such as blood– brain barrier disruption, memory malfunction, changes in gene expression [53], autophagy, apoptosis [53,84] (especially due to modulation [85]), lifespan reduction, DNA damage, and cancer [18].

Theory and Research Perspectives for a Conclusive Linking of EMFs with ROS/RNS

EMF induction of OS via increased concentration of free radicals, has been challenged (by ICNIRP) mainly due to (i) the claimed non-ionizing nature of EMF (ELF/Rf), where no covalent bonds are broken at non-thermal intensities, or so the argument goes, and because (ii) the measurement of OS is performed by non-specific methods. Indeed, OS is measured, either by methods that are not specific to the identification of generated free radicals, or indirectly by certain oxidative modifications they cause on key biological molecules (e.g., DNA damage, lipid/protein peroxidation, etc.).

Man-made EMFs do not possess high enough energy to generate free radicals, e.g., on freely moving single H₂O molecules by a single photon. However, the individual EMF of such photons are fully synchronized (in terms of frequency, polarization, phase, and propagation direction), thereby producing cumulative macroscopic electric and magnetic fields and electromagnetic radiation (EMR). Nonetheless, these may be additively high enough to break covalent bonds and may directly generate free radicals. Secondly, concentrations of naturally occurring free radicals can increase by the prevention of either (i) reactions between them (e.g., appearing as the aforementioned reactants $\bullet\text{NO} + \text{O}_2\bullet^-$ and products $\bullet\text{OH} + \bullet\text{NO}_2$), or (ii) the reassociation of free radical pairs generated enzymically as transition states. Such prevention can be assisted by the EMF-induced free radical pair mechanism [86,87]. Here, EMFs can prevent the reassociation of free radical pairs by reversing the spin direction of the single electron in one of these free radicals by flipping the direction of its magnetic field component. Thus, the magnetically affected free radical pair ends up consisting of two free radicals, the electron spins of which have become parallel, thereby preventing their re-binding and indirectly increasing their concentration. The free radical pair mechanism has

been accounted for by the International Agency for Research on Cancer for the classification of the RF EMFs in the Group 2B category of “possibly carcinogenic to humans” [88]. These two mechanisms of free radical-concentration increase corroborate with the preliminary finding that ELF EMFs increase the concentration of $O_2^{\bullet-}$ by many fold in various organs of mice exposed to the ICNIRP limit of 100 μT at 50 Hz (pending publication by Dr. Georgiou’s lab). Therefore, methods for the in vivo specific detection of the key biological free radicals $\bullet OH$ and $O_2^{\bullet-}$ ([89,90]) are needed in order to unequivocally prove the generation of carcinogenic OS by EMFs. Open access paper: <https://www.mdpi.com/1422-0067/22/18/10041>

RESIDENTIAL EXTREMELY LOW FREQUENCY MAGNETIC FIELDS AND SKIN CANCER

Muhammad Waseem Khan, Jukka Juutilainen, Jonne Naarala, Päivi Roivainen. Residential extremely low frequency magnetic fields and skin cancer. *Occup Environ Med.* 2021 Sep 30; oemed-2021-107776. doi: 10.1136/oemed-2021-107776.

Abstract

Objective: Photoinduced radical reactions have a fundamental role in skin cancer induced by ultraviolet radiation, and changes in radical reactions have also been proposed as a mechanism for the putative carcinogenic effects of extremely low frequency (ELF) magnetic fields (MFs). We assessed the association of melanoma and squamous cell carcinoma with residential MF exposure.

Methods: All cohort members had lived in buildings with indoor transformer stations (TSs) during the period from 1971 to 2016. MF exposure was assessed based on apartment location. Out of the 225 492 individuals, 8617 (149 291 person-years of follow-up) living in apartments next to TSs were considered as exposed, while individuals living on higher floors of the same buildings were considered as referents. Associations between MF exposure and skin cancers were examined using Cox proportional hazard models.

Results: The HR for MF exposure ≥ 6 month was 1.05 (95% CI 0.72 to 1.53) for melanoma and 0.94 (95% CI 0.55 to 1.61) for squamous cell carcinoma. Analysis of the age at the start of residence showed an elevated HR (2.55, 95% CI 1.15 to 5.69) for melanoma among those who lived in the apartments when they were less than 15 years old. This finding was based on seven exposed cases.

Conclusions: The results of this study suggested an association between childhood ELF MF exposure and adult melanoma. This is in agreement with previous findings suggesting that the carcinogenic effects of ELF MFs may be associated particularly with childhood exposure <https://pubmed.ncbi.nlm.nih.gov/34593542/>

ELECTROMAGNETIC FIELDS MODIFY REDOX BALANCE IN THE RAT GASTROINTESTINAL TRACT

Sieroń Karolina, Knapik Katarzyna, Onik Grzegorz, Romuk Ewa, Birkner Ewa, Kwiatek Sebastian, Sieroń Aleksander. Electromagnetic Fields Modify Redox Balance in the Rat Gastrointestinal Tract. *Frontiers in Public Health*. 9:1283. 2021. DOI:10.3389/fpubh.2021.710484.

Abstract

Objective: The aim of the study was to assess the influence of electromagnetic fields with divergent physical properties on the prooxidative and antioxidative balances in homogenates of the tongue, salivary glands, esophagus, stomach, and small and large intestines of rats.

Material and Methods: Forty rats were randomly divided into four equal groups, namely, a control group, a group exposed to low-frequency electromagnetic fields (LF-EMFs; frequency: 50 Hz; intensity: 10 kV/m; magnetic induction: 4.3 pT), a group exposed to radiofrequency electromagnetic fields (RF-EMFs) emitted by mobile phones (frequency: 900 MHz), and a group exposed simultaneously to LF-EMFs and RF-EMFs emitted by mobile phones. After 28 consecutive days of the experiment, the following pro- and antioxidative markers were assessed in the gastrointestinal tract homogenates: superoxide dismutase (SOD) and its two isoenzymes (Mn- SOD, Cu,Zn-SOD) catalase (CAT), glutathione peroxidase (GPx), glutathione reductase (GR), glutathione S- transferase (GST), total antioxidative capacity (TAC), total oxidative status (TOS), and malondialdehyde (MDA).

Results: In rats exposed to LF-EMFs, higher concentrations of the markers of prooxidant processes, MDA or TOS, were observed in the salivary glands, esophagus, and small intestine homogenates in comparison with the control group. Additionally, in the group of rats opposite to the control, antioxidant activity was observed. The main differences included a higher activity of Cu,Zn-SOD in homogenates of the tongue, salivary glands, and esophagus as well as decreased activity of CAT in homogenates of the tongue, esophagus, and small intestine. In animals exposed to RF-EMFs, the concentration of TOS was higher in the large intestine than in control rats. The main difference of antioxidant activity was presented by decreased Cu,Zn-SOD in homogenates of the salivary glands, stomach, small and large intestine as well as CAT in homogenates of the tongue, esophagus, stomach, and small and large intestine. Moreover, in rats exposed simultaneously to LF-EMFs and RF-EMFs, a lower concentration of TOS was observed. Antioxidant activity was presented by a decreased activity of CAT in homogenates of the tongue, esophagus, stomach, and small and large intestine in comparison to the control group.

Conclusion: Among those applied in the study, electromagnetic fields of a low-frequency caused the most significant disturbances of oxidative stress in the rat gastrointestinal tract. Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.710484>

NEAR FIELD RADIO-FREQUENCY ELECTROMAGNETIC FIELD EXPOSURE OF A WESTERN HONEY BEE

D. Toribio, W. Joseph, A. Thielens. Near Field Radio-Frequency Electromagnetic Field Exposure of a Western Honey Bee. IEEE Transactions on Antennas and Propagation, doi: 10.1109/TAP.2021.3111286.

Abstract

It has been suggested that the wireless network evolution to smaller carrier wavelengths (from 2G to 5G) increases radio-frequency electromagnetic field (RF-EMF) absorption in Western Honey Bees (*Apis mellifera*). It is unknown whether the radiation performance of antennas is stable when an insect appears in their vicinity. In this research, the absorbed power in a worker honey bee and the influence of the bee's presence on antennas' radiation performance is investigated for the newly used frequencies in 5G networks, from 6-240 GHz. To these aims, numerical simulations using the finite-difference time-domain method were performed, in which a bee model, obtained by micro-CT scanning, was employed. These simulations showed that in the near field, the absorbed power can increase by a factor of 53, from 6-240 GHz. This is a factor of 7 higher than the increase reported in the far field, in previous studies. Furthermore, the simulations revealed that antennas' radiation efficiency can decrease by up to -40 % when a bee appears in the near field. Likewise, it was found that the gain pattern depends on the separation distance between the bee and the antenna, with a stronger dependency for higher frequencies.

CONCLUSION

Numerical simulations using finite-difference time-domain analysis were executed to calculate the radio- frequency electromagnetic fields (RF-EMFs) in and around a worker Western Honey Bee (*Apis mellifera*). This lead to the quantification of the whole-body averaged absorbed radio-frequency power Pabs, under near-field exposure, in the frequency range of 6-240 GHz. The simulations showed that, in the near field, Pabs decreases as the separation distance between the bee and the dipole increases, and increases as frequency increases. The frequency behavior of Pabs in the near field is thus different to its far-field behavior, since, for a given accepted power and distance, in the near field Pabs increases with frequency (an average of 30.5 dB from 6-240 GHz), while in the far field Pabs in the worker bee is maximized at 12 GHz. This near-field increase in Pabs as a function of frequency, can be a factor of 7 higher than the far-field increase.

Moreover, these simulations allowed the study of the influence of the bee's position ($0.1 - 10 \lambda$ from the dipoles) in the radiation performance of the dipoles. In particular, it was found that as the separation distance increases, the dipoles' isotropic gain decreases and approaches their free-space gain. Also, it was noted that as a result of the near-field frequency behavior of Pabs, the dipoles' radiation efficiency decreases with increasing frequency and increases with increasing separation distance. Additionally, it was acknowledged that the gain pattern in the direction where the bee approaches the antenna depends on the separation distance between the bee and the antenna, with a stronger dependency for higher frequencies. This is important for 5G telecommunication networks, because they rely on infrastructure (antennas) whose performance can be influenced by the presence of free-flying insects.

<https://ieeexplore.ieee.org/document/9539081>

SUBLETHAL NEONICOTINOID EXPOSURE ATTENUATES THE EFFECTS OF ELECTROMAGNETIC FIELDS ON HONEY BEE FLIGHT AND LEARNING

S. Shepherd, M.A.P. Lima, E.E. Oliveira, S.M. Sharkh, H. Aonuma, C.W. Jackson, P.L. Newland. Sublethal neonicotinoid exposure attenuates the effects of electromagnetic fields on honey bee flight and learning, *Environmental Advances*. Volume 4. 2021. 100051. doi:10.1016/j.envadv.2021.100051.

Abstract

Many environmental stressors are currently implicated in the decline of flying insects in general, and important pollinator species such as the honey bee. Recent studies have shown that extremely low frequency electromagnetic fields (ELF EMFs) affect many aspects of insect behaviour and cognition which raises the potential that ELF EMFs could interact with other environmental stressors such as neonicotinoid insecticides to have even greater impact on the decline of flying insects. Here we analysed the effects of individual and combined exposure of the honey bee to 50 Hz EMFs and sublethal exposure to clothianidin, in a tethered flight assay and an olfactory appetitive learning assay. Clothianidin was significantly toxic to bees and exposure to field-realistic levels (2.00 ng/bee) led to 25% mortality. Exposure to ELF EMFs alone led to a significant increase in wingbeat frequency at levels above 100 μ T. Prior exposure to clothianidin attenuated the effects of EMFs on wingbeat frequency. Exposure to EMFs alone reduced learning of a proboscis extension reflex (PER). Prior exposure to low doses of clothianidin attenuated the effects of EMFs on PER. These results indicate no evidence of synergy between clothianidin and ELF EMFs as environmental stressors but suggest the potential for EMFs to affect the same susceptible fraction of the bee population that have been affected by clothianidin. Results lay the foundation to further explore the interactions of ELF EMFs with other environmental stressors and consider the key factors that may make bees susceptible to ELF EMFs.

Open access paper: <https://www.sciencedirect.com/science/article/pii/S2666765721000223>

EFFECTS OF LONG-TERM EXPOSURE TO L-BAND (~2.0 GHZ) HIGH-POWER (0.5, 1.0, AND 1.5 W/M2) MICROWAVE ON THE BRAIN FUNCTION OF MALE MICE

Lin Y, Gao P, Guo Y, Chen Q, Lang H, Guo Q, Miao X, Li J, Zeng L, Guo G. Effects of Long-Term Exposure to L-Band High-Power Microwave on the Brain Function of Male Mice. *Biomed Res Int*. 2021 Sep 4;2021:2237370. doi: 10.1155/2021/2237370.

Abstract

Currently, the impact of electromagnetic field (EMF) exposure on the nervous system is an increasingly arousing public concern. The present study was designed to explore the effects of continuous long-term exposure to L- band (~ 2.0 GHz) high-power microwave (L-HPM) on brain function and related mechanisms. Forty-eight male Institute of Cancer Research (ICR) mice were exposed to L-HPM at various power densities (0.5, 1.0, and 1.5 W/m²) and the brain function was examined at different time periods after exposure. The morphology of the brain was examined by hematoxylin-eosin (HE) and deoxynucleotidyl transferase-mediated dUTP nick-end labeling (TUNEL) staining. Furthermore, cholinergic markers, oxidative stress markers, and the expression of c- fos were evaluated to identify a "potential" mechanism. The results showed that exposure to L-HPM at 1.5 W/m² can cause generalized injuries in the hippocampus (CA1 and CA3) and cerebral cortex (the first somatosensory cortex) of mice, including cell apoptosis, cholinergic dysfunction, and oxidative damage.

Moreover, the deleterious effects were closely related to the power density and exposure time, indicating that long-term and high-power density exposure may be detrimental to the nervous system.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8437633/>

ELECTROMAGNETIC POLLUTION AS A POSSIBLE EXPLANATION FOR THE DECLINE OF HOUSE SPARROWS IN INTERACTION WITH OTHER FACTORS

Alfonso Balmori. Electromagnetic Pollution as a Possible Explanation for the Decline of House Sparrows in Interaction with Other Factors. *Birds* 2021, 2(3), 329-337; doi: 10.3390/birds2030024

Simple Summary

In recent decades, there has been a decline of the house sparrow (*Passer domesticus*), mainly in European cities, and several hypotheses have been proposed. The objective of this article is to delve into the reasons why an increase in electromagnetic radiation especially in cities, may be intervening in some way. Previous studies indicated that house sparrows were significantly negatively associated with increasing electromagnetic radiation and sparrows disappeared from areas most polluted. Electromagnetic radiation is the most plausible factor and is the only one that affects the other hypotheses proposed so far. Additionally, the recent sparrow decline matches the deployment of mobile telephony networks. For these reasons, electromagnetic radiation is not only a plausible but a probable hypothesis that must be seriously considered, probably in synergy with the other factors previously proposed.

Abstract

In recent decades, there has been a decline of the House Sparrow (*Passer domesticus*), mainly in European cities, and several hypotheses have been proposed that attempt to determine the causes of this rapid decline. Previous studies indicated that house sparrows were significantly negatively associated with increasing electromagnetic radiation and sparrows disappeared from areas most polluted. In addition, there are many studies on the impact of radiation on other bird and non-bird species, as well as numerous laboratory studies that demonstrated detrimental effects at electric field strength levels that can be found in cities today.

Electromagnetic radiation is the most plausible factor for multiple reasons, including that this is the only one that affects the other hypotheses proposed so far. It is a type of pollution that affects productivity, fertility, decreases insects (chicken feed), causes loss of habitat, decreases immunity and can promote disease.

Additionally, the recent sparrow decline matches the deployment of mobile telephony networks. Further, there are known mechanisms of action for non-thermal effects of electromagnetic radiation that may affect sparrows causing their decline. Thus, electromagnetic radiation must be seriously considered as a factor for house sparrows' decline, probably in synergy with the other factors previously proposed.

Open access paper: <https://www.mdpi.com/2673-6004/2/3/24/htm>

ONE-TIME ELECTROMAGNETIC IRRADIATION MODIFIES STRESS-SENSITIVE GENE EXPRESSIONS IN RICE PLANT

Ardhendu Kundu, Sathish Vangaru, Sucharita Bhowmick, Somnath Bhattacharyya, Amirul I Mallick, Bhaskar Gupta. One-time Electromagnetic Irradiation Modifies Stress-sensitive Gene Expressions in Rice Plant. Bioelectromagnetics. 2021 Sep 24. doi: 10.1002/bem.22374.

Abstract

Electromagnetic energy is utilized over multiple frequency bands to provide seamless wireless communication services. Plants can well perceive electromagnetic energy present in open environment due to reasonably high permittivity and electrical conductivity of constituent tissues. Moreover, higher surface-to-volume ratio of plant structure facilitates increased interaction with the incident electromagnetic waves. To date, a few well-designed studies have been conducted inside controlled electromagnetic reverberation chambers to investigate either short duration-low amplitude or long duration-periodic electromagnetic irradiation-induced molecular responses in plants. However, as far as is known, studies investigating molecular responses particularly at the mid-vegetative stage in plants following one-time (hours-long) electromagnetic irradiation have not been reported earlier. Hence, the present study aimed at investigating molecular responses in 40-day-old Swarnaprabha rice plants following one-time 1837.50 MHz, 2.75 mW/m² electromagnetic irradiation of 2 h 30 min duration. Controlled electromagnetic irradiation inside a simple reverberation chamber was ensured to achieve pure electromagnetic environment at 1837.50 MHz with deterministic electromagnetic power density at selected position. Swarnaprabha rice plant was chosen for this investigation since the rice variety is widely cultivated and consumed in the Indian subcontinent. Subsequent alterations in some selected stress-sensitive gene expressions were assayed using real-time quantitative polymerase chain reaction technique-significant upregulation in calmodulin and phytochrome B gene expressions were noted. This investigation was purposefully focused on subsequent molecular responses immediately following electromagnetic irradiation so that the possible effects of secondary stimulations could be avoided. Observed molecular responses strongly suggested that plants perceive 1837.50 MHz, 2.75 mW/m² electromagnetic irradiation similar to other injurious stimuli. <https://pubmed.ncbi.nlm.nih.gov/34559898/>

SHORT-TERM EXTREMELY LOW-FREQUENCY ELECTROMAGNETIC FIELD INHIBITS SYNAPTIC PLASTICITY OF SCHAFER COLLATERAL- CA1 SYNAPSES IN RAT HIPPOCAMPUS VIA THE CA 2+/CALCINEURIN PATHWAY

Pei Xia, Yu Zheng, Lei Dong, Chunxiao Tian. Short-Term Extremely Low-Frequency Electromagnetic Field Inhibits Synaptic Plasticity of Schaffer Collateral-CA1 Synapses in Rat Hippocampus via the Ca²⁺/Calcineurin Pathway. ACS Chem Neurosci. 2021 Sep 9. doi: 10.1021/acchemneuro.1c00500.

Abstract

In this study, we investigate the intrinsic mechanism by which an extremely low-frequency electromagnetic field (ELF-EMF) influences neurons in the Schaffer collateral-CA1 (SC-CA1) region of rat hippocampus using electrophysiological techniques. ELF-EMF has an interesting effect on synaptic plasticity: it weakens long-term potentiation and enhances long-term depression. Here, the magnetic field effect disappeared after a blockade of voltage-gated calcium channels and calcineurin, which are key components in the Ca²⁺/calcineurin pathway,

with two blockers, cadmium chloride and cyclosporin A. This fully establishes that the effect of ELF-EMF on synaptic plasticity is mediated by the Ca²⁺/calcineurin pathway and represents a novel technique for studying the specific mechanisms of action of ELF-EMF on learning and memory. <https://pubmed.ncbi.nlm.nih.gov/34498467/>

DNMT1 AND MIRNAS: POSSIBLE EPIGENETICS FOOTPRINTS IN ELECTROMAGNETIC FIELDS UTILIZATION IN ONCOLOGY

Mohadeseh Shayeghan, Flora Forouzesh, Alireza Madjid Ansari, Mohammad Amin Javidi. DNMT1 and miRNAs: possible epigenetics footprints in electromagnetic fields utilization in oncology. Med Oncol. 2021 Sep 8;38(10):125. doi: 10.1007/s12032-021-01574-y.

Abstract

Many studies were performed to unravel the effects of different types of Electromagnetic fields (EMFs) on biological systems. Some studies were conducted to exploit EMFs for medical purposes mainly in cancer therapy. Although many studies suggest that the EMFs exposures can be effective in pre-clinical cancer issues, the treatment outcomes of these exposures on the cancer cells, especially at the molecular level, are challenging and overwhelmingly complicated yet. This article aims to review the epigenetic mechanisms that can be altered by EMFs exposures with the main emphasis on Extremely low frequency electromagnetic field (ELF-EMF). The epigenetic mechanisms are reversible and affected by environmental factors, thus, EMFs exposures can modulate these mechanisms. According to the reports, ELF-EMF exposures affect epigenetic machinery directly or through the molecular signaling pathways. ELF-EMF in association with DNA methylation, histone modification, miRNAs, and nucleosome remodeling could affect the homeostasis of cancer cells and play a role in DNA damage repairing, apoptosis induction, prevention of metastasis, differentiation, and cell cycle regulation. In general, the result of this study shows that ELF-EMF exposure probably can be effective in cancer epigenetic therapy, but more molecular and clinical investigations are needed to clarify the safe and specific dosimetric characteristics of ELF-EMF in practice. <https://pubmed.ncbi.nlm.nih.gov/34495398/>

SAFETY OF WIRELESS POWER TRANSFER

James C. Lin. Safety of Wireless Power Transfer. IEEE Access. 10 September 2021. doi: 10.1109/ACCESS.2021.3108966.

Abstract

Wireless power transfer (WPT) systems are developed to provide electric power either directly or via battery- charging. The optimism on WPT technology is driven by the ubiquity of cellphones, laptops, and mobile communication devices. Aside from not having to plug in a cellphone or laptop, WPT battery charging offers the potential for mobile devices to get electrical power the same way they get data through harvesting ambient electromagnetic radiation. The dream is a truly wireless mobility scenario with tether-free electric power for cellphones, laptops, appliances, and transportation systems. Beyond wireless communication, the electromagnetic power required for large-scale commercial WPT implementation is substantial. A key facet of the system design and research should include consideration of health effects and safety of radiofrequency electromagnetic radiation.

<https://ieeexplore.ieee.org/document/9535368>

CELLULAR EFFECTS OF TERAHERTZ WAVES

Cherkasova OP, Serdyukov DS, Nemova EF, Ratushnyak AS, Kucheryavenko AS, Dolganova IN, Xu G, Skorobogatiy M, Reshetov IV, Timashev PS, Spektor IE, Zaytsev KI, Tuchin VV. Cellular effects of terahertz waves. J Biomed Opt. 2021 Sep;26(9). doi: 10.1117/1.JBO.26.9.090902.

Abstract

Significance: An increasing interest in the area of biological effects at exposure of tissues and cells to the terahertz (THz) radiation is driven by a rapid progress in THz biophotonics, observed during the past decades. Despite the attractiveness of THz technology for medical diagnosis and therapy, there is still quite limited knowledge about safe limits of THz exposure. Different modes of THz exposure of tissues and cells, including continuous-wave versus pulsed radiation, various powers, and number and duration of exposure cycles, ought to be systematically studied.

Aim: We provide an overview of recent research results in the area of biological effects at exposure of tissues and cells to THz waves.

Approach: We start with a brief overview of general features of the THz-wave-tissue interactions, as well as modern THz emitters, with an emphasis on those that are reliable for studying the biological effects of THz waves. Then, we consider three levels of biological system organization, at which the exposure effects are considered:

- (i) solutions of biological molecules;
- (ii) cultures of cells, individual cells, and cell structures; and
- (iii) entire organs or organisms; special attention is devoted to the cellular level.

We distinguish thermal and nonthermal mechanisms of THz-wave-cell interactions and discuss a problem of adequate estimation of the THz biological effects' specificity. The problem of experimental data reproducibility, caused by rareness of the THz experimental setups and an absence of unitary protocols, is also considered.

Results: The summarized data demonstrate the current stage of the research activity and knowledge about the THz exposure on living objects.

Conclusions: This review helps the biomedical optics community to summarize up-to-date knowledge in the area of cell exposure to THz radiation, and paves the ways for the development of THz safety standards and THz therapeutic applications.

Conclusion

Recent research results in the area of THz-wave effects on biological systems of the different organization levels, such as biomolecules, cells, and organism, were discussed. Despite the considerable data accumulated in this demanding research direction, we still possess quite limited knowledge about biological effects of THz waves.

Further research and engineering efforts are required to develop adequate safe limits of THz-wave exposure and to objectively uncover strengths and weaknesses of THz technology in different branches of medial therapy.

Thus, this review summarizes up-to-date knowledge in the area of cell exposure to THz radiation and paves the ways to the THz dosimetry and therapeutic avenues. Open access paper:

<https://www.spiedigitallibrary.org/journals/journal-of-biomedical-optics/volume-26/issue-09/090902/Cellular-effects-of-terahertzwaves/10.1117/1.JBO.26.9.090902.full?SSO=1>

NUMERICAL MODELLING AND EXPERIMENTAL VERIFICATION OF THERMAL EFFECTS IN LIVING CELLS EXPOSED TO HIGH-POWER PULSES OF THZ RADIATION

D S Sitnikov, A A Pronkin, I V Ilina, V A Revkova, M A Konoplyannikov, V A Kalsin, V P Baklaushev. Numerical modelling and experimental verification of thermal effects in living cells exposed to high-power pulses of THz radiation. Sci Rep. 2021 Sep 9;11(1):17916. doi: 10.1038/s41598-021-96898-0.

Abstract

Exposure of cells or biological tissues to high-power pulses of terahertz (THz) radiation leads to changes in a variety of intracellular processes. However, the role of heating effects due to strong absorption of THz radiation by water molecules still stays unclear. In this study, we performed numerical modelling in order to estimate the thermal impact on water of a single THz pulse as well as a series of THz pulses. A finite-element (FE) model that provides numerical solutions for the heat conduction equation is employed to compute the temperature increase. A simple expression for temperature estimation in the center of the spot of THz radiation is presented for given frequency and fluence of the THz pulse. It has been demonstrated that thermal effect is determined by either the average power of radiation or by the fluence of a single THz pulse depending on pulse repetition rate. Human dermal fibroblasts have been exposed to THz pulses (with an energy of [Formula: see text] and repetition rate of 100 Hz) to estimate the thermal effect. Analysis of heat shock proteins expression has demonstrated no statistically significant difference ([Formula: see text]) between control and experimental groups after 3 h of irradiation.

Open access paper: <https://www.nature.com/articles/s41598-021-96898-0>

ELECTRIC FIELD DETECTION AS FLORAL CUE IN HOVERFLY POLLINATION

Khan SA, Khan KA, Kubik S, Ahmad S, Ghramh HA, Ahmad A, Skalicky M, Naveed Z, Malik S, Khalofah A, Aljedani DM. Electric field detection as floral cue in hoverfly pollination. Sci Rep. 2021 Sep 21;11(1):18781. doi: 10.1038/s41598-021-98371-4.

Abstract

Pollinators can detect the color, shape, scent, and even temperature of the flowers they want to visit. Here, we present the previously unappreciated capacity of hoverflies (*Eristalis tenax* and *Cheilosia albipila*) to detect the electric field surrounding flowers. Using hoverflies as key dipteran pollinators, we explored the electrical interactions between flies and flowers-how a hoverfly acquired a charge and how their electrical sensing ability for target flowers contributed to nectar identification and pollination. This study revealed that rapid variations in a floral electric field were related to a nectar reward and increased the likelihood of the fly's return visits. We found that thoracic hairs played a role in the polarity of hoverfly charge, revealing their electro-mechanosensory capability, as in bumblebees (*Bombus terrestris*). Electrophysiological analysis of the hoverfly's antennae did not reveal neural sensitivity to the

electric field, which favors the mechanosensory hairs as putative electroreceptive organs in both species of hoverflies.

Open access paper: <https://www.nature.com/articles/s41598-021-98371-4>

EFFECTS OF MOBILE PHONE RADIOFREQUENCY RADIATION ON SPERM QUALITY (REVIEW)

Romualdo Sciorio, Luca Tramontano, Sandro C Esteves. Effects of mobile phone radiofrequency radiation on sperm quality. *Zygote*. 2021 Aug 13;1-10. doi: 10.1017/S096719942100037X

Abstract

In the last decades, the universal use of mobile phones has contributed to radiofrequency electromagnetic radiation environmental pollution. The steady growth in mobile phone usage has raised concerns about the effects of phone radiation on male reproductive health. Epidemiological studies report a sharp decline in sperm counts in developing countries, and worldwide with c. 14% of couples having difficulties to conceive, many of which are attributed to a male infertility factor. Environment and lifestyle factors are known to contribute to male infertility. Exposure to heat, radiation, or radioactivity might induce damage to biological tissue organs, including the testis. Given the ubiquitous use of mobile phones, the potential adverse effects of the resulting environmental radiation needs to be elucidated further. It seems to be an apparent relationship between the increased exposure to mobile phone radiofrequency and sperm quality decline, but the evidence is not conclusive. Our review summarizes the evidence concerning the possible adverse effects of cell phone radiation on the male reproductive system, with a focus on sperm quality. Also, we critically analyze the effects of elevated testicular temperature and oxidative stress on male fertility and how these factors could interfere with the physiological activities of the testis.

<https://pubmed.ncbi.nlm.nih.gov/34384508/>

Future perspectives and conclusions

The rapid technological advances in personal computers and communication devices might pose a risk for human health. Cell phone devices emit radiofrequency electromagnetic waves that seem to affect male reproductive health and other body functions (McClelland 3rd and Jaboin, 2018; Sage and Burgio, 2018; Wall et al., 2019). Although the current data are not unequivocal, it seems sound to speculate that mobile phone exposure might be contributing to subfertility. However, the existing evidence primarily relates to adverse effects on sperm motility and morphology, which are limited endpoints for evaluating the male fertility potential.

The exact mechanisms of how RF-EMR might affect the testis, epididymis, and sperm have not yet been fully understood. Additional studies are warranted, particularly prospective studies assessing sperm functional markers, such as sperm DNA integrity and OS, in fertile and subfertile men. Equally important will be to analyze whether the decreased sperm quality associated with mobile phone exposure translates into impaired pregnancy chances. The effects of short-term and long-term exposure and energy intensity should be also investigated in more detail, taking into account relevant confounders. Only then will scientific societies and regulatory bodies be able to provide users with transparent information concerning the risks and guidance for proper use.

THRESHOLD OF RADIOFREQUENCY ELECTROMAGNETIC FIELD EFFECT ON HUMAN BRAIN (REVIEW)

Hiie Hinrikus, Jaanus Lass, Maie Bachmann. Threshold of radiofrequency electromagnetic field effect on human brain. *Int J Radiat Biol.* 2021 Aug 17;1-38. doi: 10.1080/09553002.2021.1969055.

Abstract

Purpose: This review aims to estimate the threshold of radiofrequency electromagnetic field (RF EMF) effects on human brain based on analyses of published research results. To clarify the threshold of the RF EMF effects, two approaches have been applied: 1) the analyses of restrictions in sensitivity for different steps of the physical model of low-level RF EMF mechanism and 2) the analyses of experimental data to clarify the dependence of the RF EMF effect on exposure level based on the results of published original neurophysiological and behavioral human studies for 15 years 2007-2021.

Conclusions: The analyses of the physical model of non-thermal mechanisms of RF EMF effect leads to conclusion that no principal threshold of the effect can be determined. According to the review of experimental data, the rate of detected RF EMF effects is 76.7% in resting EEG studies, 41.7% in sleep EEG and 38.5% in

Behavioral studies : The changes in EEG probably appear earlier than alterations in behavior become evident. The lowest level of RF EMF at which the effect in EEG was detected is 2.45 V/m (SAR =0.003 W/kg). There is a preliminary indication that the dependence of the effect on the level of exposure follows rather field strength than SAR alterations. However, no sufficient data are available for clarifying linearity-nonlinearity of the dependence of effect on the level of RF EMF. The finding that only part of people are sensitive to RF EMF exposure can be related to immunity to radiation or hypersensitivity. The changes in EEG caused by RF EMF appeared similar in the majority of analyzed studies and similar to these in depression. The possible causal relationship between RF EMF effect and depression among young people is highly important problem. <https://pubmed.ncbi.nlm.nih.gov/34402382/>

Excerpt

Does the RF EMF has a threshold, lower of which the RF EMF does not have biological effect? This is an important question and crucial to avoid possible consequences on health.

Theoretical estimations for the sensitivity of living cells to electric field provided the threshold values about 10^{-6} – 10^{-7} V/m Hz^{1/2} (Weaver and Astumian, 1990; Hinrikus et al., 1998). In the case of wide-band telecommunication technology, the threshold rises: at 1 MHz bandwidth, the sensitivity is 10^{-3} – 10^{-4} V/m. However, these estimations used a simple single cell model. Realistic model involving combinations of different cells, molecules and partly nonlinear physiological processes is highly complex. To the best of our knowledge, the calculations using complex model have been not performed.

Conclusions

The analyses of the model of the non-thermal mechanism of RF EMF effect shows that the steps of the model contain no principal threshold for the effect. Therefore, the only way to estimate the possible threshold is analyses of experimental data. The review of experimental data of human RF EMF neurophysiological effects results in following main conclusions that indicated the directions of future research:

1)The lowest field strength that has caused the effect in EEG, according to the reviewed studies, is 2.45 V/m (SAR=0.003 W/kg), close to the radio and TV broadcasting RF EMF field strength about 0.1 V/m. The future large- scale human, animal and in vitro studies are required to clarify the level and to increase the reliability of the experimentally determined threshold of RF EMF effect.

2)There is a preliminary indication that the intensity of the effect follows rather the field strength than SAR alteration. However, no sufficient data are available for clarifying regularity and linearity-nonlinearity of the relationship. The studies with systematic variations in exposure level (electric and magnetic field strengths, power density) would help to advance the field. The research from cellular to humans is needed in this direction.

3)Very limited data are available about the repair and adaptive phenomena important in the interpretation of the RF EMF effects on health. Special studies are required to clarify the mechanisms and possibilities of the repair processes.

4)The finding that only part of people are sensitive to RF EMF exposure can be related to immunity to radiation or hypersensitivity. The variability of sensitivity between people and longterm stability of the status required large-scale long-term experimental studies.

5)The indication that the changes in EEG caused by RF EMF appeared similar to these in depression need a special attention. The fast increase of depression with the highest prevalence among young individuals and more intense use of mobile telecommunication technology with prevalence in young people during the same period needs attention. The possible causal relationship between RF EMF effect and depression among young people is the problem of high importance. The methodology and large-scale investigations in this direction are required.

EFFECTS OF RADIOFREQUENCY ELECTROMAGNETIC RADIATION ON NEUROTRANSMITTERS IN THE BRAIN

Cuicui Hu, Hongyan Zuo, Yang Li. Effects of Radiofrequency Electromagnetic Radiation on Neurotransmitters in the Brain. *Frontiers in Public Health*. 9:1139. 2021. doi: 10.3389/fpubh.2021.691880.

Abstract

With the rapid development of electronic information in the past 30 years, technical achievements based on electromagnetism have been widely used in various fields pertaining to human production and life.

Consequently, electromagnetic radiation (EMR) has become a substantial new pollution source in modern civilization. The biological effects of EMR have attracted considerable attention worldwide. The possible interaction of EMR with human organs, especially the brain, is currently where the most attention is focused. Many studies have shown that the nervous system is an important target organ system sensitive to EMR. In recent years, an increasing number of studies have focused on the neurobiological effects of EMR, including the metabolism and transport of neurotransmitters. As messengers of synaptic transmission, neurotransmitters play critical roles in cognitive and emotional behavior. Here, the effects of EMR on the metabolism and receptors of neurotransmitters in the brain are summarized. Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.691880>

THE EFFECT OF EXPOSURE TO RADIOFREQUENCY FIELDS ON CANCER RISK IN THE GENERAL AND WORKING POPULATION: A PROTOCOL FOR A SYSTEMATIC REVIEW OF HUMAN OBSERVATIONAL STUDIES

Susanna Lagorio, Maria Blettner, Dan Baaken, Maria Feychting, Ken Karipidis, Tom Loney, Nicola Orsini, Martin Röösli, Marilia Silva Paulo, Mark Elwood. The effect of exposure to radiofrequency fields on cancer risk in the general and working population: A protocol for a systematic review of human observational studies. *Environ Int.* 2021 Aug 22;157:106828. doi: 10.1016/j.envint.2021.106828.

Highlights

- RF-EMF was classified by IARC as possibly carcinogenic to humans (2B) in May 2011
- A systematic review of all subject-relevant epidemiological studies is now needed.
- A detailed protocol ensures the review's transparency, utility and credibility.
- Original study validity will be evaluated with a customized OHAT risk of bias tool.
- Internal coherence and external plausibility will inform conclusions.

Abstract

Background: The World Health Organization (WHO) has an ongoing project to assess potential health effects of exposure to radiofrequency electromagnetic fields (RF-EMF) in the general and working population. Here we present the protocol for a systematic review of the scientific literature on cancer hazards from exposure to RF-EMF in humans, commissioned by the WHO as part of that project.

Objective: To assess the quality and strength of the evidence provided by human observational studies for a causal association between exposure to RF-EMF and risk of neoplastic diseases.

Eligibility Criteria: We will include cohort and case-control studies investigating neoplasia risks in relation to three types of exposure to RF-EMF: near-field, head-localized, exposure from wireless phone use (SR-A); far-field, whole body, environmental exposure from fixed-site transmitters (SR-B); near/far-field occupational exposures from use of handheld transceivers or RF-emitting equipment in the workplace (SR-C). While no restriction on tumour type will be applied, we will focus on selected neoplasms of the central nervous system (brain, meninges, pituitary gland, acoustic nerve) and salivary gland tumours (SR-A); brain tumours and leukaemias (SR-B, SR-C).

Information sources: Eligible studies will be identified through Medline, Embase, and EMF-Portal.

Risk-of-bias assessment: We will use a tailored version of the OHAT's tool to evaluate the study's internal validity.

Data synthesis: We will consider separately studies on different tumours, neoplasm-specific risks from different exposure sources, and a given exposure-outcome pair in adults and children. When a quantitative synthesis of findings can be envisaged, the main aims of the meta-analysis will be to assess the strength of association and the shape of the exposure-response relationship; to quantify the degree of heterogeneity across studies; and explore the sources of inconsistency (if any). When a meta-analysis is judged inappropriate, we will perform a narrative synthesis, complemented by a structured tabulation of results and appropriate visual displays.

Evidence assessment: Confidence in evidence will be assessed in line with the GRADE approach.

Funding: This project is supported by the World Health Organization. Co-financing was provided by the New Zealand Ministry of Health; the Istituto Superiore di Sanità in its capacity as a WHO Collaborating Centre for Radiation and Health; ARPANSA as a WHO Collaborating Centre for Radiation Protection. Registration: PROSPERO CRD42021236798. <https://pubmed.ncbi.nlm.nih.gov/34433115/>

EFFECTS OF RADIOFREQUENCY FIELD EXPOSURE ON MALE FERTILITY & PREGNANCY & BIRTH OUTCOMES: PROTOCOLS FOR A SYSTEMATIC REVIEW OF EXPERIMENTAL STUDIES IN NON-HUMAN MAMMALS & IN HUMAN SPERM EXPOSED IN VITRO

Francesca Pacchierotti, Lucia Ardoino, Barbara Benassi, Claudia Consales, Eugenia Cordelli, Patrizia Eleuteri, Carmela Marino, Maurizio Sciortino, Martin H. Brinkworth, Guangdi Chen, James P. McNamee, Andrew William Wood, Carlijn R. Hooijmans, Rob B.M. de Vries. *Environment Intl.* Volume 157, December 2021, 106806.

Highlights

- Male infertility and adverse pregnancy outcomes are relevant human health problems.
- Radiofrequency electromagnetic fields are widespread in the human environment.
- A link between radiofrequency and adverse reproductive outcomes is controversial.
- This is the protocol of WHO-funded systematic review and meta-analysis on this issue.

Abstract

Background: Radiofrequency Electromagnetic Fields (RF-EMF) at environmental level have been reported to induce adverse effects on the male reproductive system and developing embryos. However, despite the number of experiments conducted since the 1970s, the diversity of testing approaches and exposure conditions, inconsistencies among results, and dosimetric flaws have not yet permitted a solid assessment of the relationship between RF-

EMF exposure and such effects, warranting a more systematic and methodologically rigorous approach to the evaluation of available data.

Objectives: This study aims at evaluating the effects of RF-EMF exposure on male fertility and pregnancy outcomes by a systematic review (SR) of experimental studies, conducted in compliance with international guidelines. The evidence will be organized into three streams:

- 1) Studies evaluating the impact of RF-EMF on the male reproductive system of experimental mammals;
- 2) studies evaluating the impact of RF-EMF on human sperm exposed in vitro;
- 3) studies evaluating the impact of RF-EMF on adverse pregnancy, birth outcomes and delayed effects in experimental mammals exposed in utero.

Study eligibility and criteria Eligible studies will include peer-reviewed articles reporting of original results about effects of controlled exposures to RF-EMF in the frequency range 100 kHz–300 GHz on the selected outcomes without any language or year-of-publication restrictions. Eligible studies will be retrieved by calibrated search strings applied to three electronic databases, PubMed, Scopus and EMF Portal and by manual search of the list of references of included papers and published reviews.

Study appraisal and synthesis method The internal validity of the studies will be evaluated using the Risk of Bias (RoB) Rating Tool developed by National Toxicology Program/Office of Health Assessment and Translation (NTP/OHAT) integrated with input from the SYRCLE RoB tool. Given sufficient commensurate data, meta- analyses will be performed, otherwise narrative syntheses will be produced. Finally, the certainty of the effects of RF-EMF exposure on male fertility and pregnancy and birth outcomes will be established following GRADE.

Funding The study is financially supported by the World Health Organization. Registration OSF Registration DOI <https://doi.org/10.17605/OSF.IO/7MUS3>; PROSPERO CRD42021227729, CRD42021227746.

<https://www.sciencedirect.com/science/article/pii/S0160412021004311>

THE EFFECTS OF RADIOFREQUENCY ELECTROMAGNETIC FIELDS EXPOSURE ON TINNITUS, MIGRAINE AND NON-SPECIFIC SYMPTOMS IN THE GENERAL AND WORKING POPULATION: A PROTOCOL FOR A SYSTEMATIC REVIEW ON HUMAN OBSERVATIONAL STUDIES

Martin Rööslī, Stefan Dongus, Hamed Jalilian, Maria Feychting, John Evers, Ekpereonne Esu, Chioma Moses Oringanje, Martin Meremikwu, Xavier Bosch-Capblanch. Environment International, Volume 157, 2021. doi: 10.1016/j.envint.2021.106852.

Highlights

- There is public concern to develop non-specific symptoms from EMF.
- No up to date comprehensive systematic review is available.
- Priority outcomes for head exposure are tinnitus, migraine, and headaches.
- Further priority outcomes are sleep disturbances and composite symptom scores.

Abstract

Background Applications emitting radiofrequency electromagnetic fields (RF-EMF; 100 kHz to 300 GHz) are widely used for communication (e.g. mobile phones), in medicine (diathermy) and in industry (RF heaters). Concern has been raised that RF-EMF exposure affects health related quality of life, because a part of the population reports to experience a variety of symptoms related to low exposure levels below regulatory limits.

Objectives To systematically review the effects of longer-term or repeated local and whole human body RF-EMF exposure on the occurrence of symptoms evaluating migraine, tinnitus, headaches, sleep disturbances and composite symptom scores as primary outcomes.

Methods We will follow the WHO handbook for guideline development. For the development of the systematic review protocol we considered handbook for conducting systematic reviews for health effects evaluations from the National Toxicology Program-Office of Health Assessment and Translation (NTP-OHAT) and COSTER (Recommendations for the conduct of systematic reviews in toxicology and environmental health research).

Eligibility criteria Peer-reviewed epidemiological studies in the general population or workers aiming to investigate the association between local or whole-body RF-EMF exposure for at least one week and symptoms are eligible for inclusion. Only cohort, case-control and panel studies will be included.

Information Sources We will search the scientific literature databases Medline, Web of Science, PsycInfo, Cochrane Library, Epistemonikos and Embase, using a predefined search strategy. This search will be supplemented by a search in the EMF-Portal and checks of reference lists of relevant papers and reviews.

Study Appraisal And Synthesis Method Data from included papers will be extracted according to predefined forms. Findings will be summarized in tables, graphical displays and in a narrative synthesis of the available evidence, complemented with meta-analyses. We will separately review effects of local, far field and occupational exposure.

Risk of bias The internal validity of included studies will be assessed using the NTP-OHAT Risk of Bias Rating Tool for Human and Animal Studies, elaborated to observational RF-EMF studies.

Evidence appraisal To rate certainty of the evidence, we will use the OHAT GRADE-based approach for epidemiological studies.

Framework And Funding This protocol concerns one of the ten different systematic reviews considered in a larger systematic review of the World Health Organization to assess potential health effects of exposure to RF- EMF in the general and working population.

Registration PROSPERO CRD42021239432.

Open access paper:

<https://www.sciencedirect.com/science/article/pii/S0160412021004773?via%3Dihub>

MORE RELIABLE BIOEFFECTS STUDIES AT “HIGH BAND” 5G FREQUENCIES

Kenneth R Foster, Vijayalaxmi. More Reliable Bioeffects Studies at “High Band” 5G Frequencies. *Frontiers in Communications and Networks*. 2:39. 2021. doi:10.3389/frcmn.2021.721925.

Abstract

One major source of controversy related to possible health effects of radiofrequency radiation (RFR) is the large number of reported statistically significant effects of exposure, over the entire RF part of the spectrum and over a wide range of exposure levels, even as health agencies do not find clear evidence for health hazards of exposure at levels within current IEEE and ICNIRP exposure limits. This Perspective considers 31 studies related to genetic damage produced by exposure to RFR at frequencies above 6 GHz, including at millimeter-wave (mm-wave) frequencies. Collectively, the papers report many statistically significant effects related to genetic damage, many at exposure levels below current exposure limits. However, application of five risk of bias (RoB) criteria and other considerations suggest that the studies in many cases are vulnerable to false discovery (nonreplicable results). The authors call for improvements in study design, analysis and reporting in future bioeffects research to provide more reliable information for health agencies and regulatory decision makers. This Perspective is a companion to another Perspective by Mattsson et al. elsewhere in this volume (Mattsson et al., 2021).

Conflict of interest

KF has received minor support for research on an unrelated topic (thermal dosimetry) by Microwave and Wireless Forum, an industry group.

The remaining author declares that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Open access paper: <https://www.frontiersin.org/article/10.3389/frcmn.2021.721925>

DOSIMETRY OF VARIOUS HUMAN BODIES EXPOSED TO MICROWAVE BROADBAND ELECTROMAGNETIC PULSES

Jerdvisanop Chakarothai, Wake Kanako, Fujii Katsumi. Dosimetry of Various Human Bodies Exposed to Microwave Broadband Electromagnetic Pulses. *Frontiers in Public Health*. 9:1182. 2021. doi:10.3389/fpubh.2021.725310.

Abstract

In this paper, human exposures to ultra-wideband (UWB) electromagnetic (EM) pulses in the microwave region are assessed using a frequency-dependent FDTD scheme previously proposed by the authors. Complex permittivity functions of all biological tissues used in the numerical analyses are accurately expressed by the four-term Cole–Cole model. In our method, we apply the fast inverse Laplace transform to determine the time-domain impulse response, utilize the Prony method to find the Z-domain representation, and extract residues and poles for use in the FDTD formulation. Update equations for the electric field are then derived via the Z-transformation. Firstly, we perform reflection and transmission analyses of

a multilayer composed of six different biological tissues and then confirm the validity of the proposed method by comparison with analytical results. Finally, numerical dosimetry of various human bodies exposed to EM pulses from the front in the microwave frequency range is performed, and the specific energy absorption is evaluated and compared with that prescribed in international guidelines.

Conclusion

We have performed numerical dosimetry on human bodies illuminated by an EM pulse from the front by using the (FD)2TD method, previously proposed by the authors. The method fully considers broadband characteristics of the complex relative permittivity of the biological media used in the analysis model via the application of the FILT and the Prony method. Firstly, we demonstrated the validity of the update coefficients, i.e., the residues and poles of the expression for the IIR in the z-domain, by comparing the numerical reflection coefficients with those derived from the EM theory. It was clarified that the numerical results within 2% of those obtained theoretically over a broad frequency range from 50 MHz to 10 GHz, demonstrating the validity of the proposed approach. It was also found that the transmission characteristics of the EM pulse into the CSF layer of a multilayer mimicking a human head are almost flat over a frequency range between 300 and 800 MHz and that the transmission decreases with increasing EM traveling distance from the skin boundary due to higher energy absorption at superficial biological tissues such as “Skin” and “Fat” when the frequency is higher than 1 GHz.

Therefore, most of the pulse energy that penetrates into the biological body has a frequency below 1 GHz. Then, numerical dosimetry of various human models exposed to an EM pulse having a frequency component of up to ~1.3 GHz was performed. The whole-body average SAR at 24 frequencies was determined by a single run of broadband FDTD simulations. The results matched those published in the literature, demonstrating the validity and availability of the proposed FDTD method. Then the SA distribution of each numerical model was determined, and it was found that the maximum SA occurs at the hands and neck for the ungrounded model, while they appear at the hands and ankle when the model is grounded. The maximum SA value was 0.290 pJ/kg for an incident electric field strength of 1 V/m or 0.437 nJ/kg for an incident power density of 2 W/m². It has been shown for the first time that, by using our proposed FDTD approach with the FILT and the Prony method, we can obtain quantitatively detailed information on SA that can be compared with the limits prescribed in international guidelines or standards.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.725310>

SIMULTANEOUS EXPOSURE TO ELECTROMAGNETIC FIELD FROM MOBILE PHONE AND UNIMPEDED FRUCTOSE DRINKING DURING PRE-, PERI-, AND POST-PUBERTAL STAGES PERTURBS THE HYPOTHALAMIC AND HEPATIC REGULATION OF ENERGY HOMEOSTASIS BY EARLY ADULthood: EXPERIMENTAL EVIDENCE

Ruchi Tripathi, Sanjay Kumar Banerjee, Jay Prakash Nirala, Rajani Mathur. Environ Sci Pollut Res Int. 2021 Sep 2. doi: 10.1007/s11356-021-15841-y.

Abstract

The present-day children-adolescents ubiquitously use the mobile phones and unrestrictedly consume fructose- laden diet. Unfortunately, a rise in the incidence of insulin resistance and fatty liver syndrome in young adults has also been recorded. To delineate a possible correlate, the effect of exposure to electromagnetic field (EMF) from the mobile phone and unrestricted fructose intake during pre-, peri-, and post-pubertal stages of development on orexigenic and anorexigenic signals arising from the hypothalamus and liver of rats is investigated here. The study design included four arms, i.e., "Normal", "Exposure Only (ExpO)", "Fructose Only (FruO)", and "Exposure with Fructose (EF)", wherein weaned rats received either "normal chow and drinking water" or "normal chow and fructose (15%) drinking solution" in presence and absence of EMF exposure (2 h/day) for 8 weeks. The results indicate that the total calories consumed by the EF were higher by early adulthood than normal, possibly under the influence of the raised levels of the orexigenic hormone, i.e., ghrelin, and it reflected as raised rate of weight gain. At early adulthood, the EF recorded mitigated response and sensitivity of insulin. Despite EF being a "fed-state", both centrally and peripherally, the glycolysis was restrained, but the gluconeogenesis was raised. Additionally, the altered lipid profile and the glycogen levels indicate that the EF developed fatty liver. The energy homeostasis of the EF was compromised as evidenced by

- (a) reduced expression of the glucosensors-GLUT2 and glucokinase in the hypothalamus and liver and
- (b) reduced expression of the cellular energy regulator-AMPK, orexigenic peptide-NPY, and anorexigenic peptide- POMC in the hypothalamus.

Taken together, the present study evidences that the exposure to EMF from the mobile phone and unrestricted fructose intake during childhood-adolescence impairs the central and peripheral pathways that mediate the glucosensing, glucoregulation, feeding, and satiety behavior by early adulthood.

<https://pubmed.ncbi.nlm.nih.gov/34476698/>

OXIDATIVE DAMAGE IN THE LIVER AND BRAIN OF THE RATS EXPOSED TO FREQUENCY-DEPENDENT RADIOFREQUENCY ELECTROMAGNETIC EXPOSURE: BIOCHEMICAL AND HISTOPATHOLOGICAL EVIDENCE

Anjali Sharma, Sadhana Shrivastava, Sangeeta Shukla. Free Radic Res. 2021 Aug 17;1-30. doi: 10.1080/10715762.2021.1966001.

Highlights

- EMR exposure showed frequency-dependent toxicity.
- Alterations in blood profile and modifications in the serological markers.
- Increasing lipid peroxidation indicating membrane damage.
- Inhibition of acetylcholinesterase activity affecting cholinergic neurotransmission.
- EMR exposure resulted in the loss of cellular energy and production of excess amounts of ROS thereby altering several antioxidant enzymes.
- Histopathological evidence of severe degenerative changes in the liver and brain.

Abstract

The study aimed to discover a link between the liver and brain's functional status due to frequency dependent- radiofrequency electromagnetic radiation (RF-EMR).

40 Wistar rats were randomly classified as control (sham-exposed) and EMR exposed groups. Animals were exposed to 900, 1800, and 2100 MHz with the specific absorption rate (SAR) 0.434 (W/Kg), 0.433 (W/Kg), and 0.453 (W/Kg) respectively. Animal exposure was limited at 1 hour/day, 5 days/week for 1 month with a restricted power density (900 MHz- 11.638 μ W/m², 1800- 11.438 μ W/m² and 2100 MHz frequency- 8.237 μ W/m²).

Exposure at various frequencies showed a frequency-dependent change in the body weight and hematologic parameters (RBCs, WBCs, platelets, hemoglobin, and hematocrit) as compared with the control group ($P \leq 0.01$)($P \leq 0.001$). A significant elevation in serum transaminases and bilirubin, urea, uric acid, and creatinine was noted, whereas albumin significantly decreased after EMR exposure ($P \leq 0.01$)($P \leq 0.001$). The blood glucose, lipid peroxidation, triglycerides, and cholesterol were elevated while adenosine triphosphatases, acetylcholinesterase, and tissue antioxidants such as glutathione, superoxide dismutase, catalase, glutathione reductase, glutathione Peroxidase, glutathione-S-transferase, and glucose-6-phosphate dehydrogenases were decreased significantly ($P \leq 0.001$). Histopathological observations of the liver showed centrilobular mononuclear cell infiltration and swelling in sinusoidal spaces, while in the brain degenerated pyramidal and Purkinje neurons were seen. Furthermore, Substantial evidence was found that the brain is more susceptible to oxidative mutilation compared to the liver of exposed animals.

In conclusion, RF-EMR exposure showed oxidative damage to the liver, increasing the incidence of brain damage in a frequency-dependent manner.

Highlights EMR exposure showed frequency-dependent toxicity. Alterations in blood profile and modifications in the serological markers. Increasing lipid peroxidation indicating membrane damage. Inhibition of acetylcholinesterase activity affecting cholinergic neurotransmission. EMR exposure resulted in the loss of cellular energy and production of

excess amounts of ROS thereby altering several antioxidant enzymes. Histopathological evidence of severe degenerative changes in the liver and brain.

<https://pubmed.ncbi.nlm.nih.gov/34404322/>

Conclusion

Our findings documented that the frequency of EMR exposure is a significant factor associated with its toxicity, the extent of toxicity is increased with a simultaneous increase in the frequency of exposure. In addition, this study confirms that EMR exposure initiates hepatic and neural injury, but the potency of damage was more prominent in the brain. Our results will provide support for future studies in the analysis of electromagnetic radiation and its associated systemic injury.

900 MHz Radiofrequency Field Induces Mitochondrial Unfolded Protein Response in Mouse Bone Marrow Stem Cells

Xie Wen, Xu Rui, Fan Caiyun, Yang Chunyu, Chen Haiyan, Cao Yi. 900 MHz Radiofrequency Field Induces Mitochondrial Unfolded Protein Response in Mouse Bone Marrow Stem Cells. *Frontiers in Public Health*. 9:1265. 2021. doi: 10.3389/fpubh.2021.724239

Abstract

Objective: To examine whether exposure of mouse bone marrow stromal cells (BMSC) to 900 MHz radiofrequency fields used in mobile communication devices can induce mitochondrial unfolded protein response (UPR^{mt}).

Methods: BMSCs were exposed to continuous wave 900 MHz radiofrequency fields (RF) at 120 $\mu\text{W}/\text{cm}^2$ power intensity for 4 h/d for 5 consecutive days. Cells in sham group (SH) were cultured in RF exposure system, but without RF radiation. The positive control cells were irradiated with 6 Gy X-ray at a dose rate of 1.103 Gy/min (XR). To inhibit the upstream molecular JNK2 of UPR^{mt}, cells in siRNA + RF, and siRNA + XR group were also pretreated with 100 nM siRNA-JNK2 for 48 h before RF/XR exposure. Thirty minutes, 4 h, and 24 h post-RF/XR exposure, cells were collected, the level of ROS was measured with flow cytometry, the expression levels of UPR^{mt}-related proteins were detected using western blot analysis.

Results: Compared with Sham group, the level of ROS in RF and XR group was significantly increased 30 min and 4 h post-RF/XR exposure ($P < 0.05$), however, the RF/XR-induced increase of ROS level reversed 24 h post-RF/XR exposure. Compared with Sham group, the expression levels of HSP10/HSP60/ClpP proteins in cells of RF and XR group increased significantly 30 min and 4 h post-RF/XR exposure ($P < 0.05$), however, the RF/XR-induced increase of HSP10/HSP60/ClpP protein levels reversed 24 h post-RF exposure. After interfering with siRNA-JNK2, the RF/XR exposures could not induce the increase of HSP10/HSP60/ClpP protein levels any more.

Conclusions: The exposure of 900 MHz RF at 120 $\mu\text{W}/\text{cm}^2$ power flux density could increase ROS level and activate a transient UPR^{mt} in BMSC cells. Mitochondrial homeostasis in term of protein folding ability is restored 24 h post-RF exposure.

Exposure to RF in our experimental condition did not cause permanent and severe mitochondrial dysfunctions. However, the detailed underlying molecular mechanism of RF-induced UPR^{mt} remains to be further studied.

Open access paper: <https://www.frontiersin.org/article/10.3389/fpubh.2021.724239>

EFFECTS OF MICROWAVE 10 GHZ RADIATION EXPOSURE IN THE SKIN OF RATS: AN INSIGHT ON MOLECULAR RESPONSES

Saurabh Verma, Gaurav K Keshri, Santanu Karmakar, Kumar Vyonkesh Mani, Satish Chauhan, Anju Yadav, Manish Sharma, Asheesh Gupta. Effects of Microwave 10 GHz Radiation Exposure in the Skin of Rats: An Insight on Molecular Responses. Radiat Res. 2021 Aug 18. doi: 10.1667/RADE-20-00155.1.

Abstract

Microwave (MW) radiation poses the risk of potential hazards on human health. The present study investigated the effects of MW 10 GHz exposure for 3 h/day for 30 days at power densities of 5.23 ± 0.25 and 10.01 ± 0.15 mW/cm² in the skin of rats. The animals exposed to 10 mW/cm² (corresponded to twice the ICNIRP-2020 occupational reference level of MW exposure for humans) exhibited significant biophysical, biochemical, molecular and histological alterations compared to sham-irradiated animals. Infrared thermography revealed an increase in average skin surface temperature by 1.8°C and standard deviation of 0.3°C after 30 days of 10 mW/cm² MW exposure compared to the sham-irradiated animals. MW exposure also led to oxidative stress (ROS, 4-HNE, LPO, AOPP), inflammatory responses (NFkB, iNOS/NOS2, COX-2) and metabolic alterations [hexokinase (HK), lactate dehydrogenase (LDH), citrate synthase (CS) and glucose-6-phosphate dehydrogenase (G6PD)] in 10 mW/cm² irradiated rat skin. A significant alteration in expression of markers associated with cell survival (Akt/PKB) and HSP27/p38MAPK-related stress-response signaling cascade was observed in 10 mW/cm² irradiated rat skin compared to sham-irradiated rat skin. However, MW-irradiated groups did not show apoptosis, evident by unchanged caspase-3 levels. Histopathological analysis revealed a mild cytoarchitectural alteration in epidermal layer and slight aggregation of leukocytes in 10 mW/cm² irradiated rat skin. Altogether, the present findings demonstrated that 10 GHz exposure in continuous-wave mode at 10 mW/cm² (3 h/day, 30 days) led to significant alterations in molecular markers associated with adaptive stress-response in rat skin.

Furthermore, systematic scientific studies on more prevalent pulsed-mode of MW-radiation exposure for prolonged duration are warranted.

<https://pubmed.ncbi.nlm.nih.gov/34407201/>

High Power Electromagnetic Waves Exposure of Healthy and Tumor Bearing Mice: Assessment of Effects on Mice Growth, Behavior, Tumor Growth, and Vessel Permeabilization

Kolosnjaj-Tabi J, Golzio M, Bellard E, Catrain A, Chretiennot T, Saurin Q, Tarayre J, Vezinet R, Rols MP. High Power Electromagnetic Waves Exposure of Healthy and Tumor Bearing Mice: Assessment of Effects on Mice Growth, Behavior, Tumor Growth, and Vessel Permeabilization. *Int J Mol Sci.* 2021 Aug 7;22(16):8516. doi: 10.3390/ijms22168516.

Abstract

High power radiofrequencies may transiently or permanently disrupt the functioning of electronic devices, but their effect on living systems remains unknown. With the aim to evaluate the safety and biological effects of narrow-band and wide-band high-power electromagnetic (HPEM) waves, we studied their effects upon exposure of healthy and tumor-bearing mice. In field experiments, the exposure to 1.5 GHz narrow-band electromagnetic fields with the incident amplitude peak value level in the range of 40 kV/m and 150 MHz wide-band electric fields with the amplitude peak value in the range of 200 kV/m, did not alter healthy and tumor-bearing animals' growth, nor it had any impact on cutaneous murine tumors' growth. While we did not observe any noticeable behavioral changes in mice during the exposure to narrow-band signals when wide-band HPEM signals were applied, mice could behave in a similar way as they respond to loud noise signals: namely, if a mouse was exploring the cage prior to signal application, it returned to companion mates when wide-band HPEM signals were applied. Moreover, the effect of wide-band signals was assessed on normal blood vessels permeability in real-time in dorsal-chamber-bearing mice exposed in a pilot study using wide-band signal applicators. Our pilot study conducted within the applicator and performed at the laboratory scale suggests that the exposure to wide-band signals with the amplitude of 47.5 kV/m does not result in increased vessel permeability.

<https://pubmed.ncbi.nlm.nih.gov/34445226/>

HEALTH EFFECTS OF WIFI RADIATION: A REVIEW BASED ON SYSTEMATIC QUALITY EVALUATION

Stefan Dongus, Hamed Jalilian, David Schürmann & Martin Rösli (2021) Health effects of WiFi radiation: a review based on systematic quality evaluation, *Critical Reviews in Environmental Science and Technology*, DOI: 10.1080/10643389.2021.1951549

Abstract

Although WiFi contributes little to total radiofrequency electromagnetic field (RF-EMF) exposure in our everyday environment, concern has raised whether this specific type of modulated RF-EMF causes health problems. The aim of this review is to evaluate all types of studies that investigated biological and health effects of WiFi exposure and fulfilled basic quality criteria. Eligible for inclusion were epidemiological, human experimental, in vivo and in vitro studies using realistic WiFi exposure settings. We conducted a systematic literature search for all papers published between January 1997 and August 2020 followed by a quality review addressing blinding and dosimetry in experimental studies and various types of biases in epidemiological studies. All studies fulfilling the quality criteria were descriptively summarized in terms of observation or absence of associations. From 1385 articles identified by the literature search, 23 fulfilled basic quality criteria: 6 epidemiological papers, 6 human experimental articles, 9 in vivo articles, and 2 in vitro articles. Whereas in vivo and in vitro

studies applied exposure levels up to 4 W/kg, human studies dealt with exposure levels several orders of magnitude below the ICNIRP guidelines, which are typical for WiFi exposure situations in the everyday environment. Numerous outcomes ranging from biological markers to symptoms were mostly found not to be associated with WiFi exposure. Sporadic findings were not consistent in terms of outcomes or exposure-response associations. This review based on a systematic literature search and quality evaluation does not suggest detrimental health effects from WiFi exposure below regulatory limits.

<https://doi.org/10.1080/10643389.2021.1951549>

‘PROOF OF EHS BEYOND ALL REASONABLE DOUBT’

Michael Bevington. ‘Proof of EHS beyond all reasonable doubt’. Comment on: Leszczynski D. Review of the scientific evidence on the individual sensitivity to electromagnetic fields (EHS). Rev Environ Health 2021; doi: 10.1515/reveh-2021-0038. Reviews on Environmental Health, vol. , no. , 2021, pp. 000010151520210101. <https://doi.org/10.1515/reveh-2021-0101>

No abstract

Open access letter: <https://www.degruyter.com/document/doi/10.1515/reveh-2021-0101/html>

POOLED ANALYSIS OF RECENT STUDIES OF MAGNETIC FIELDS AND CHILDHOOD LEUKEMIA

Aryana T. Amoon, John Swanson, Corrado Magnani, Christoffer Johansen, Leeka Kheifets. Pooled analysis of recent studies of magnetic fields and childhood leukemia. Environmental Research, 2021, 111993, doi: 10.1016/j.envres.2021.111993.

Highlights

- No association in recent studies between magnetic fields and childhood leukemia.
- Current results are not in line with previous pooled analyses.
- A small elevated risk for the aggregate evidence remains.

Abstract

Background Over forty epidemiologic studies have addressed an association between measured or calculated extremely-low-frequency magnetic fields (MF) and childhood leukemia. These studies have been aggregated in a series of pooled analyses, but it has been 10 years since the last such.

Methods We present a pooled analysis combining individual-level data (24,994 cases, 30,769 controls) from four recent studies on MF and childhood leukemia.

Results Unlike previous pooled analyses, we found no increased risk of leukemia among children exposed to greater MF: odds ratio (OR) = 1.01, for exposure $\geq 0.4 \mu\text{T}$ (μT) compared with exposures $< 0.1 \mu\text{T}$. Similarly, no association was observed in the subset of acute lymphoblastic leukemia, birth homes, studies using calculated fields, or when geocoding accuracy was ignored. In these studies, there is a decline in risk over time, also evident when we compare three pooled analyses. A meta-analysis of the three pooled analyses overall presents an OR of 1.45 (95% CI: 0.95–2.20) for exposures $\geq 0.4 \mu\text{T}$.

Conclusions Our results are not in line with previous pooled analysis and show a decrease in effect to no association between MF and childhood leukemia. This could be due to methodological issues, random chance, or a true finding of disappearing effect.
<https://pubmed.ncbi.nlm.nih.gov/34481821/>

HUMAN DETECTION THRESHOLDS OF DC, AC, AND HYBRID ELECTRIC FIELDS: A DOUBLE-BLIND STUDY

Michael Kursawe, Dominik Stunder, Thomas Krampert, Andrea Kaifie, Sarah Drießen, Thomas Kraus, Kathrin Jankowiak. Human detection thresholds of DC, AC, and hybrid electric fields: a double-blind study. *Environ Health*. 2021 Aug 21;20(1):92. doi: 10.1186/s12940-021-00781-4.

Abstract

Background: In the course of the ongoing transition of electric energy systems, transmission corridors are often upgraded to higher voltages and other technologies leading to another quality of human exposure. The study aims to determine human detection thresholds for direct current (DC), alternating current (AC), and hybrid electric fields (various DC; constant AC).

Methods: A total of 203 participants were exposed to DC, AC, and hybrid electric fields (EFs) in a highly specialized whole-body exposure laboratory using a double-blind experimental setting. Additionally, the participants were exposed to ion currents in part of the DC and hybrid sessions. To investigate environmental influences, relative humidity was changed in two subgroups during EF perception. Methods derived from the signal detection theory and the adaptive staircase procedure based on the single interval adjustment matrix were used to assess individual sensitivity and detection thresholds, respectively.

Results: The results indicated that detection thresholds of hybrid EF were lower compared to single EF presentation of DC or AC. Ion current exposure enhanced EF perception. High relative humidity facilitated DC EF perception, whereas low relative humidity reinforced the perception of AC EFs.

Conclusions: With this systematic investigation of human perception of DC, AC, and hybrid EFs, detection thresholds were provided, which can help improve the construction processes of energy transmission systems and the prevention of unwanted sensory perception by contributing to the determination of limit values.

Conclusion

This study is the first systematic investigation of hybrid EF perception in humans using a double-blind experimental setup. While detection thresholds of DC and AC EF were lower compared to previous studies, the synergistic effect of both field types on human perception was shown. Especially, in the minimal hybrid EF combination (2 kV/m DC and 4 kV/m AC), 40% of the participants were able to successfully detect the EF. In terms of unwanted sensory perception, this should be considered in the discussion on reference levels and recommendations for hybrid EFs, which current remain non-existent. Furthermore, we suggest a further investigation on low hybrid EFs in detail to obtain a fine-grained picture of the lower bound of human EF perception. Although the average detection thresholds do not

undercut the existing reference levels for DC and AC EFs, the study found evidence for successful EF perception around these reference levels in a small subset of participants. As the reference levels were mainly designed to prevent adverse health effects during EF exposure [14, 16], the current results do not question these values. Furthermore, the findings can be utilized in the discussion on whether sensory perception, as a byproduct of EF exposure, is acceptable.

We enhanced the external validity of the results by investigating subgroups across different levels of relative humidity. Hints for an enhanced perception of DC EF in a high-humidity environment and a facilitated AC perception under low-humidity conditions were confirmed. These results may further contribute to the understanding of the underlying mechanisms and should be considered as well in the discussion on recommendations for EF exposure. Thus, this study provided reliable EF detection thresholds that add quantitative information on sensory perception to the discourse on the impact of EF exposure. In this manner, it can help to improve the construction processes of planned HVDC and hybrid overhead power lines.

Open access paper: <https://ehjournal.biomedcentral.com/articles/10.1186/s12940-021-00781-4>

EFFECTS OF WI-FI RADIOFREQUENCY RADIATION ON CARBAPENEM-RESISTANT KLEBSIELLA PNEUMONIAE

Ilham Said-Salman, Wissam Yassine, Ali Rammal, Mohammad Hneino, Hoda Yusef, Mohamed Moustafa. Effects of Wi-Fi Radiofrequency Radiation on Carbapenem-Resistant Klebsiella pneumoniae. Bioelectromagnetics. 2021 Aug 1. doi: 10.1002/bem.22364.

Abstract

The hazardous consequences of electromagnetic field (EMF) exposure represent a public health concern. Common sources of EMF include smartphones and wireless fidelity (Wi-Fi). The aim of our study is to assess whether exposure to Wi-Fi radiofrequency radiation influences the pathogenic traits of carbapenem-resistant Klebsiella pneumoniae. The susceptibility to antibiotics was evaluated by the determination of minimum inhibitory concentrations (MIC). In this study, K. pneumoniae showed a non-linear response to treatments with Colistin and Gentamycin following different Wi-Fi exposure periods. Transmission electron microscopy revealed morphological changes in the bacterial cell membrane within 24 h of Wi-Fi exposure. Crystal violet quantification and quantitative real-time polymerase chain reaction showed that the ability to form biofilms was greater in Wi-Fi exposed K. pneumoniae when compared to control. Moreover, higher levels of bcsA, mrkA, and luxS messenger RNAs were observed. Our data suggest that Wi-Fi exposure can influence bacteria in a stressful way, leading to an alteration in their antibiotic susceptibility, morphological changes, and cumulative biofilm formation.

<https://pubmed.ncbi.nlm.nih.gov/34337771/>

Excerpts

"Several opportunistic bacterial pathogens with multidrug resistance were identified in the area of telecommunication stations [Adebayo et al., 2015]. Research in this area seems to be of big interest and mandatory in order to improve the measures of protection assessment against the occurrence of antibiotic-resistant organisms in the environment."

"We measured a constant field of 6 V/m at the position of the exposed sample corresponding to a power flux density of 0.05 W/m². The influence of the exposure to Wi-Fi waves was investigated, and the protocol consisted of continuous radiation exposures for different periods (1, 5, 10, 20, 24, and 48 h). A control measurement was done for each experimental condition in a Faraday bag at an identical temperature to avoid any external radiation."

"K. pneumonia is a rod-shaped Gram-negative opportunistic pathogenic bacterium causing nosocomial infections. K. pneumonia has emerged in recent years with enhanced virulence. K. pneumoniae carbapenemase (KPC) is the main enzymatic resistance mechanism used by K. pneumoniae against antibiotics [Wang et al., 2011]. Carbapenem-resistant K. pneumoniae carries the blaKPC gene which confers resistance to carbapenem antibiotics. Bacterial virulence factors such as antimicrobial resistance (AMR) and biofilm formation play key roles in infection settlement. The World Health Organization (WHO) stated that AMR is a "global health crisis" [WHO, 2018]."

"In summary, exposure of K. pneumoniae to 2.4 GHz Wi-Fi radiofrequency radiation affected the pathogenic traits of multidrug-resistant bacteria. There was a nonlinear response of K. pneumoniae to Colistin and Gentamycin antibiotics after Wi-Fi pre-exposure. The 24 h exposure period to Wi-Fi radiation disrupted the cell membrane and disintegrated its protoplasm. In addition, the transcription of representative genes implicated in biofilm formation and quorum sensing in K. pneumoniae was upregulated due to Wi-Fi exposure.

The different bacterial responses to EMF observed among reported studies may result from the differences found in cell wall composition between Gram-positive and Gram-negative bacteria as well as the duration and type of the radiation. Low and continuous far-field exposure such as Wi-Fi may cause unexpected antibiotic resistance in the future. Under stress, bacteria tend to regulate their membrane properties through homeostasis by altering the lipid composition, thus resulting in higher permeability [Zhang and Rock, 2008]. While there might be a modification in membrane permeability by such low fields, allowing increased susceptibility to antibiotics, it is not, however, sufficient to induce cell death. Thus, bacteria are more easily affected by lower doses of antibiotics in the environment. Further molecular experiments at gene and protein levels are required for understanding more details on the mechanisms by which these biological changes are taking place in bacteria following exposure to Wi-Fi radiation. Considering the impact of EMF on bacteria is an essential approach for controlling public health threats and may be beneficial for medical applications."

MEDICINAL PLANTS IN MITIGATING ELECTROMAGNETIC RADIATION-INDUCED NEURONAL DAMAGE: A CONCISE REVIEW

Raghu SV, Kudva AK, Rajanikant GK, Baliga MS. Medicinal plants in mitigating electromagnetic radiation-induced neuronal damage: a concise review. Electromagn Biol Med. 2021 Aug 12:1-14. doi: 10.1080/15368378.2021.1963762.

Abstract

Although the evidence is inconclusive, epidemiological studies strongly suggest that increased exposure to electromagnetic radiation (EMR) increases the risk of brain tumors, parotid gland tumors, and seminoma. The International Agency for Research on Cancer (IARC) has classified mobile phone radiofrequency radiation as possibly carcinogenic to humans (Group 2B).

Humans being are inadvertently being exposed to EMR as its prevalence increases, mainly through mobile phones. Radiation exposure is unavoidable in the current context, with mobile phones being an inevitable necessity. Prudent usage of medicinal plants with a long history of mention in traditional and folklore medicine and, more importantly, are safe, inexpensive, and easily acceptable for long-term human use would be an appealing and viable option for mitigating the deleterious effects of EMR. Plants with free radical scavenging, anti-oxidant and immunomodulatory properties are beneficial in maintaining salubrious health. Green tea polyphenols, Ginkgo biloba, lotus seedpod procyanidins, garlic extract, Loranthus longiflorus, Curcuma amada, and Rosmarinus officinalis have all been shown to confer neuroprotective effects in validated experimental models of study. The purpose of this review is to compile for the first time the protective effects of these plants against mobile phone-induced neuronal damage, as well as to highlight the various mechanisms of action that are elicited to invoke the beneficial effects.

<https://pubmed.ncbi.nlm.nih.gov/34382485/>

Conclusions

Recent research suggests green tea polyphenols, Ginkgo biloba, lotus seedpod procyanidins, garlic extract, Loranthus longiflorus, Curcuma amada, and Rosmarinus officinalis may be beneficial in preventing EMR-induced brain damage. The brain-protective effects of these plants are thought to be mediated by a variety of mechanisms, including scavenging of radiation-induced free radicals, increased cellular anti-oxidant status, inhibition of neuronal apoptosis, and improved neurobehavioral outcomes. Given the increasing prevalence of intentional and unintentional EMR exposure in daily life, more comprehensive cell culture, animal, and clinical studies using pharmacologically validated models that incorporate multiple biochemical, physiological, pathological, and neurobehavioral end points are required. Such studies will be highly beneficial because these agents have been validated for various health benefits and are known to improve/maintain general health; if they are also protective against EMR-induced damage in multiple studies, they will be extremely useful to both the scientific fraternity and society at large. The most important factors to consider when starting clinical studies are that these medicinal plants have been used since antiquity, are non-toxic, have a wide range of medicinal properties, and are easily acceptable for human use. Green tea polyphenols could be particularly useful because it is one of the most widely consumed beverages, has a wide range of medicinal applications, and is a relatively non-toxic agent with widespread acceptance. According to data from various clinical trial registries, green tea is being studied at various stages (phases 1 to 4) in humans for various ailments. A well-designed study aimed at collecting secondary endpoints with a checklist for daily mobile phone use in volunteers enrolled in clinical trials will be a cost- and time-effective way to validate the neuroprotective effects of medicinal plants against EMR-induced damage.

MICROARRAY PROFILING OF LNCRNA EXPRESSION IN THE TESTIS OF PUBERTAL MICE FOLLOWING MORNING AND EVENING EXPOSURE TO 1800 MHZ RADIOFREQUENCY FIELDS

Qin F, Cao H, Feng C, Zhu T, Zhu B, Zhang J, Tong J, Pei H. Microarray profiling of LncRNA expression in the testis of pubertal mice following morning and evening exposure to 1800 MHz radiofrequency fields. *Chronobiol Int.* 2021Aug8; 1-16.doi:10.1080/07420528.2021.1962902.

Abstract

In this paper, the chronotoxicity of radiofrequency fields (RF) in the pubertal testis development and the involved molecular pathways were investigated by exposing four-week-old mice to RF (1800 MHz, SAR, 0.50 W/kg) in the morning and evening of each day for three weeks. Then, pathological changes and functional indices within the testis were determined. We also used a long non-coding RNA (lncRNA) microarray and GO/KEGG pathway analyses to determine lncRNA expression profiles and predict their potential functions. The cis and trans regulation of lncRNAs were investigated, and an interaction network was constructed using Cytoscape software. RF exposure led to a range of pathological changes in the testes of adolescent mice, as testicular weights and daily sperm productions decreased, and the testosterone secretion reduced.

Furthermore, RF induced dysregulation in the expression of testicular lncRNAs. We identified 615 and 183 differentially expressed lncRNAs that were associated with morning and evening exposure to RF, respectively. From 15 differential expression lncRNAs both in morning RF group and evening RF group, we selected 6 lncRNAs to be validated by quantitative reverse transcription PCR (qRT-PCR). The differentially expressed lncRNAs induced by morning RF exposure were highly correlated with many different pathways, including Fanconi syndrome, metabolic processes, cell cycle, DNA damage, and DNA replication. Trans-regulation analyses further showed that differentially expressed lncRNAs were involved in multiple transcription factor-regulated pathways, such as TCFAP4, NFkB, HINFP, TFDP2, FoxN1, and PAX5. These transcription factors have all been shown to be involved in the modulation of testis development, cell cycle progression, and spermatogenesis. These findings suggest that the extent to which 1800 MHz RF induced toxicity in the testes and changed the expression of lncRNAs showed differences between morning exposure and evening exposure. These data indicate that differentially expressed lncRNAs play crucial roles in the RF exposure damage to the developing pubertal testis. Collectively, our findings provide a better understanding of the mechanisms underlying the toxic effects of RF exposure on testicular development.

<https://pubmed.ncbi.nlm.nih.gov/34369206/>

PROTEOMIC AND BIOLOGICAL ANALYSES REVEAL THE EFFECT ON GROWTH UNDER FLOODING STRESS OF CHICKPEA IRRADIATED WITH MILLIMETER WAVES

Setsuko Komatsu, Junya Maruyama, Takashi Furuya, Xiaojian Yin, Hisateru Yamaguchi, Keisuke Hitachi, Natsuki Miyashita, Kunihiro Tsuchida, Masahiko Tani. Proteomic and Biological Analyses Reveal the Effect on Growth under Flooding Stress of Chickpea Irradiated with Millimeter Waves. *J Proteome Res.* 2021 Aug 30. doi: 10.1021/acs.jproteome.1c00368.

Abstract

Chickpea cultivated on marginal lands in arid and semiarid tropics is one of the food legumes, and its growth is reduced by flooding stress. Millimeter-wave irradiation has influences on organisms, and it improves the growth of plants such as soybean. To reveal the dynamic effects of millimeter-wave irradiation on chickpea under flooding, gel- and label-free proteomic analysis was conducted. Millimeter-wave irradiation improved chickpea growth and its tolerance to flooding stress. According to functional categorization, oppositely changed proteins were correlated with photosynthesis, fermentation, and protein degradation. Immunoblot analysis confirmed that RuBisCO activase and large subunits decreased in leaves under flooding; however, they are recovered in irradiated chickpea even if it was in this condition. The activity and accumulation of alcohol dehydrogenase increased in roots under flooding; however, this followed the same pattern. Cell death was significantly increased and decreased by flooding on unirradiated and irradiated chickpeas, respectively. These findings suggest that irradiation with millimeter waves on chickpea seeds improves the recovery of plant growth through regulation of photosynthesis in leaves and fermentation in roots. Furthermore, millimeter-wave irradiation might promote chickpea tolerance under flooding via the regulation of cell death.

<https://pubmed.ncbi.nlm.nih.gov/34455783/>

TRANSCRIPTOMIC PROFILE REVEALS DEREGULATION OF HEARING-LOSS RELATED GENES IN VESTIBULAR SCHWANNOMA CELLS FOLLOWING ELECTROMAGNETIC FIELD EXPOSURE (50 HZ)

Colciago A, Audano M, Bonalume V, Melfi V, Mohamed T, Reid AJ, Faroni A, Greer PA, Mitro N, Magnaghi V. Transcriptomic Profile Reveals Deregulation of Hearing-Loss Related Genes in Vestibular Schwannoma Cells Following Electromagnetic Field Exposure. *Cells.* 2021 Jul 20;10(7):1840. doi: 10.3390/cells10071840.

Abstract

Hearing loss (HL) is the most common sensory disorder in the world population. One common cause of HL is the presence of vestibular schwannoma (VS), a benign tumor of the VIII cranial nerve, arising from Schwann cell (SC) transformation. In the last decade, the increasing incidence of VS has been correlated to electromagnetic field (EMF) exposure, which might be considered a pathogenic cause of VS development and HL. Here, we explore the molecular mechanisms underlying the biologic changes of human SCs and/or their oncogenic transformation following EMF exposure. Through NGS technology and RNA-Seq transcriptomic analysis, we investigated the genomic profile and the differential display of HL-related genes after chronic EMF. We found that chronic EMF exposure modified the cell proliferation, in parallel with intracellular signaling and metabolic pathways changes, mostly related to translation and mitochondrial activities. Importantly, the expression of HL-related genes such as NEFL, TPRN, OTOGL, GJB2, and REST appeared to be deregulated in chronic

EMF exposure. In conclusion, we suggest that, at a preclinical stage, EMF exposure might promote the transformation of VS cells and contribute to HL.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8307028/>

EXPOSURE TO 60 HZ MAGNETIC FIELD CAN AFFECT MEMBRANE PROTEINS AND MEMBRANE POTENTIAL IN HUMAN CANCER CELLS

Seiya Hayashi, Makiko Kakikawa. Exposure to 60 Hz magnetic field can affect membrane proteins and membrane potential in human cancer cells. *Electromagn Biol Med*. 2021 Aug 15;1-8. doi: 10.1080/15368378.2021.1958340.

Abstract

The experimental data support the hypothesis that extremely low frequency magnetic field (ELF-MF) can affect cell membranes. Since our previous studies suggested that MF changes the permeability of cell membrane, in this study we focused on the cell membrane and investigated the effect of 60 Hz, 50 mT MF on the membrane potential and membrane proteins. The membrane potentials of three cultured human cancer cell lines, A549, MES-SA, and MES-SA/Dx5, were increased by exposure to ELF-MF. When exposed to MF and an anticancer drug, changes in the membrane potentials were detected in A549 and MES-SA cells, but not in the multi drug-resistant cells, MES-SA/Dx5. We examined whether MF has an influence on the membrane proteins extracted from cultured A549 cells, using DiBAC4(3) dye enhanced fluorescence binding to a hydrophobic site. The increase in fluorescence observed following MF exposure for 10 min indicated that the structure of the hydrophobic site on the membrane proteins changed and became more likely to bind the probe dye. A decrease in fluorescence was detected following exposure to MF for 240 min. These results indicated that 60 Hz, 50 mT MF causes changes in the membrane potential of cultured cancer cells and the conformation of membrane proteins extracted from cultured cancer cells, and has different effects depending on the exposure time. <https://pubmed.ncbi.nlm.nih.gov/34396886/>

SELECTIVE INDUCTION OF RAPID CYTOTOXIC EFFECT IN GLIOBLASTOMA CELLS BY OSCILLATING MAGNETIC FIELDS

Helekar SA, Hambarde S, Ijare OB, Pichumani K, Baskin DS, Sharpe MA. Selective induction of rapid cytotoxic effect in glioblastoma cells by oscillating magnetic fields. *J Cancer Res Clin Oncol*. 2021 Sep 3. doi: 10.1007/s00432-021-03787-0.

Abstract

Purpose: The mechanisms underlying anticancer effects of electromagnetic fields are poorly understood. An alternating electric field-generating therapeutic device called Optune™ device has been approved for the treatment of glioblastoma (GBM). We have developed a new device that generates oscillating magnetic fields (OMF) by rapid rotation of strong permanent magnets in specially designed patterns of frequency and timing and have used it to treat an end-stage recurrent GBM patient under an expanded access/compassionate use treatment protocol. Here, we ask whether OMF causes selective cytotoxic effects in GBM and whether it is through generation of reactive oxygen species (ROS).

Methods: We stimulated patient derived GBM cells, lung cancer cells, normal human cortical neurons, astrocytes, and bronchial epithelial cells using OMF generators (oncoscillators) of our Oncomagnetic Device and compared the results to those obtained under unstimulated or

sham-stimulated control conditions. Quantitative fluorescence microscopy was used to assess cell morphology, viability, and ROS production mechanisms.

Results: We find that OMF induces highly selective cell death of patient derived GBM cells associated with activation of caspase 3, while leaving normal tissue cells undamaged. The cytotoxic effect of OMF is also seen in pulmonary cancer cells. The underlying mechanism is a marked increase in ROS in the mitochondria, possibly in part through perturbation of the electron flow in the respiratory chain.

Conclusion: Rotating magnetic fields produced by a new noninvasive device selectively kill cultured human glioblastoma and non-small cell lung cancer cells by raising intracellular reactive oxygen species, but not normal human tissue cells.
<https://pubmed.ncbi.nlm.nih.gov/34477946/>

COMBINED EFFECT OF CO-EXPOSURE TO DI (2-ETHYLHEXYL) PHTHALATES AND 50-HZ MAGNETIC-FIELDS ON PROMOTING HUMAN AMNIOTIC CELLS PROLIFERATION

Liangjing Chen, Anfang Ye, Xiaochen Liu, Jingchun Lu, Qixin Xie, Yixin Guo, Wenjun Sun. *Ecotoxicol Environ Saf.* 2021 Aug 26;224:112704. doi: 10.1016/j.ecoenv.2021.112704.

Abstract

Di (2-ethylhexyl) phthalate (DEHP) and extremely low-frequency electromagnetic fields (ELF-EMFs) exist far and wide in our surroundings. Studies have reported that both of DEHP and ELF-EMFs could promote cell proliferation which is related with adverse bioeffects. In this study, we investigated whether there is the combined effect between DEHP and 50-Hz magnetic fields (MFs) on cell proliferation in human amniotic (FL) cells. Results revealed that the low-concentration DEHP (1 μ M) could promote FL cell proliferation, whereas the high-dose DEHP (100 μ M) inhibited cell proliferation. When FL cells were treated jointly by a 50-Hz, 0.2-mT MF and 0.1 μ M DEHP, the proliferation rate of cells was significantly higher than that of single factor exposure.

Additionally, co-exposure to under-threshold MF and DEHP could cooperatively activate protein kinase B (Akt), sphingosine kinase 1 (SphK1) and extracellular signal regulated kinase (ERK) in a cascade manner, and finally mediate cell proliferation. Taken together, the findings of this study indicated that the co-exposure to under- threshold MF and DEHP could jointly promote cell proliferation through activating proliferation-related signal pathway, which warned us that it should be cautious about assessing the underlying health hazards of co-exposure to MFs and DEHP at under-threshold levels.

Open access paper: <https://pubmed.ncbi.nlm.nih.gov/34455183/>

EVALUATION OF PULSED ELECTROMAGNETIC FIELD EFFECTS: A SYSTEMATIC REVIEW AND META-ANALYSIS ON HIGHLIGHTS OF TWO DECADES OF RESEARCH IN VITRO STUDIES

Mahsa Mansourian, Ahmad Shanei. *Int.* 2021 Jul 29;2021:6647497. doi: 10.1155/2021/6647497.

Abstract

Pulsed electromagnetic field (PEMF) therapy is a type of physical stimulation that affects biological systems by producing interfering or coherent fields. Given that cell types are

significantly distinct, which represents an important factor in stimulation, and that PEMFs can have different effects in terms of frequency and intensity, time of exposure, and waveform. This study is aimed at investigating if distinct positive and negative responses would correspond to specific characteristics of cells, frequency and flux density, time of exposure, and waveform. Necessary data were abstracted from the experimental observations of cell-based in vitro models. The observations were obtained from 92 publications between the years 1999 and 2019, which are available on PubMed and Web of Science databases. From each of the included studies, type of cells, pulse frequency of exposure, exposure flux density, and assayed cell responses were extracted. According to the obtained data, most of the experiments were carried out on human cells, and out of 2421 human cell experiments, cell changes were observed only in 51.05% of the data. In addition, the results pointed out the potential effects of PEMFs on some human cell types such as MG-63 human osteosarcoma cells (p value < 0.001) and bone marrow mesenchymal stem cells. However, human osteogenic sarcoma SaOS-2 ($p < 0.001$) and human adipose-derived mesenchymal stem cells (AD-MSCs) showed less sensitivity to PEMFs. Nevertheless, the evidence suggests that frequencies higher than 100 Hz, flux densities between 1 and 10 mT, and chronic exposure more than 10 days would be more effective in establishing a cellular response. This study successfully reported useful information about the role of cell type and signal characteristic parameters, which were of high importance for targeted therapies using PEMFs. Our findings would provide a deeper understanding about the effect of PEMFs in vitro, which could be useful as a reference for many in vivo experiments or preclinical trials.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8342182/>

GROWTH ALTERATION OF ALLIUM CEPA L. ROOTS EXPOSED TO 1.5 MT, 25 HZ PULSED MAGNETIC FIELD

Alejandro González-Vidal, Silvia Mercado-Sáenz, Antonio M Burgos-Molina, Francisco Sendra-Portero, Miguel J Ruiz-Gómez. *Int J Environ Health Res.* 2021 Sep 2;1-13. doi: 10.1080/09603123.2021.1972090.

Abstract

The response of plants to magnetic fields (MF) is not fully understood. This work studies the effects of pulsed MF on the germination and growth of *Allium cepa* roots. Onions were exposed to 25Hz, 1.5mT, 33h. Pulsed MF was generated by a Helmholtz-type equipment that generated rectangular voltage pulses. The results showed that fewer roots grew in the specimens exposed to pulsed MF (14 ± 6 roots on day 1 to 21 ± 8 on day 4) than in the control groups (32 ± 17 to 48 ± 23) ($p < 0.05$ Friedman). Control specimens showed a root mean length of 7 ± 4 mm (day 1) and 24 ± 10 mm (day 4). The specimens treated with pulsed MF showed a length of 4 ± 2 mm (day 1), reaching 18 ± 9 mm on day 4 ($p < 0.001$ ANOVA). In conclusion, the exposure of *Allium cepa* specimens to 25Hz, 1.5mT pulsed MF during 33h produces a decrease in the germination and growth of roots.

<https://pubmed.ncbi.nlm.nih.gov/34474627/>

CAN STATIC ELECTRIC FIELDS INCREASE THE ACTIVITY OF NITRIC OXIDE SYNTHASE AND INDUCE OXIDATIVE STRESS AND DAMAGE OF SPLEEN?

Li Dong, Junli Xiang, Jingyi Guo, Guangdi Chen, Guoqing Di. Can static electric fields increase the activity of nitric oxide synthase and induce oxidative stress and damage of spleen? *Environ Sci Pollut Res Int.* 2021 Aug 16. doi: 10.1007/s11356-021-15853-8.

Abstract

With the rapid development of ultra-high-voltage (UHV) direct-current (DC) transmissions, the impact of static electric fields (SEF) in the vicinity of overhead UHV DC transmission lines on health has aroused much public concern. This study explored the effects of 56.3kV/m SEF on the spleen of mice. Results showed that SEF exposure of 21days significantly increased malonic dialdehyde content, superoxide dismutase activity, calcineurin activity, nitric oxide synthase (NOS) activity, and the mRNA expression levels of tumor necrosis factor- α (TNF- α) and nuclear factor- κ B (NF- κ B) in the spleen and caused the separation of nucleus and nuclear membrane, the disappearance of mitochondrial membrane, and the deficiency of mitochondrial cristae in splenic lymphocytes. By analysis and discussion, it was deduced that SEF could induce oxidative stress of the spleen by increasing the activity of NOS. Oxidative stress could further cause ultrastructural changes of splenic lymphocytes. Moreover, oxidative stress could cause the increase of the mRNA expression levels of TNF- α and NF- κ B, which contributed to the occurrence of spleen inflammation.

<https://pubmed.ncbi.nlm.nih.gov/34398376/>

COMBINED EFFECTS OF PESTICIDES AND ELECTROMAGNETIC-FIELDS ON HONEYBEES: MULTI-STRESS EXPOSURE

Daniela Lupi, Marco Palamara Mesiano, Agnese Adani, Roberto Benocci, Roberto Giacchini, Paolo Parenti, Giovanni Zambon, Antonio Lavazza, Maria B. Boniotti, Stefano Bassi, Mario Colombo, Paolo Tremolada. 2021. Combined Effects of Pesticides and Electromagnetic-Fields on Honeybees: Multi-Stress Exposure. *Insects.*

12(8):716. doi: 10.3390/insects12080716.

Simple Summary

Multi-stress conditions are considered the most putative cause of honeybee decline. The ongoing reduction of domestic and natural pollinators is considered a very severe signal of the current loss of biodiversity, and it requires a broad research effort to clarify the causes. In this research, the combined effects of two possible stress sources for bees, pesticides and electromagnetic fields (multi-stress conditions) were analyzed by a field trial. After one year of monitoring, a complex picture of several induced effects was present, especially in the multi-stress site, such as disease appearance (American foulbrood), higher mortality in the underbaskets (common to pesticide-stress site), behavioral alterations (queen changes, excess of both drone-brood deposition and honey storage) and biochemical anomalies (higher ALP activity at the end of the season). The multi-stress site showed the worst health condition of the bee colonies, with only one alive at the end of the experimentation out of the four ones present at the beginning.

Abstract

Honeybee and general pollinator decline is extensively reported in many countries, adding new concern to the general biodiversity loss. Many studies were addressed to assess the causes of pollinator decline, concluding that in most cases multi-stress effects were the most probable ones. In this research, the combined effects of two possible stress sources for bees, pesticides and electromagnetic fields (multi-stress conditions), were analyzed in the field. Three experimental sites were chosen: a control one far from direct anthropogenic stress sources, a pesticide-stress site and multi-stress one, adding to the same exposure to pesticides the presence of an electromagnetic field, coming from a high-voltage electric line. Experimental apiaries were monitored weekly for one year (from April 2017 to April 2018) by means of colony survival, queen activity, storage and brood amount, parasites and pathogens, and several biomarkers in young workers and pupae. Both exposure and effect biomarkers were analysed: among the first, acetylcholinesterase (AChE), catalase (CAT), glutathione S-transferase (GST) and alkaline phosphatase (ALP) and Reactive Oxygen Species (ROS); and among the last, DNA fragmentation (DNAFRAGM) and lipid peroxidation (LPO). Results showed that bee health conditions were the worst in the multi-stress site with only one colony alive out of the four ones present at the beginning. In this site, a complex picture of adverse effects was observed, such as disease appearance (American foulbrood), higher mortality in the underbaskets (common to pesticide-stress site), behavioral alterations (queen changes, excess of honey storage) and biochemical anomalies (higher ALP activity at the end of the season). The overall results clearly indicate that the multi-stress conditions were able to induce biochemical, physiological and behavioral alterations which severely threatened bee colony survival.

Open access paper: <https://www.mdpi.com/2075-4450/12/8/716>

EFFECTS OF EXTREMELY LOW FREQUENCY ELECTROMAGNETIC FIELD ON DIFFERENT DEVELOPMENTAL STAGES OF DROSOPHILA MELANOGASTER

Neha Agrawal, Kalyani Verma, Doli Baghel, Amitabh Chauhan, Dipti N Prasad, Sanjeev K Sharma, Ekta Kohli. Effects of Extremely Low Frequency Electro Magnetic Field on different developmental stages of *Drosophila melanogaster*. *Int J Radiat Biol.* 2021 Aug 17;1-35. doi: 10.1080/09553002.2021.1969465.

Abstract

Purpose: The model biological organism *Drosophila melanogaster* has been utilized to assess effect of extremely low-frequency electromagnetic field (ELF-EMF) on locomotion, longevity, developmental dynamics, cell viability and oxidative stress.

Materials and method: Developmental stages of *Drosophila melanogaster* (Oregon R strain) individually exposed to ELF-EMF (75 Hz, 550 μ T) for 6 h once for acute exposure. For chronic exposure, complete life cycle of fly, that is, egg to adult fly was exposed to ELF-EMF for 6 h daily. The effect of exposure on their crawling and climbing ability, longevity, development dynamics, cellular damage and oxidative stress (generation of Reactive Oxygen Species (ROS)) was evaluated.

Results: The crawling ability of larvae was significantly ($p < 0.05$) reduced on acute (third stage instar larvae) as well as chronic exposure (F0 and F1 larvae). When locomotion of flies were

tested using climbing assay, no alteration was observed in their climbing ability under both acute and chronic exposure, however, when their speed of climbing was compared, a significant decrease in speed of F1 flies was observed (p value 0.0027) on chronic exposure. The survivability of flies was significantly affected under chronic and acute exposure (at third stage instar larvae). In case of acute exposure of the third stage instar larvae, though all the flies were eclosed by the 17th day, but there was a significant decline in the number of flies (p value 0.007) in comparison to control. While, in case of chronic exposure apart from low number of flies eclosed in comparison to control, there was delay in eclosion by one day (p value 0.0004). Using trypan blue assay the internal gut damage of third stage instar larvae was observed. Under acute exposure condition at third stage instar larvae, 30% larvae has taken up trypan blue, while only 10% larvae from acute exposure at adult stage. On chronic exposure, 50% larvae of the F1 generation have taken up trypan blue. On evaluation of oxidative stress, there is significant rise in ROS in case of acute exposure at third stage instar larvae (p value 0.0004), adult fly stage (p value 0.0004) and chronic exposure (p value 0.0001).

Conclusion: ELF-EMF has maximum effects on acute exposure of third stage instar larvae and chronic exposure (egg to adult fly stage). These results suggest that electromagnetic radiations, though have become indispensable part of our lives but they plausibly effect our health.

<https://pubmed.ncbi.nlm.nih.gov/34402374/>

ELECTROMAGNETIC FIELDS AND CALCIUM SIGNALING BY THE VOLTAGE DEPENDENT ANION CHANNEL

Volker Ullrich, Hans-Jürgen Apell. Electromagnetic Fields and Calcium Signaling by the Voltage Dependent Anion Channel. Open Journal of Veterinary Medicine. 11(1):57-86, January 2021. DOI: 10.4236/ojvm.2021.111004.

Abstract

Electromagnetic fields (EMFs) can interact with biological tissues exerting positive as well as negative effects on cell viability, but the underlying sensing and signaling mechanisms are largely unknown. So far in excitable cells EMF exposure was postulated to cause Ca²⁺ influx through voltage-dependent Ca channels (VDCC) leading to cell activation and an antioxidant response. Upon further activation oxidative stress causing DNA damage or cell death may follow. Here we report collected evidence from literature that voltage dependent anion channels (VDAC) located not only in the outer microsomal membrane but also in the cytoplasmic membrane convert to Ca²⁺ conducting channels of varying capacities upon subtle changes of the applied EMF even in non-excitabile cells like erythrocytes. Thus, VDAC can be targeted by external EMF in both types of membranes to release Ca²⁺ into the cytosol. The role of frequency, pulse modulation or polarization remains to be investigated in suitable cellular models. VDACS are associated with several other proteins, among which the 18 kDa translocator (TSPO) is of specific interest since it was characterized as the central benzodiazepine receptor in neurons. Exhibiting structural similarities with magnetoreceptors we propose that TSPO could sense the magnetic component of the EMF and thus together with VDAC could trigger physiological as well as pathological cellular responses. Pulsed EMFs in the frequency range of the brain-wave communication network may explain psychic disturbances of electromagnetic hypersensitive persons. An important support is provided

from human psychology that states deficits like insomnia, anxiety or depression can be treated with diazepines that indicates apparent connections between the TSPO/VDAC complex and organismic responses to EMF.

Open access paper: <https://www.scirp.org/journal/paperinformation.aspx?paperid=106911>

INDUCED RADIOFREQUENCY FIELDS IN PATIENTS UNDERGOING MRI (MAGNETIC RESONANCE IMAGING) EXAMS: INSIGHTS FOR RISK ASSESSMENT

Aiping Yao, Manuel Murbach, Tolga Goren, Earl Zastrow, Wolfgang Kainz, Niels Kuster. Induced radiofrequency fields in patients undergoing MR examinations: insights for risk assessment. *Phys Med Biol*. 2021 Aug 25. doi: 10.1088/1361-6560/ac212d.

Abstract

Purpose: To characterize and quantify the induced radiofrequency (RF) E-fields and B1+rms fields in patients undergoing magnetic resonance (MR) examinations; to provide guidance on aspects of RF heating risks for patients with and without implants; and to discuss some strengths and limitations of safety assessments in current ISO, IEC, and ASTM standards to determine the RF heating risks for patients with and without implants.

Methods: Induced E-fields and B1+rms fields during 1.5T and 3T MR examinations were numerically estimated for high-resolution patient models of the Virtual Population exposed to ten two-port birdcage RF coils from head to feet imaging landmarks over the full polarization space, as well as in surrogate ASTM phantoms.

Results: Worst-case B1+rms exposure greater than 3.5 μ T (1.5T) and 2 μ T (3T) must be considered for all MR examinations at the Normal Operating Mode limit. Representative induced electric (E)-field and specific absorption rate (SAR) distributions under different clinical scenarios allow quick estimation of clinical factors of high and reduced exposure. B1 shimming can cause +6dB enhancements to E-fields along implant trajectories. The distribution and magnitude of induced E-fields in the ASTM phantom differ from clinical exposures and are not always conservative for typical implant locations.

Conclusions: Field distributions in patient models are condensed, visualized for quick estimation of risks, and compared to those induced in the ASTM phantom. Induced E-fields in patient models can significantly exceed those in the surrogate ASTM phantom in some cases. In the recent 19e2 revision of the ASTM F2182 standard, the major shortcomings of previous versions have been addressed by requiring that the relationship between ASTM test conditions and in vivo tangential E-fields be established, e.g. numerically. With this requirement, the principal methods defined in the ASTM standard for passive implants are reconciled with those of the ISO 10974 standard for active implantable medical devices.

Open access paper: <https://iopscience.iop.org/article/10.1088/1361-6560/ac212d>

HOW THE GEOMAGNETIC FIELD INFLUENCES LIFE ON EARTH - AN INTEGRATED APPROACH TO GEOMAGNETO BIOLOGY

Weronika Erdmann, Hanna Kmita, Jakub Z Kosicki, Łukasz Kaczmarek. How the Geomagnetic Field Influences Life on Earth - An Integrated Approach to Geomagnetobiology. *Orig Life Evol Biosph.* 2021 Aug 7. doi: 10.1007/s11084-021-09612-5.

Abstract

Earth is one of the inner planets of the Solar System, but - unlike the others - it has an oxidising atmosphere, relatively stable temperature, and a constant geomagnetic field (GMF). The GMF does not only protect life on Earth against the solar wind and cosmic rays, but it also shields the atmosphere itself, thus creating relatively stable environmental conditions. What is more, the GMF could have influenced the origins of life: organisms from archaea to plants and animals may have been using the GMF as a source of spatial information since the very beginning. Although the GMF is constant, it does undergo various changes, some of which, e.g. a reversal of the poles, weaken the field significantly or even lead to its short-term disappearance. This may result in considerable climatic changes and an increased frequency of mutations caused by the solar wind and cosmic radiation. This review analyses data on the influence of the GMF on different aspects of life and it also presents current knowledge in the area. In conclusion, the GMF has a positive impact on living organisms, whereas a diminishing or disappearing GMF negatively affects living organisms. The influence of the GMF may also be an important factor determining both survival of terrestrial organisms outside Earth and the emergence of life on other planets.

<https://pubmed.ncbi.nlm.nih.gov/34363564/>

HEALTH IMPACT OF 5G: CURRENT STATE OF KNOWLEDGE OF 5G-RELATED CARCINOGENIC AND REPRODUCTIVE/DEVELOPMENTAL HAZARDS AS THEY EMERGE FROM EPIDEMIOLOGICAL STUDIES AND IN VIVO EXPERIMENTAL STUDIES

Fiorella Belpoggi. Health impact of 5G: Current state of knowledge of 5G-related carcinogenic and reproductive/developmental hazards as they emerge from epidemiological studies and in vivo experimental studies. Panel for the Future of Science and Technology. European Parliamentary Research Service. Scientific Foresight Unit (STOA). PE 690.012. June 2021.

Abstract

The upcoming deployment of 5G mobile networks will allow for significantly faster mobile broadband speeds and increasingly extensive mobile data usage. Technical innovations include a different transmission system (MIMO: use of multiple-input and multiple-output antennas), directional signal transmission or reception (beamforming), and the use of other frequency ranges. At the same time, a change is expected in the exposure to electromagnetic fields (EMF) of humans and the environment. In addition to those used to date, the 5G pioneer bands identified at EU level have frequencies of 700 MHz, 3.6 GHz (3.4 to 3.8 GHz) and 26 GHz (24.25 to 27.5 GHz). The first two frequencies (FR1) are similar to those used for 2G to 4G technologies and have been investigated in both epidemiological and experimental studies for different end points (including carcinogenicity and reproductive/developmental effects), while 26 GHz (FR2) and higher frequencies have not been adequately studied for the same end points.

The International Agency for Research on Cancer (IARC) classified radiofrequency (RF) EMF as 'possibly carcinogenic to humans' (Group 2B) and recently recommended RF exposure for re-evaluation 'with high priority' (IARC, 2019). Since 2011 a great number of studies have been performed, both epidemiological and experimental. The present review addresses the current knowledge regarding both carcinogenic and reproductive/ developmental hazards of RF as exploited by 5G. There are various in vivo experimental and epidemiological studies on RF at a lower frequency range (450 to 6000 MHz), which also includes the frequencies used in previous generations' broadband cellular networks, but very few (and inadequate) on the higher frequency range (24 to 100 GHz, centimetre/MMW).

The review shows: 1) 5G lower frequencies (700 and 3 600 MHz): a) sufficient evidence of carcinogenicity in epidemiological studies; b) sufficient evidence of carcinogenicity in experimental bioassays; c) sufficient evidence of reproductive/developmental adverse effects in humans; d) sufficient evidence of reproductive/ developmental adverse effects in experimental animals; 2) 5G higher frequencies (24.25-27.5 GHz): the systematic review found no adequate studies either in humans or in experimental animals.

Conclusions: 1) cancer: FR1 (450 to 6 000 MHz): EMF are probably carcinogenic for humans, in particular related to gliomas and acoustic neuromas; FR2 (24 to 100 GHz): no adequate studies were performed on the higher frequencies; 2) reproductive developmental effects: FR1 (450 to 6 000 MHz): these frequencies clearly affect male fertility and possibly female fertility too. They may have possible adverse effects on the development of embryos, fetuses and newborns; FR2 (24 to 100 GHz): no adequate studies were performed on non-thermal effects of the higher frequencies.

Open access report:

[https://www.europarl.europa.eu/RegData/etudes/STUD/2021/690012/EPRS_STU\(2021\)690012_EN.pdf](https://www.europarl.europa.eu/RegData/etudes/STUD/2021/690012/EPRS_STU(2021)690012_EN.pdf)

See also: European Parliament: 5G Health Effects and Environmental Impact

ENVIRONMENTAL IMPACTS OF 5G: A LITERATURE REVIEW OF EFFECTS OF RADIO-FREQUENCY ELECTROMAGNETIC FIELD EXPOSURE OF NON-HUMAN VERTEBRATES, INVERTEBRATES AND PLANTS

Arno Thielens. Environmental impacts of 5G: A literature review of effects of radio-frequency electromagnetic field exposure of non-human vertebrates, invertebrates and plants. Panel for the Future of Science and Technology (STOA). European Parliament. 2021, 137 pp. PE 690.021, ISBN 9789284680337. doi: 10.2861/318352.

Abstract

Telecommunication networks use radio-frequency electromagnetic fields to enable wireless communication. These networks have evolved over time, and have been launched in successive generations. The fifth generation of telecommunication networks will operate at frequencies that were not commonly used in previous generations, changing the exposure of wildlife to these waves. This report reviews the literature on the exposure of vertebrates, invertebrates and plants to radio-frequency electromagnetic fields in anticipation of this change.

The review shows that dielectric heating can occur at all considered frequencies (0.4-300 GHz) and for all studied organisms. Summarising and discussing the results of a series of studies of radio-frequency electromagnetic field exposure of wildlife, the review shows that several studies into the effects of radio-frequency electromagnetic field exposure on invertebrates and plants in the frequency bands considered demonstrate experimental shortcomings. Furthermore, the literature on invertebrate and plant exposure to radio-frequency electromagnetic fields above 6 GHz is very limited. More research is needed in this field.

Open access report:

[https://www.europarl.europa.eu/RegData/etudes/STUD/2021/690021/EPRS_STU\(2021\)690021_EN.pdf](https://www.europarl.europa.eu/RegData/etudes/STUD/2021/690021/EPRS_STU(2021)690021_EN.pdf)

DEVELOPMENT OF HEALTH-BASED EXPOSURE LIMITS FOR RADIOFREQUENCY RADIATION FROM WIRELESS DEVICES USING A BENCHMARK DOSE APPROACH

Uloma Igara Uche, Olga V Naidenko. Development of health-based exposure limits for radiofrequency radiation from wireless devices using a benchmark dose approach. *Environ Health* 20, 84 (2021). doi: 10.1186/s12940-021-00768-1.

Abstract

Background Epidemiological studies and research on laboratory animals link radiofrequency radiation (RFR) with impacts on the heart, brain, and other organs. Data from the large-scale animal studies conducted by the U.S. National Toxicology Program (NTP) and the Ramazzini Institute support the need for updated health-based guidelines for general population RFR exposure.

Objectives The development of RFR exposure limits expressed in whole-body Specific Absorption Rate (SAR), a metric of RFR energy absorbed by biological tissues.

Methods Using frequentist and Bayesian averaging modeling of non-neoplastic lesion incidence data from the NTP study, we calculated the benchmark doses (BMD) that elicited a 10% response above background (BMD10) and the lower confidence limits on the BMD at 10% extra risk (BMDL10). Incidence data for individual neoplasms and combined tumor incidence were modeled for 5% and 10% response above background.

Results Cardiomyopathy and increased risk of neoplasms in male rats were the most sensitive health outcomes following RFR exposures at 900 MHz frequency with Code Division Multiple Access (CDMA) and Global System for Mobile Communications (GSM) modulations. BMDL10 for all sites cardiomyopathy in male rats following 19 weeks of exposure, calculated with Bayesian model averaging, corresponded to 0.27–0.42 W/kg whole-body SAR for CDMA and 0.20–0.29 W/kg for GSM modulation. BMDL10 for right ventricle cardiomyopathy in female rats following 2 years of exposure corresponded to 2.7–5.16 W/kg whole-body SAR for CDMA and 1.91–2.18 W/kg for GSM modulation. For multi-site tumor modeling using the multistage cancer model with a 5% extra risk, BMDL5 in male rats corresponded to 0.31 W/kg for CDMA and 0.21 W/kg for GSM modulation.

Conclusion BMDL10 range of 0.2–0.4 W/kg for all sites cardiomyopathy in male rats was selected as a point of departure. Applying two ten-fold safety factors for interspecies and intraspecies variability, we derived a whole-body SAR limit of 2 to 4 mW/kg, an exposure level that is 20–40-fold lower than the legally permissible level of 0.08 W/kg for whole-body SAR under the current U.S. regulations. Use of an additional ten-fold children’s health safety factor points to a whole-body SAR limit of 0.2–0.4 mW/kg for young children.

Open access paper: <https://ehjournal.biomedcentral.com/articles/10.1186/s12940-021-00768-1>

EFFECTS OF MOBILE PHONE USAGE ON SPERM QUALITY - NO TIME-DEPENDENT RELATIONSHIP ON USAGE: A SYSTEMATIC REVIEW AND UPDATED META-ANALYSIS

Sungjoon Kim, Donghyun Han, Jiwoo Ryu, Kihun Kim, Yun Hak Kim. Effects of mobile phone usage on sperm quality - No time-dependent relationship on usage: A systematic review and updated meta-analysis. *Environ Res.* 2021 Jul 29;111784. doi: 10.1016/j.envres.2021.111784.

Abstract

Background: Mobile phones emit radiofrequency (RF) electromagnetic waves (EMWs), a low-level RF that can be absorbed by the human body and exert potential adverse effects on the brain, heart, endocrine system, and reproductive function. Owing to the novel findings of numerous studies published since 2012 regarding the effect of mobile phone use on sperm quality, we conducted a systematic review and updated meta-analysis to determine whether the exposure to RF-EMWs affects human sperm quality.

Methods: This study was conducted in accordance with the PRISMA guidelines. The outcome measures depicting sperm quality were motility, viability, and concentration, which are the most frequently used parameters in clinical settings to assess fertility.

Results: We evaluated 18 studies that included 4280 samples. Exposure to mobile phones is associated with reduced sperm motility, viability, and concentration. The decrease in sperm quality after RF-EMW exposure was not significant, even when the mobile phone usage increased. This finding was consistent across experimental in vitro and observational in vivo studies.

Discussion: Accumulated data from in vivo studies show that mobile phone usage is harmful to sperm quality. Additional studies are needed to determine the effect of the exposure to EMWs from new mobile phone models used in the present digital environment.

<https://pubmed.ncbi.nlm.nih.gov/34333014/>

Excerpts

... 18 studies fulfilled all inclusion criteria and were included in the meta-analysis (Table 1 and Fig. 1) (Agarwal et al., 2008, 2009; Ahmad and Baig, 2011; Al-Bayyari, 2017b; De Iuliis et al., 2009; Ding et al., 2018a; Dkhil et al., 2011; Erogul et al., 2006; Falzone et al., 2008; Fejes et al., 2005; Kaya et al., 2020; Malini, 2017b; Rago et al., 2013; Sajeda and Al-Watter, 2011; Veerachari and Vasan, 2012; Wdowiak et al., 2018; Yildirim et al., 2015; Zalata et al., 2015). Nine studies from a previous meta-analysis and nine new studies that included 4280 samples

were used for analysis. One conference paper included in the previous study was excluded. The sperm quality parameters established in each paper varied and were subjected to a meta-analysis; 16 papers provided data on sperm motility, 6 provided data on sperm viability, and 12 provided data on sperm concentration. All in vitro studies were experimental, whereas all in vivo studies were observational. We identified the MD values of the entire 4280 samples and analyzed the MD values of each group after classifying them according to four criteria: control group setting (non-exposure vs. less exposure), study design (in vivo and in vitro), participant group (fertility clinic and population), and storage location (trousers or not).

Conclusion

Mobile phone use decreased the overall sperm quality by affecting the motility, viability, and concentration. It was further reduced in the group with high mobile phone usage. In particular, the decrease was remarkable in in vivo studies with stronger clinical significance in subgroup analysis. Therefore, long-term cell phone use is a factor that must be considered as a cause of sperm quality reduction. Additional studies are needed to determine the effect of the exposure to EMWs emitted from new mobile phone models in the present digital environment.

ASSOCIATION BETWEEN MOBILE PHONE USE AND HEARING IMPAIRMENT: A SYSTEMATIC REVIEW AND META-ANALYSIS

Mohammad Hosein Taziki Balajelini, Masoud Mohammadi, Abdolhalim Rajabi. *Rev Environ Health*. 2021 Jul 22. doi: 10.1515/reveh-2021-0062.

Abstract

Objectives: To investigate whether a possible association of mobile phone use with hearing impairment was conducted a systematic review and meta-analysis.

Content: This is a systematic review and meta-analysis. A comprehensive literature search was carried out based on the Meta-analysis of Observational Studies in Epidemiology (MOOSE) methodology using PubMed, Scopus, Web of Science, OVID, and Cochrane. The Robins-I tool was used for quality assessment and risk of bias. Two investigators independently reviewed all articles. Pooled effect size was calculated and meta-analysis was performed to compute an overall effect size.

Summary: Overall, five relevant studies (two cross-sectional and three cohort studies) with 92,978 participants were included in the analysis. The studies were stratified by design, there was no significant association between mobile phone use and hearing impairment in cross-sectional studies (OR=0.94, 95% CI=0.57-1.31) and cohort studies (OR=1.09, 95% CI=0.93-1.25). In addition, the effect estimates did not differ significantly between cross-sectional and cohort studies (Q=0.50, p=0.48). Overall, the pooled odds ratio (OR) of hearing impairment was 1.07 (95% CI: 0.94-1.20), which indicates no significant association between mobile phone use and hearing impairment.

Outlook: Our findings indicate no association between mobile phone use and hearing impairment. However, these findings must be interpreted with caution.

<https://pubmed.ncbi.nlm.nih.gov/34293837/>

EFFECTS OF NON-IONIZING ELECTROMAGNETIC FIELDS ON FLORA AND FAUNA, PART 2 IMPACTS: HOW SPECIES INTERACT WITH NATURAL AND MAN-MADE EMF

B. Blake Levitt, Henry C. Lai, Albert M Manville. Effects of non-ionizing electromagnetic fields on flora and fauna, Part 2 impacts: how species interact with natural and man-made EMF. *Rev Environ Health*. 2021 Jul 8.doi: 10.1515/reveh-2021-0050.

Abstract

Ambient levels of nonionizing electromagnetic fields (EMF) have risen sharply in the last five decades to become a ubiquitous, continuous, biologically active environmental pollutant, even in rural and remote areas. Many species of flora and fauna, because of unique physiologies and habitats, are sensitive to exogenous EMF in ways that surpass human reactivity. This can lead to complex endogenous reactions that are highly variable, largely unseen, and a possible contributing factor in species extinctions, sometimes localized. Non-human magnetoreception mechanisms are explored. Numerous studies across all frequencies and taxa indicate that current low-level anthropogenic EMF can have myriad adverse and synergistic effects, including on orientation and migration, food finding, reproduction, mating, nest and den building, territorial maintenance and defense, and on vitality, longevity and survivorship itself. Effects have been observed in mammals such as bats, cervids, cetaceans, and pinnipeds among others, and on birds, insects, amphibians, reptiles, microbes and many species of flora. Cyto- and geno-toxic effects have long been observed in laboratory research on animal models that can be extrapolated to wildlife. Unusual multi-system mechanisms can come into play with non-human species — including in aquatic environments — that rely on the Earth's natural geomagnetic fields for critical life-sustaining information. Part 2 of this 3-part series includes four online supplement tables of effects seen in animals from both ELF and RFR at vanishingly low intensities. Taken as a whole, this indicates enough information to raise concerns about ambient exposures to nonionizing radiation at ecosystem levels. Wildlife loss is often unseen and undocumented until tipping points are reached. It is time to recognize ambient EMF as a novel form of pollution and develop rules at regulatory agencies that designate air as 'habitat' so EMF can be regulated like other pollutants. Long-term chronic low-level EMF exposure standards, which do not now exist, should be set accordingly for wildlife, and environmental laws should be strictly enforced — a subject explored in Part 3.

<https://pubmed.ncbi.nlm.nih.gov/34243228/>

Conclusion

Effects from both natural and man-made EMF over a wide range of frequencies, intensities, wave forms, and signaling characteristics have been observed in all species of animals and plants investigated. The database is now voluminous with in vitro, in vivo, and field studies from which to extrapolate. The majority of studies have found biological effects at both high and low-intensity man-made exposures, many with implications for wildlife health and viability. It is clear that ambient environmental levels are biologically active in all non-human species which can have unique physiological mechanisms that require natural geomagnetic information for their life's most important activities. Sensitive magnetoreception allows living organisms, including plants, to detect small variations in environmental EMF and react immediately as well as over the long term, but it can also make some organisms exquisitely vulnerable to man-made fields. Anthropogenic EMF may be contributing more than we currently realize to species' diminishment and extinction. Exposures continue to escalate

without understanding EMF as a potential causative and/or co-factorial agent. It is time to recognize ambient EMF as a potential novel stressor to other species, design technology to reduce exposures to as low as reasonably achievable, keep systems wired as much as possible to reduce ambient RFR, and create laws accordingly — a subject explored more thoroughly in Part 3.

<https://www.degruyter.com/document/doi/10.1515/reveh-2021-0050/html>

THE EFFECT OF EXPOSURE TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS ON COGNITIVE PERFORMANCE IN HUMAN EXPERIMENTAL STUDIES: A PROTOCOL FOR A SYSTEMATIC REVIEW

Blanka Pophof, Jacob Burns, Heidi Danker-Hopfe, Hans Dorn, Cornelia Egblomassé-Roidl, Torsten Eggert, Kateryna Fuks, Bernd Henschenmacher, Jens Kuhne, Cornelia Sauter, Gernot Schmid. *Environ Int.* 2021 Jul 29;157:106783. doi: 10.1016/j.envint.2021.106783.

Abstract

Background: The World Health Organization (WHO) is currently assessing the potential health effects of exposure to radiofrequency electromagnetic fields (RF-EMFs) in the general and working population. Related to one such health effect, there is a concern that RF-EMFs may affect cognitive performance in humans. The systematic review (SR) aims to identify, summarize and synthesize the evidence base related to this question. Here, we present the protocol for the planned SR.

Objectives: The main objective is to present a protocol for a SR which will evaluate the associations between short-term exposure to RF-EMFs and cognitive performance in human experimental studies.

Data sources: We will search the following databases: PubMed, Embase, Web of Science, Scopus, and the EMF- Portal. The reference lists of included studies and retrieved review articles will be manually searched.

Study eligibility and criteria: We will include randomized human experimental studies that assess the effects of RF-EMFs on cognitive performance compared to no exposure or lower exposure. We will include peer-reviewed articles of any publication date in any language that report primary data.

Data extraction and analysis: Data will be extracted according to a pre-defined set of forms developed and piloted by the review author team. To assess the risk of bias, we will apply the Rating Tool for Human and Animal Studies developed by NTP/OHAT, supplemented with additional questions relevant for cross-over studies. Where sufficiently similar studies are identified (e.g. the heterogeneity concerning population, exposure and outcome is low and the studies can be combined), we will conduct random-effects meta-analysis; otherwise, we will conduct a narrative synthesis.

Assessment of certainty of evidence: The certainty of evidence for each identified outcome will be assessed according to Grading of Recommendations Assessment, Development, and Evaluation (GRADE). Performing the review according to this protocol will allow the identification of possible effects of RF-EMFs on cognitive performance in humans. The protocol has been registered in PROSPERO, an open-source protocol registration system, to foster transparency.

Open access paper:

<https://www.sciencedirect.com/science/article/pii/S0160412021004086?via%3Dihub>

IMPROVING THE QUALITY OF RADIOFREQUENCY BIOEFFECTS RESEARCH: THE NEED FOR A CARROT AND A STICK

Vijayalaxmi, Kenneth R Foster. Improving the Quality of Radiofrequency Bioeffects Research: The Need for a Carrot and a Stick. *Radiat Res.* 2021 Jul 16. doi: 10.1667/RADE-21-00079.1.

Abstract

This commentary considers research needs for radiofrequency (RF) energy above 6 GHz, including in the "high band" of 5G New Radio (NR) communications systems that exists just beneath the mm-wave band (30-300 GHz). As of late 2020, approximately 100 RF bioeffects studies have been published involving exposures above 6 GHz, encompassing a wide range of exposure levels and frequencies. A majority of these studies report statistically significant effects of exposure, many at exposures within international safety limits. This commentary examines 31 genetic damage studies involving RF exposures above 6 GHz in the context of two sets of quality-assessment criteria: 1. "Risk of bias" (RoB) criteria used for systematic reviews of health-related studies; and 2. a broader set of criteria for research quality from a different scholarly approach (metascience). The 31 studies report several statistically significant effects of exposure on different markers for genetic damage. These effects, if real, would have great potential significance for carcinogen risk assessment. However, the studies as a group have significant technical weaknesses, including small size, failure to meet multiple RoB criteria, naïve use of statistics, and lack of prespecified hypotheses and methods of analysis, all of which increase the chances of false discovery. Here we propose a "carrot" (adequate funding to support high-quality research) and a "stick" (more stringent review of bioeffects manuscripts, including explicit instructions to reviewers to assess study quality) approach to increase the reliability of RF bioeffects studies to facilitate health agency reviews of this socially controversial topic.

<https://pubmed.ncbi.nlm.nih.gov/34270779/>

AMERICAN NATIONAL STANDARD FOR METHODS OF MEASUREMENTS OF RADIO-FREQUENCY EMISSIONS FROM WIRELESS POWER TRANSFER EQUIPMENT

"American National Standard for Methods of Measurements of Radio-Frequency Emissions from Wireless Power Transfer Equipment," in ANSI C63.30-2021 , vol., no., pp.1-253, 15 July 2021, doi: 10.1109/IEEESTD.2021.9491984.

Abstract

U.S. consensus standard methods, instrumentation, and facilities for measurement of radio-frequency (RF) emissions and signals emitted from wireless power transfer equipment in the frequency range from 9 kHz to 40 GHz are specified. This standard does not include generi

nor product-specific emission limits. Where possible, the specifications herein are harmonized with other national and international standards used for similar purposes.

Scope This standard includes procedures for evaluating the compliance of wireless power transfer (WPT) equipment with applicable electromagnetic compatibility (EMC) requirements. Test procedures for radiated field strength and conducted disturbance measurements are included, with reference to established standards, where applicable. WPT RF exposure compliance procedures are not included. This document covers measurement methodologies but is not intended to specify regulatory limits. This standard does not consider test methods for the transmitter or receiver portion of any radio apparatus subcomponents that might be included in the equipment under test (EUT), other than those operating at the WPT frequency (or frequencies). These non-WPT radio subcomponents, which can be related or not to the WPT function of the EUT, are covered by other standards (e.g., ANSI C63.10 [B1] and ANSI C63.26 [B2]).

Purpose This standard specifies the methods of measurement to be used for evaluating compliance with the applicable radiated and ac mains power-line conducted emissions requirements for WPT devices that are either already in wide use or soon to be introduced on the market, at the time of publishing this first edition of ANSI C63.30 (such as Qi, PMA, A4WP2, and SAE3 J2954 [B26]) compliant models -- see Annex A). This document includes procedures developed from an exhaustive investigation of such devices, inclusive of field propagation properties, simulation-based analysis, and test site dynamics (see Annex B).

<https://ieeexplore.ieee.org/servlet/opac?punumber=9491982>

USE OF MACHINE LEARNING FOR THE ESTIMATION OF DOWN- AND UP-LINK FIELD EXPOSURE IN MULTI-SOURCE INDOOR WIFI SCENARIOS

Gabriella Tognola, David Plets, Emma Chiaramello, Silvia Gallucci, Marta Bonato, Serena Fiocchi, Marta Parazzini, Luc Martens, Wout Joseph, Paolo Ravazzani. Bioelectromagnetics. 2021 Jul 23. doi: 10.1002/bem.22361.

Abstract

A novel Machine Learning (ML) method based on Neural Networks (NN) is proposed to assess radio-frequency (RF) exposure generated by WiFi sources in indoor scenarios. The aim was to build an NN capable of addressing the complexity and variability of real-life exposure setups, including the effects of not only down-link transmission access points (APs) but also up-link transmission by different sources (e.g. laptop, printers, tablets, and smartphones). The NN was fed with easy to be found data, such as the position and type of WiFi sources (APs, clients, and other users) and the position and material characteristics (e.g. penetration loss) of walls. The NN model was assessed using an additional new layout, distinct from that one used to build and optimize the NN coefficients. The NN model achieved a remarkable field prediction accuracy across exposure conditions in both layouts, with a median prediction error of -0.4 to 0.6 dB and a root mean square error of 2.5-5.1 dB, compared with the target electric field estimated by a deterministic indoor network planner. The proposed approach performs well for the different layouts and is thus generally used to assess RF exposure in indoor scenarios.

Open access paper: <https://onlinelibrary.wiley.com/doi/10.1002/bem.22361>

TIME-TEMPERATURE THRESHOLDS AND SAFETY FACTORS FOR THERMAL HAZARDS FROM RADIOFREQUENCY ENERGY ABOVE 6 GHZ

Kenneth R Foster, Marvin C Ziskin, Quirino Balzano. Time-temperature Thresholds and Safety Factors for Thermal Hazards from Radiofrequency Energy above 6 GHz. *Health Phys.* 2021 Jul 9. doi: 10.1097/HP.0000000000001447.

Abstract

Two major sets of exposure limits for radiofrequency (RF) radiation, those of the International Commission on Nonionizing Radiation Protection (ICNIRP 2020) and the Institute of Electrical and Electronics Engineers (IEEE C95.1-2019), have recently been revised and updated with significant changes in limits above 6 GHz through the millimeter wave (mm-wave) band (30-300 GHz). This review compares available data on thermal damage and pain from exposure to RF energy above 6 GHz with corresponding data from infrared energy and other heat sources and estimates safety factors that are incorporated in the IEEE and ICNIRP RF exposure limits. The benchmarks for damage are the same as used in ICNIRP IR limits: minimal epithelial damage to cornea and first- degree burn (erythema in skin observable within 48 h after exposure). The data suggest that limiting thermal hazard to skin is cutaneous pain for exposure durations less than ≈ 20 min and thermal damage for longer exposures. Limitations on available data and thermal models are noted. However, data on RF and IR thermal damage and pain thresholds show that exposures far above current ICNIRP and IEEE limits would be required to produce thermally hazardous effects. This review focuses exclusively on thermal hazards from RF exposures above 6 GHz to skin and the cornea, which are the most exposed tissues in the considered frequency range.

<https://pubmed.ncbi.nlm.nih.gov/34261892/>

EFFECTS OF 5.8 GHZ MICROWAVE ON HIPPOCAMPAL SYNAPTIC PLASTICITY OF RATS

Gang Rui, Li-Yuan Liu, Ling Guo, Yi-Zhe Xue, Pan-Pan Lai, Peng Gao, Jun-Ling Xing, Jing Li, Gui-Rong Ding. Effects of 5.8 GHz microwave on hippocampal synaptic plasticity of rats. *Int J Environ Health Res.* 2021 Jul 22;1-13. doi: 10.1080/09603123.2021.1952165.

Abstract

Objective 5.8 GHz spectrum is gaining more attention in wireless technology. To explore the potential hazards, we investigated the effect of exposure to 5.8 GHz microwave on learning and memory ability of rats.

Methods Morris Water maze (MWM), Novel object recognition (NOR) and Fear conditioning test (FCT) were used to evaluate the ability of spatial and non-spatial memory of rats. The hippocampal morphology, the level of brain injury factors in serum and the mitochondrial membrane potential of hippocampal neurons was examined to evaluate the damage of hippocampal neurons. The density of dendritic spines, the ultrastructure of synapses and the level of PSD95, Synaptophysin, p-CREB and CREB were detected to evaluate the hippocampal synaptic plasticity.

Results Compared with Sham group, there was no significant difference in the performance of ethology (in MWM, NOR, FCT) in Microwave 2 h group or Microwave 4 h group. The hippocampal morphology, the serum level of brain injury factors and the content of mitochondrial JC-1 monomer in Microwave 2 h group or Microwave 4 h group did not change

obviously, compared with Sham group. The density of dendritic spines, the ultrastructure of synapse and the level of PSD95, Synaptophysin, p-CREB and CREB in hippocampus in Microwave 2 h group or Microwave 4 h group did not significantly change, compared with Sham group.

Conclusion Under this experimental condition, exposure to 5.8 GHz microwave could not affect the hippocampal synaptic plasticity of rats.

<https://pubmed.ncbi.nlm.nih.gov/34293966/>

SPECIFIC ELECTROMAGNETIC RADIATION IN THE WIRELESS SIGNAL RANGE INCREASES WAKEFULNESS IN MICE

Lingyu Liu, Hu Deng, Xiaping Tang, Yingxian Lu, Jiayao Zhou, Xiaofei Wang, Yanyu Zhao, Bing Huang, Yigong Shi. *Proc Natl Acad Sci U S A.* 2021 Aug 3;118(31):e2105838118. doi: 10.1073/pnas.2105838118.

Abstract

Electromagnetic radiation (EMR) in the environment has increased sharply in recent decades. The effect of environmental EMR on living organisms remains poorly characterized. Here, we report the impact of wireless-range EMR on the sleep architecture of mouse. Prolonged exposure to 2.4-GHz EMR modulated by 100-Hz square pulses at a nonthermal output level results in markedly increased time of wakefulness in mice. These mice display corresponding decreased time of nonrapid eye movement (NREM) and rapid eye movement (REM). In contrast, prolonged exposure to unmodulated 2.4-GHz EMR at the same time-averaged output level has little impact on mouse sleep. These observations identify alteration of sleep architecture in mice as a specific physiological response to prolonged wireless-range EMR exposure.

Open access paper: <https://www.pnas.org/content/118/31/e2105838118>

GLIOBLASTOMA CELL MIGRATION IS DIRECTED BY ELECTRICAL SIGNALS

Hannah Clancy, Michal Pruski, Bing Lang, Jared Ching, Colin D McCaig. *Exp Cell Res.* 2021 Jul 14;112736. doi: 10.1016/j.yexcr.2021.112736.

Abstract

Electric field (EF) directed cell migration (electrotaxis) is known to occur in glioblastoma multiforme (GBM) and neural stem cells, with key signalling pathways frequently dysregulated in GBM. One such pathway is EGFR/PI3K/Akt, which is down-regulated by peroxisome proliferator activated receptor gamma (PPAR γ) agonists. We investigated the effect of electric fields on primary differentiated and glioma stem cell (GSCs) migration, finding opposing preferences for anodal and cathodal migration, respectively. We next sought to determine whether chemically disrupting Akt through PTEN upregulation with the PPAR γ agonist, pioglitazone, would modulate electrotaxis of these cells. We found that directed cell migration was significantly inhibited with the addition of pioglitazone in both differentiated GBM and GSCs subtypes. Western blot analysis did not demonstrate any change in PPAR γ expression with and without exposure to EF. In summary we demonstrate opposing EF responses in primary GBM differentiated cells and GSCs can be inhibited chemically by pioglitazone, implicating GBM EF modulation as a potential target in preventing tumour recurrence.<https://pubmed.ncbi.nlm.nih.gov/34273404/>

HEISENBERG UNCERTAINTY OF SPATIALLY GATED ELECTROMAGNETIC FIELDS

Vladimir Y Chernyak, Shaul Mukamel. Heisenberg uncertainty of spatially gated electromagnetic fields. J Chem Phys. 2021 May 7;154(17):174110. doi: 10.1063/5.0045352.

Abstract

A Heisenberg uncertainty relation is derived for spatially-gated electric ΔE and magnetic ΔH field fluctuations. The uncertainty increases for small gating sizes, which implies that in confined spaces, the quantum nature of the electromagnetic field must be taken into account. Optimizing the state of light to minimize ΔE at the expense of ΔH and vice versa should be possible. Spatial confinements and quantum fields may alternatively be realized without gating by interaction of the field with a nanostructure. Possible applications include nonlinear spectroscopy of nanostructures and optical cavities and chiral signals. <https://pubmed.ncbi.nlm.nih.gov/34241063/>

REFLECTION PROPERTIES OF THE HUMAN SKIN FROM 40 TO 110 GHz: A CONFIRMATION STUDY

Andreas Christ, Adrian Aeschbacher, Fereshteh Rouholahnejad, Theodoros Samaras, Bernadetta Tarigan, Niels Kuster. Reflection Properties of the Human Skin From 40 to 110 GHz: A Confirmation Study. Bioelectromagnetics. 2021 Jul 21. doi: 10.1002/bem.22362.

Abstract

Several recent theoretical dosimetric studies above 6 GHz apply generic layered skin models. For this frequency range, new experimental phantoms for over-the-air performance of wireless devices were proposed that simulate the impedance matching effects of the stratum corneum layer (SCL) with a low-loss coating layer. The aim of this study was to verify the skin models by comparing their reflection coefficients S_{11} with measurements of 37 human volunteers (21 males, 16 females, 5-80 years) at 21 body locations (10 at palm, 11 at arm/face) with different SCL thicknesses, using waveguides covering frequencies from 40 to 110 GHz. Such measurements were also carried out with the phantom material. The statistical analysis showed strong evidence that S_{11} depends on the SCL thickness and no evidence that S_{11} depends on sex. The measured S_{11} values for thin and thick skin can be represented by SCL layers of 15 and 140 μm , respectively. These values correspond well to the assumptions of previous studies. (The cohort did not include volunteers doing heavy manual work.) The phantom material mimics the matching effect of the SCL with deviations from the waveguide measurements of less than 0.85 dB (22%), which confirms the suitability of layered phantoms to represent the electromagnetic reflection/absorption of human skin <https://pubmed.ncbi.nlm.nih.gov/34289515/>

ELECTROMAGNETIC FIELD EXPOSURE IN KINDERGARTEN CHILDREN: RESPONSIVE HEALTH RISK CONCERN

Shiva Raj Acharya, Yong Chul Shin, Deog Hwan Moon, Sandip Pahari. Electromagnetic Field Exposure in Kindergarten Children: Responsive Health Risk Concern. *Front Pediatr.* 2021 Jul 5;9:694407. doi: 10.3389/fped.2021.69.

Abstract

Long-term exposure to physical agents can be detrimental to children due to their vulnerability. This study aimed to assess and compare the electromagnetic field (EMF) exposure level around the kindergartens from the underground transmission line (UGTL). We investigated randomly selected 24 kindergartens based on the location of the UGTL. The EMF emission levels were measured using an EMDEX II (Electric and Magnetic Digital Exposure Meter). The maximum mean value of the EMF emission level was 13.5 mG around the kindergartens and 17.7 mG from the point of UGTL to kindergartens. EMF emission level around the kindergartens was significantly associated with the location of the UGTL ($t = -7.35$, $P < 0.001$). These estimates are not trivial, as long-term exposure to EMF among kindergarten children can lead to different health problems. Routine monitoring of EMF emission levels is recommended including the awareness of EMF exposure to public citizens.

Open access paper: <https://www.frontiersin.org/articles/10.3389/fped.2021.694407/full>

RADIOFREQUENCY EXPOSURE LEVELS FROM MOBILE PHONE BASE STATIONS IN OUTDOOR ENVIRONMENTS AND AN UNDERGROUND SHOPPING MALL IN JAPAN

Teruo Onishi, Miwa Ikuyo, Kazuhiro Tobita, Sen Liu, Masao Taki, Soichi Watanabe. Radiofrequency Exposure Levels from Mobile Phone Base Stations in Outdoor Environments and an Underground Shopping Mall in Japan. *Int. J. Environ. Res. Public Health* 2021, 18(15), 8068; doi: 10.3390/ijerph18158068.

(This article belongs to the Special Issue Occupational and General Public Exposure to Electromagnetic Fields)

Abstract

Recent progress in wireless technologies has made human exposure to electromagnetic fields (EMFs) increasingly complex. The situation can increase public concerns related to possible health effects due to EMF exposure. Monitoring EMF exposure levels and characterizing them are indispensable for risk communications of human exposure to EMFs. From this background, a project on the acquisition, accumulation, and applications of EMF exposure monitoring data in Japan was started in 2019. One of the objectives of this project is to obtain a comprehensive picture of EMF exposure in actual daily lives. In 2019 and 2020, we measured the electric field (E-field) strength from mainly mobile phone base stations in the same areas as those in measurements conducted in 2006 and 2007 by the Ministry of Internal Affairs and Communications (MIC), Japan, and compared the data to investigate the time-course of the EMF environment. The number of measured points was 100 (10 × 10 grids) in an area of 1 km × 1 km in two urban and two suburban areas, and that in an underground shopping mall was 158. This large-scale study is the first in Japan. As a result, we found that the measured E-field strengths tended to be higher in 2019 and 2020 than those in 2006 and 2007, especially in the mall. However, the median ratios to the Japanese radio wave protection guideline values for urban areas and malls are lower than −40 dB.

Excerpts

"Measurement results were compared with limits in general environments in the radio wave protection guidelines in Japan, as shown in Table 2. The E-field strength in the radio wave protection guidelines is proportional to the square root of the frequency in between 300 MHz and 1.5 GHz and is constant at 61.4 V/m (155.76 dB μ V/m) at frequencies above 1.5 GHz [25]. The minimum E-field strength in the guidelines at the target frequency of this measurement is 44.1 V/m (152.89 dB μ V/m). Therefore, the ratio to the guideline value in each frequency band and the sum of squares were calculated. A measurement of 0 dB means the same level as the limits. As shown in Table 2, the maximum ratios in the urban areas and the mall are about 10 times higher than those in the suburban areas; however, they were lower than –20 dB from the level of the Japanese guidelines. If we focus on the median value, the ratio is approximately on the order of the –40 dB from the limit."

"The results of E-field strength measurements in mobile phone base stations, which are one of the main sources of EMF exposure in the general environment, namely in outdoor environments and an underground shopping mall in Japan in 2019 and 2020, have been presented. The measurements were conducted in the same areas as those conducted by MIC, Japan, in 2006 and 2007.

As a result, we found that the total median E-field strengths in the urban areas are about 7 dB larger than those in the suburban areas. The E-field strengths in the urban areas also tend to be larger than those in the suburban areas in individual frequency bands. For the shopping mall, it is clear that the differences in E-field strengths between daytime and night-time are marginal. The E-field strengths in the 2000 and 3500 MHz bands are larger than those in other bands. The E-field strengths in the 2400 MHz ISM band are the same as those in the 2000 MHz band, whereas the median E-field strengths in the 5000 MHz ISM bands are 5–9 dB smaller.

The measured results were compared with the Japanese radio wave protection guidelines. As a result, we found amounts lower than the limits by –20 dB. If the median E-field strength is focused on, the ratio is approximately on the order of –40 dB with respect to the limit. Compared with the previous data in 2006 and 2007, it is clear that the E-field strengths in both outdoor environments and the mall increased, especially in the mall, whose ratio of the E-field strength in 2007 to that in 2020 is 20.3 dB. Although the present measurement conditions are not precisely the same as those in the previous measurements, it is useful to investigate the time-course of EMFs in various environments. Additionally, the E-field strengths we obtained in outdoor environments are slightly smaller than those in another country.

Since our measurements were carried out before the start of 5G commercial service, the exposure level at the frequency band used in 5G was not included. However, measurements of exposure levels for 5G and new radio waves used will be continued. Due to the limited time and space of the measurements in this study, exposure level data under various conditions of daily life will be accumulated in the future."

Open access paper: <https://www.mdpi.com/1660-4601/18/15/8068>

EPIGENETIC DYSREGULATION IN VARIOUS TYPES OF CELLS EXPOSED TO EXTREMELY LOW-FREQUENCY MAGNETIC FIELDS (REVIEW)

Gianfranco Giorgi, Brunella Del Re. Epigenetic dysregulation in various types of cells exposed to extremely low- frequency magnetic fields. *Cell Tissue Res.* 2021 Jul 21. doi: 10.1007/s00441-021-03489-6.

Abstract

Epigenetic mechanisms regulate gene expression, without changing the DNA sequence, and establish cell-type- specific temporal and spatial expression patterns. Alterations of epigenetic marks have been observed in several pathological conditions, including cancer and neurological disorders. Emerging evidence indicates that a variety of environmental factors may cause epigenetic alterations and eventually influence disease risks. Humans are increasingly exposed to extremely low-frequency magnetic fields (ELF-MFs), which in 2002 were classified as possible carcinogens by the International Agency for Research on Cancer. This review summarizes the current knowledge of the link between the exposure to ELF-MFs and epigenetic alterations in various cell types. In spite of the limited number of publications, available evidence indicates that ELF-MF exposure can be associated with epigenetic changes, including DNA methylation, modifications of histones and microRNA expression. Further research is needed to investigate the molecular mechanisms underlying the observed phenomena.

<https://pubmed.ncbi.nlm.nih.gov/34287715/>

EFFECT OF THE ELECTROMAGNETIC FIELD (EMF) RADIATION ON TRANSCRIPTOMIC PROFILE OF PIG MYOMETRIUM DURING THE PERI-IMPLANTATION PERIOD-AN IN VITRO STUDY

Ewa Monika Drzewiecka, Wiktoria Kozłowska, Lukasz Paukszto, Agata Zmijewska, Pawel Jozef Wydorski, Jan Pawel Jastrzebski, Anita Franczak. Effect of the Electromagnetic Field (EMF) Radiation on Transcriptomic Profile of Pig Myometrium during the Peri-Implantation Period-An In Vitro Study. *Int J Mol Sci.* 2021 Jul 7;22(14):7322. doi: 10.3390/ijms22147322.

Abstract

The electromagnetic field (EMF) affects the physiological processes in mammals, but the molecular background of the observed alterations remains not well established. In this study was tested the effect of short duration (2 h) of the EMF treatment (50 Hz, 8 mT) on global transcriptomic alterations in the myometrium of pigs during the peri-implantation period using next-generation sequencing. As a result, the EMF treatment affected the expression of 215 transcript active regions (TARs), and among them, the assigned gene protein-coding biotype possessed 90 ones (differentially expressed genes, DEGs), categorized mostly to gene ontology terms connected with defense and immune responses, and secretion and export. Evaluated DEGs enrich the KEGG TNF signaling pathway, and regulation of IFNA signaling and interferon-alpha/beta signaling REACTOME pathways. There were evaluated 12 differentially expressed long non-coding RNAs (DE-lnc-RNAs) and 182 predicted single nucleotide variants (SNVs) substitutions within RNA editing sites. In conclusion, the EMF treatment in the myometrium collected during the peri-implantation period affects the expression of genes involved in defense and immune responses. The study also gives new insight into the mechanisms of the EMF action in the regulation of the transcriptomic profile through lnc-RNAs and SNVs. <https://pubmed.ncbi.nlm.nih.gov/34298942/>

MAGNETIC FIELD AFFECTS GROWTH AND YIELD OF SUNFLOWER UNDER DIFFERENT MOISTURE STRESS CONDITIONS

Ananta Vashisth, Neetu Meena, Prameela Krishnan. Magnetic Field Affects Growth and Yield of Sunflower Under Different Moisture Stress Conditions. Bioelectromagnetics. 26 June 2021/ doi: 10.1002/bem.22354

Abstract

Magnetic field treatments of seeds have shown significant effects on the enhancement of crop growth. Soil moisture stress is the major constraint in the production of the sunflower crop. Therefore, the experiment was conducted to investigate the effect of a 200 mT magnetic field for 2 h on crop growth, and yield of sunflower crops raised from magnetically treated seeds sown under different moisture stress conditions. Results showed that plants from magnetically treated seeds had higher leaf area index, shoot length, number of leaves, chlorophyll content, biomass, 1000-seed mass, and seed yield as compared with untreated. Radiation use efficiency and water productivity were significantly higher in plants raised from magnetically treated seeds than untreated seeds. Crop raised from magnetically treated seeds had 6.2% more seed yield, 7.1% more protein, and oil content as compared with crops raised from untreated seeds. Hence, it may be concluded that exposure of dry sunflower seeds to the static magnetic field of 200 mT for 2 h could be used for improving crop growth and yield under different moisture stress conditions.

<https://onlinelibrary.wiley.com/doi/10.1002/bem.22354>

CHANGES IN BODY TEMPERATURE OF SMALL MAMMALS AND BIRDS IN A FEW MINUTES RANGE AS REFLECTION OF ENVIRONMENTAL INFLUENCES

M E Diatroptov. Changes in Body Temperature of Small Mammals and Birds in a Few Minutes Range as Reflection of Environmental Influences. Bull Exp Biol Med. 2021 Jul 23. doi: 10.1007/s10517-021-05234-z.

Abstract

The study examined the changes in intraperitoneal body temperature of laboratory mice, Jungar hamsters, European greenfinch *Chloris chloris*, and starlings. In a few minutes range, these changes significantly correlated not only between the animals of the same species, but also between the different classes such as birds and mammals, which were isolated from each other and maintained under different illumination regimen. This phenomenon indicates some external influence(s) on the central mechanisms of the thermal control system not related to illumination regiment. In 80% cases, the phases of most pronounced rhythms of body temperature oscillating with the periods of 8-9 and 12-13 min coincided with those of geomagnetic field within the accuracy of ± 1 min. However, the amplitude of body temperature oscillations did not depend on the amplitude of geomagnetic field (GMF) oscillations. Synchronicity of the changes in body temperature and GMF was observed at the amplitude of GMF oscillation of 0.4 nT, which is extremely low value. In contrast, there was no reaction of body temperature to greater (6-10 nT) but irregular and abrupt perturbations of GMF.

<https://pubmed.ncbi.nlm.nih.gov/34297296/>

METHODS AND EXPERIMENTS FOR SENSING VARIATIONS IN SOLAR ACTIVITY AND DEFINING THEIR IMPACT ON HEART VARIABILITY

Michael Hanzelka, Jiří Dan, Zoltán Szabó, Zdeněk Roubal, Přemysl Dohnal, Radim Kadlec. Methods and Experiments for Sensing Variations in Solar Activity and Defining Their Impact on Heart Variability. *Sensors (Basel)*. 2021 Jul 14;21(14):4817. doi: 10.3390/s21144817.

Abstract

This paper evaluates variations in solar activity and their impact on the human nervous system, including the manner in which human behavior and decision-making reflect such effects in the context of (symmetrical) social interactions. The relevant research showed that solar activity, manifesting itself through the exposure of the Earth to charged particles from the Sun, affects heart variability. The evaluation methods focused on examining the relationships between selected psychophysiological data and solar activity, which generally causes major alterations in the low-level electromagnetic field. The investigation within this paper revealed that low-level EMF changes are among the factors affecting heart rate variability and, thus, also variations at the spectral level of the rate, in the VLF, ($f = 0.01\text{--}0.04$ Hz), LF ($f = 0.04\text{--}0.15$ Hz), and HF ($f = 0.15$ až 0.40 Hz) bands. The results of the presented experiments can also be interpreted as an indirect explanation of sudden deaths and heart failures.

<https://pubmed.ncbi.nlm.nih.gov/34300557/>

EFFECTS OF COMMUNICATING UNCERTAINTY DESCRIPTIONS IN HAZARD IDENTIFICATION, RISK CHARACTERIZATION, AND RISK PROTECTION

Peter Wiedemann, Franziska U Boerner, Frederik Freudenstein. Effects of communicating uncertainty descriptions in hazard identification, risk characterization, and risk protection. *PLoS One*. 2021 Jul 13;16(7):e0253762. doi: 10.1371/journal.pone.0253762.

Abstract

Uncertainty is a crucial issue for any risk assessment. Consequently, it also poses crucial challenges for risk communications. Many guidebooks advise reporting uncertainties in risk assessments, expecting that the audience will appreciate this disclosure. However, the empirical evidence about the effects of uncertainty reporting is sparse and inconclusive. Therefore, based on examples of potential health risks of electromagnetic fields (EMF), three experiments were conducted analysing the effects of communicating uncertainties separately for hazard identification, risk characterisation and risk protection. The setups aimed to explore how reporting and how explaining of uncertainty affects dependent variables such as risk perception, perceived competence of the risk assessors, and trust in risk management. Each of the three experiments used a 2x2 design with a first factor presenting uncertainty descriptions (as used in public controversies on EMF related health effects) or describing a certainty conditions; and a second factor explaining the causes of uncertainties (by pointing at knowledge gaps) or not explaining them. The study results indicate that qualitative uncertainty descriptions regarding hazard identification reduce the confidence in the professional competencies of the assessors. In contrast, a quantitative uncertainty description in risk characterisation-regarding the magnitude of the risk-does not affect any of the dependent variables. Concerning risk protection, trust in exposure limit values is not affected by qualitative uncertainty information. However, the qualitative description of

uncertainty regarding the adequacy of protection amplifies fears. Furthermore, explaining this uncertainty results in lower text understandability.

Conclusions

Our experiments provide several implications for reporting uncertainties. They show that informing about uncertainties is not always a means for improving trust and credibility. Science skeptics and activists with a particular political agenda might instrumentalize uncertainty for spreading distrust in science, primarily when uncertainty refers to hazard identification. Risk communicators should be aware that admitting uncertainty is a double-edged sword. Laypersons might attribute it to the lack of professional expertise.

Further research should focus on the issues of cognitive resources and prior beliefs. It should focus on how the recipients interpret the specific social context in which the information is given and how the recipients' cognitive resources, motivations to process information, and their prior beliefs influence uncertainty information interpretation. These issues could be tackled by dual-process theories developed in judgment and decision-making [53]. Recipients might understand the same information differently depending on whether the information is processed by elaborative or heuristic cognition. Furthermore, prior beliefs shape the interpretation of uncertainties in risk assessment. This issue can be tackled by theories that explain how motivated reasoning influences judgments and opinions about uncertainty information [54].

Finally, we would like to stress the importance of the performative goal of informing about uncertainty. It makes a massive difference whether uncertainty is used to raise fears and distrust in science or honestly report the limits of evidence-based knowledge. Further research should pay more attention to this subject. New studies could learn from the analysis of the politicization of science [16] that has focused on how uncertainty is exploited for undermining trust in science.

Open access paper:

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0253762>

EFFECTS OF PULSED ELECTROMAGNETIC FIELDS ON TUMOR CELL VIABILITY: A META-ANALYSIS OF IN VITRO RANDOMIZED CONTROLLED EXPERIMENTS

Guangzhou An, Meilun Shen, Juan Guo, Xia Miao, Yuntao Jing, Keying Zhang, Ling Guo, Junling Xing. *Electromagn Biol Med.* 2021 Jul 26;1-8. doi: 10.1080/15368378.2021.1958341.

Abstract

Malignant tumor treatment remains a big challenge till now, and expanding literature indicated that pulsed electromagnetic fields (PEMF) is promising in tumor treatment with the advantage of safety and being economical, but it is still controversial on whether PEMF could affect the tumor cell viability. Therefore, we conducted the meta-analysis to evaluate effects of PEMF on tumor cell viability. The PubMed, EMBASE, Web of Science, and Cochrane Library databases were searched for studies published up to February 2021. Studies on the direct effects of PEMF on tumor cell viability, determined using colorimetric analysis, were included. Two authors extracted the data and completed the quality assessment. A meta-analysis was performed to calculate the absorbance values and 95% confidence intervals (CIs) using

random-effects models. Seven studies, including 32 randomized controlled experiments, were analyzed. Compared with the control group, tumor cell viability in the PEMF exposure group was obviously lower (SMD, -0.67; 95% CI: -1.12 to -0.22). The subgroup meta-analysis results showed that PEMF significantly reduced epithelial cancer cell viability (SMD, -0.58; 95% CI: -0.92 to -0.23) but had no influence on stromal tumor cell viability (SMD, -0.93; 95% CI: -0.21 to 0.15). Our study demonstrated that PEMF could inhibit tumor cell proliferation to some extent, but the risk of bias and high heterogeneity ($I^2 > 75\%$) weakened the strength of the conclusions drawn from the analysis. <https://pubmed.ncbi.nlm.nih.gov/34311647/>

HEALTH COUNCIL OF THE NETHERLANDS AND EVALUATION OF THE FIFTH GENERATION, 5G, FOR WIRELESS COMMUNICATION AND CANCER RISKS

Lennart Hardell. Health Council of the Netherlands and evaluation of the fifth generation, 5G, for wireless communication and cancer risks. *World J Clin Oncol* 2021; 12(6): 393-403 doi: 10.5306/wjco.v12.i6.393.

Abstract

Currently the fifth generation, 5G, for wireless communication is about to be rolled out worldwide. Many persons are concerned about potential health risks from radiofrequency radiation. In September 2017, a letter was sent to the European Union asking for a moratorium on the deployment until scientific evaluation has been made on potential health risks (<http://www.5Gappeal.eu>). This appeal has had little success. The Health Council of the Netherlands released on September 2, 2020 their evaluation on 5G and health. It was largely based on a World Health Organization draft and report by the Swedish Radiation Safety Authority, both criticized for not being impartial. The guidelines by the International Commission on Non-Ionizing Radiation Protection were recommended to be used, although they have been considered to be insufficient to protect against health hazards (<http://www.emfscientist.org>). The Health Council Committee recommended not to use the 26 GHz frequency band until health risks have been studied. For lower frequencies, the International Commission on Non-Ionizing Radiation Protection guidelines were recommended. The conclusion that there is no reason to stop the use of lower frequencies for 5G is not justified by current evidence on cancer risks as commented in this article. A moratorium is urgently needed on the implementation of 5G for wireless communication.

Core Tip: In this comment, guidelines for radiofrequency radiation are discussed in relation to a recent evaluation by the Health Council of the Netherlands. The Committee recommends that for the deployment of 5G the frequency band 26 GHz should not be used. For lower frequencies, the International Commission on Non- Ionizing Radiation Protection guidelines are recommended. However, these guidelines are not based on an objective evaluation of health risks, which is discussed in this paper.

Conclusion

In conclusion regarding cancer, current scientific evidence clearly demonstrates an increased risk for glioma and acoustic neuroma for use of mobile and/or cordless phones. In this review other tumor types and health endpoints are not discussed. The increased risk for brain and head tumors is based on human cancer epidemiology studies and is supported by similar tumor types found in animal studies. In fact, these animal studies confirmed the earlier results in case-control studies on increased tumor risk for use of wireless phones (both mobile and

cordless phones). Mechanistic aspects on carcinogenesis come from laboratory findings on, e.g., the increase of reactive oxygen species [5] and DNA damage [4]. The current evaluation by the Health Council of the Netherlands is based on a WHO draft and SSM report. It also recommends using ICNIRP guidelines, considered to be insufficient to protect against health hazards, such as cancer, by the majority of the scientists in this field (<https://www.emfscientist.org>). The report does not represent a thorough, balanced, objective, and up-to-date evaluation of cancer risks and other hazardous effects from RF radiation. It is also strikingly contradictory as it concludes that serious health effects such as cancer and birth defects are “possible.” Yet it has no objection to the roll-out of 5G and recommends that later studies are performed to study health outcomes such as cancer and birth defects. Thus, no lessons are learned from existing observations on increased cancer risks [49]. The conclusion by the Commission that there is no reason to stop the use of lower frequencies for 5G up to 3.5 GHz because of no “proven adverse health effects,” merely reflects the biased conclusions by ICNIRP dominated groups. Thus that conclusion must be dismissed, and new guidelines for previous and new frequencies must be established considering the new technology, the different propagation pattern for 5G, and increased RF radiation. A moratorium is urgently required on the implementation of 5G for wireless communication [13]. Ultimately, wired solutions are preferred.

Open access paper: <https://www.wjgnet.com/2218-4333/full/v12/i6/393.htm>

THE CRITICAL IMPORTANCE OF MOLECULAR BIOMARKERS AND IMAGING IN THE STUDY OF ELECTROHYPERSENSITIVITY. A SCIENTIFIC CONSENSUS INTERNATIONAL REPORT

Belpomme, Dominique, George L. Carlo, Philippe Irigaray, David O. Carpenter, Lennart Hardell, Michael Kundi, Igor Belyaev, Magda Havas, Franz Adlkofer, Gunnar Heuser, Anthony B. Miller, Daniela Caccamo, Chiara De Luca, Lebrecht von Klitzing, Martin L. Pall, Priyanka Bandara, Yael Stein, Cindy Sage, Morando Soffritti, Devra Davis, Joel M. Moskowitz, S. M.J. Mortazavi, Martha R. Herbert, Hanns Moshammer, Gerard Ledoigt, Robert Turner, Anthony Tweedale, Pilar Muñoz-Calero, Iris Udasin, Tarmo Koppel, Ernesto Burgio, and André Vander Vorst. 2021. A Scientific Consensus International Report" International Journal of Molecular Sciences 22(14):7321. doi:10.3390/ijms22147321.

Abstract

Clinical research aiming at objectively identifying and characterizing diseases via clinical observations and biological and radiological findings is a critical initial research step when establishing objective diagnostic criteria and treatments. Failure to first define such diagnostic criteria may lead research on pathogenesis and etiology to serious confounding biases and erroneous medical interpretations. This is particularly the case for electrohypersensitivity (EHS) and more particularly for the so-called “provocation tests”, which do not investigate the causal origin of EHS but rather the EHS-associated particular environmental intolerance state with hypersensitivity to man-made electromagnetic fields (EMF). However, because those tests depend on multiple EMF-associated physical and biological parameters and have been conducted in patients without having first defined EHS objectively and/or endpoints adequately, they cannot presently be considered to be valid pathogenesis research methodologies. Consequently, the negative results obtained by these tests do not preclude a role of EMF exposure as a symptomatic trigger in EHS patients. Moreover, there is no proof that EHS symptoms or EHS itself are caused by psychosomatic or nocebo effects. This

international consensus report pleads for the acknowledgement of EHS as a distinct neuropathological disorder and for its inclusion in the WHO International Classification of Diseases.

Open access paper: <https://www.mdpi.com/1422-0067/22/14/7321>

REVIEW OF THE SCIENTIFIC EVIDENCE ON THE INDIVIDUAL SENSITIVITY TO ELECTROMAGNETIC FIELDS (EHS)

Dariusz Leszczynski. Review of the scientific evidence on the individual sensitivity to electromagnetic fields (EHS). *Rev Environ Health*. 2021 Jul 6. doi: 10.1515/reveh-2021-0038.

Abstract

Part of the population considers themselves as sensitive to the man-made electromagnetic radiation (EMF) emitted by powerlines, electric wiring, electric home appliance and the wireless communication devices and networks. Sensitivity is characterized by a broad variety of non-specific symptoms that the sensitive people claim to experience when exposed to EMF. While the experienced symptoms are currently considered as a real life impairment, the factor causing these symptoms remains unclear. So far, scientists were unable to find causality link between symptoms experienced by sensitive persons and the exposures to EMF. However, as presented in this review, the executed to-date scientific studies, examining sensitivity to EMF, are of poor quality to find the link between EMF exposures and sensitivity symptoms of some people. It is logical to consider that the sensitivity to EMF exists but the scientific methodology used to find it is of insufficient quality. It is time to drop out psychology driven provocation studies that ask about feelings-based non-specific symptoms experienced by volunteers under EMF exposure. Such research approach produces only subjective and therefore highly unreliable data that is insufficient to prove, or to disprove, causality link between EHS and EMF. There is a need for a new direction in studying sensitivity to EMF. The basis for it is the notion of a commonly known phenomenon of individual sensitivity, where individuals' responses to EMF depend on the genetic and epigenetic properties of the individual. It is proposed here that new studies, combining provocation approach, where volunteers are exposed to EMF, and high-throughput technologies of transcriptomics and proteomics are used to generate objective data, detecting molecular level biochemical responses of human body to EMF.

Final conclusions

- Problems and solutions for EHS research are summarized in Figure 2.
- Over the last 30 years, a sizable number of research studies has examined causality link between EMF exposures and EHS symptoms.
- The majority of the studies did not find any link between EMF and EHS.
- The EHS studies have examined acute effects but did not have capability to examine delayed EMF responses.
- The major problem is that scientists do not know whether EHS volunteers have indeed correct self-diagnosis of EHS or whether the diagnosis is incorrect and experimental groups are contaminated, to unknown degree, by non-EHS persons. In extreme situation, the small group of volunteers used in research study might have no EHS persons at all.
- Recently, research on EHS has drifted into direction of nocebo as the cause of EHS, what is incorrect.

- Instead of studying obvious impact of placebo, or likes of medical students' disease, research should focus on finding suitable biochemical and biophysical markers that could be used, in combination with single individual- focused provocation studies, to determine the sources of the EHS symptoms.
- The opinion that there is no causality link between EHS and EMF is unproven. This opinion, expressed by the World Health Organization EMF Project, the International Commission on Non-Ionizing Radiation Protection, International Committee on Electromagnetic Safety and numerous governmental organizations, should be revised because the scientific research data
- is of insufficient quality to be used as a proof of the lack of causality.
<https://www.degruyter.com/document/doi/10.1515/reveh-2021-0038/html>

NONMALIGNANT MENINGIOMA AND VESTIBULAR SCHWANNOMA INCIDENCE TRENDS IN THE UNITED STATES, 2004-2017

Diana R Withrow, Susan S Devesa, Dennis Deapen, Valentina Petkov, Alison L Van Dyke, Margaret Adamo, Terri S Armstrong, Mark R Gilbert, Martha S Linet. *Cancer*. 2021 Jun 23. doi: 10.1002/cncr.33553.

Abstract

Background: Given concerns about risks associated with the growing use of mobile phones over recent decades, the authors analyzed temporal trends in incidence rates of nonmalignant meningioma and vestibular schwannoma in the United States.

Methods: The incidence of nonmalignant meningioma and vestibular schwannoma among adults in the Surveillance, Epidemiology, and End Results 18 registries during 2004 through 2017 was evaluated according to the method of diagnosis: microscopically (MC) or radiographically confirmed (RGC). Annual percent changes (APCs) and 95% CIs were estimated using log-linear models.

Results: Overall meningioma rates (n = 108,043) increased significantly from 2004 to 2009 (APC, 5.4%; 95% CI, 4.4%-6.4%) but subsequently rose at a slower pace through 2017 (APC, 1.0%; 95% CI, 0.6%-1.5%). Rates for MC meningiomas changed little from 2004 to 2017 (APC, -0.3%; 95% CI, -0.7%, 0.1%) but rose rapidly for RGC meningiomas until 2009 (APC, 9.5%; 95% CI, 7.8%-11.1%) and rose more modestly thereafter (APC, 2.3%; 95% CI, 1.5%-3.0%). Overall vestibular schwannoma rates (n = 17,475) were stable (APC, 0.4%; 95% CI, -0.2%, 1.0%), but MC vestibular schwannoma rates decreased (APC, -1.9%; 95% CI, -2.7%, -1.1%), whereas RGC vestibular schwannoma rates rose (2006-2017: APC, 1.7%; 95% CI, 0.5%-3.0%). For each tumor, the trends by diagnostic method were similar for each sex and each racial/ethnic group, but RGC diagnosis was more likely in older patients and for smaller tumors. Meningioma trends and the proportion of RGC diagnoses varied notably by registry.

Conclusions: Overall trends obscured differences by diagnostic method in this first large, detailed assessment, but the recent stable rates argue against an association with mobile phone use. Variation among registries requires evaluation to improve the registration of these nonmalignant tumors.

Lay summary: The etiology of most benign meningiomas and vestibular schwannomas is poorly understood, but concerns have been raised about whether mobile phone use contributes to risk of developing these tumors. Descriptive studies examining temporal trends could provide insight; however, globally, few registries collect these nonmalignant cases. In the United States, reporting benign meningiomas and vestibular schwannomas became required by law in 2004. This was the first large, systematic study to quantify and characterize incidence trends for meningioma and vestibular schwannoma according to whether the tumors were diagnosed microscopically or only radiographically. Differential trends across registries and by diagnostic method suggest that caution should be used when interpreting the patterns. <https://pubmed.ncbi.nlm.nih.gov/34160068/>

Excerpts

"Etiologic factors for most benign meningiomas and vestibular schwannomas (acoustic neuromas) are poorly understood, but concerns have long been raised about whether mobile phone use may initiate or promote the occurrence of these central nervous system (CNS) tumors. 1, 2 Analytic epidemiologic studies assessing mobile phone use and the risk of meningiomas have shown no association for a duration of use >10 years (see Supporting Table 1). For vestibular schwannomas, long-term mobile phone use has not been consistently linked with risk, but there is heterogeneity among investigations, and elevated risks were observed in a few studies for a duration of use >10 years (see Supporting Table 2). 2"

"The evaluation of a statistical association between mobile phone use and risk of meningioma or vestibular schwannoma was not feasible because of the descriptive epidemiologic study methods used, along with a lack of detailed data on the history, frequency, and hours per day, per week, or per month of mobile phone use from patients with benign brain tumors and from an appropriate comparison group."

"The potentially large impact of these nonetiologic factors on the incidence trends and the modest increases in meningiomas and stable vestibular schwannoma rates in recent years argue against a substantial effect of the huge increases in mobile phone use on underlying disease risk. This conclusion assumes, however, that the latency period is not many decades in length."

My comment: Although the case-control research examining the association between mobile phone use and meningioma yields mixed results, this study found that the incidence of meningioma in the U.S. was still increasing from 2009 on, just at a slower pace.

The results from the epidemiologic research examining the association between mobile phone use and acoustic neuroma is more consistent; however, the latency for tumor identification can be 20 or more years.

The smartphone started to become popular in the U.S. in 2007 with the first iPhone and in 2008 with Android phones. In many smartphones the cellular transmission antennas are located in the bottom of the handset so the neck, not the head, is likely to receive the greatest microwave radiation exposure among those who place the phone by their ear. Two case-control studies found an association between thyroid cancer and mobile phone use, and thyroid cancer incidence has been rapidly increasing in many countries including the U.S.

IMMUNITY AND ELECTROMAGNETIC FIELDS

Piotr Piszczek, Karolina Wójcik-Piotrowicz, Krzysztof Gil, Jolanta Kaszuba-Zwoińska. Environ Res. 2021 Jun 11;111505. doi: 10.1016/j.envres.2021.111505.

Abstract

Despite many studies, the question about the positive or negative influence of electromagnetic fields (EMF) on living organisms still remains an unresolved issue. To date, the results are inconsistent and hardly comparable between different laboratories. The observed bio-effects are dependent not only on the applied EMF itself, but on many other factors such as the model system tested or environmental ones. In an organism, the role of the defense system against external stressors is played by the immune system consisting of various cell types. The immune cells are engaged in many physiological processes and responsible for the proper functioning of the whole organism. Any factor with an ability to cause immunomodulatory effects may weaken or enhance the response of the immune system. This review is focused on a wide range electromagnetic fields as a possible external factor which may modulate the innate and/or adaptive immunity. Considering the existing databases, we have compiled the bio-effects evoked by EMF in particular immune cell types involved in different types of immune response, with the common mechanistic models and mostly activated intracellular signaling cascade pathways.

<https://pubmed.ncbi.nlm.nih.gov/34126050/>

Highlights

- Immune system cells are influenced by exposure to EMFs.
- EMFs might modulate effector activities of immune response.
- Bio-effectiveness is related to the frequency range of EMFs and cell types.
- Cellular changes might be enhanced by synergic effects of EMFs and other stressors.

Conclusions

Currently it is extremely difficult to select an intracellular mechanism that could play a dominant role in viability and/or effector activities modulation of various types of immune cells under EMF exposure in a wide range of parameters. The large number of results obtained for various EMF parameters and experimental conditions do not allow for a simple comparison of findings across different laboratories. Nevertheless, most of the studies are in agreement that:

- (i) there is no generally accepted physical and/or biological mechanism of EMF action independently on type of the studies (i.e., in vivo/in vitro);
- (ii) there is lack of conclusive evidence of EMF genotoxic effects;
- (iii) findings concerning intracellular effects such as EMF-induced modulation of: gene expression, heat-shock proteins level, surface of cell membrane and cell morphology, signal transduction pathways, ions homeostasis and level of ROS cannot be excluded;
- (iv) significant bio-effects are noticed for simultaneous EMF exposure with other cell stimuli (synergic effects);
- (v) the response of various immune cells differs in an EMF type-dependent manner;
- (vi) multidirectional research on immune cell cultures are certainly needed to be continued to understand potential risk of EMF exposure;
- (vii) the influence of EMF on the innate immunity seems to be interesting issue in the context of aging process (Pawelec et al., 2020).

In summary, EMF seem to be a promising tool for modulation of various immune cell signaling pathways and immune system responses. Moreover, the studies concerning the action of electromagnetic fields alone or combined with medicaments are embedded in the mainstream of interests of EMF-related research in medicine and health care.

MALE CELLULAR TELEPHONE EXPOSURE, FECUNDABILITY, AND SEMEN QUALITY: RESULTS FROM TWO PRECONCEPTION COHORT STUDIES

E E Hatch, S K Willis, A K Wesselink, E M Mikkelsen, M L Eisenberg, G J Sommer, H T Sorensen, K J Rothman, L A Wise. Male cellular telephone exposure, fecundability, and semen quality: results from two preconception cohort studies. *Hum Reprod.* 2021 Apr 20;36(5):1395-1404. doi: 10.1093/humrep/deab001.

Abstract

Study question: To what extent is exposure to cellular telephones associated with male fertility?

Summary answer: Overall, we found little association between carrying a cell phone in the front pants pocket and male fertility, although among leaner men (BMI <25 kg/m²), carrying a cell phone in the front pants pocket was associated with lower fecundability.

What is known already: Some studies have indicated that cell phone use is associated with poor semen quality, but the results are conflicting.

Study design, size, duration: Two prospective preconception cohort studies were conducted with men in Denmark (n = 751) and in North America (n = 2349), enrolled and followed via the internet from 2012 to 2020.

Participants/materials, setting, methods: On the baseline questionnaire, males reported their hours/day of carrying a cell phone in different body locations. We ascertained time to pregnancy via bi-monthly follow-up questionnaires completed by the female partner for up to 12 months or until reported conception. We used proportional probabilities regression models to estimate fecundability ratios (FRs) and 95% confidence intervals (CIs) for the association between male cell phone habits and fecundability, focusing on front pants pocket exposure, within each cohort separately and pooling across the cohorts using a fixed-effect meta-analysis. In a subset of participants, we examined selected semen parameters (semen volume, sperm concentration and sperm motility) using a home-based semen testing kit.

Main results and the role of chance: There was little overall association between carrying a cell phone in a front pants pocket and fecundability: the FR for any front pants pocket exposure versus none was 0.94 (95% CI: 0.0.83-1.05). We observed an inverse association between any front pants pocket exposure and fecundability among men whose BMI was <25 kg/m² (FR = 0.72, 95% CI: 0.59-0.88) but little association among men whose BMI was ≥25 kg/m² (FR = 1.05, 95% CI: 0.90-1.22). There were few consistent associations between cell phone exposure and semen volume, sperm concentration, or sperm motility.

Limitations, reasons for caution: Exposure to radiofrequency radiation from cell phones is subject to considerable non-differential misclassification, which would tend to attenuate the estimates for dichotomous comparisons and extreme exposure categories (e.g. exposure 8 vs. 0 h/day). Residual confounding by occupation or other unknown or poorly measured factors may also have affected the results.

Wider implications of the findings: Overall, there was little association between carrying one's phone in the front pants pocket and fecundability. There was a moderate inverse association between front pants pocket cell phone exposure and fecundability among men with BMI <25 kg/m², but not among men with BMI ≥25 kg/m². Although several previous studies have indicated associations between cell phone exposure and lower sperm motility, we found few consistent associations with any semen quality parameters.

Study funding/competing interest(s): The study was funded by the National Institutes of Health, grant number R03HD090315. In the last 3 years, PRESTO has received in-kind donations from Sandstone Diagnostics (for semen kits), Swiss Precision Diagnostics (home pregnancy tests), Kindara.com (fertility app), and FertilityFriend.com (fertility app). Dr. L.A.W. is a fibroid consultant for AbbVie, Inc. Dr. H.T.S. reports that the Department of Clinical Epidemiology is involved in studies with funding from various companies as research grants to and administered by Aarhus University. None of these studies are related to the current study. Dr. M.L.E. is an advisor to Sandstone Diagnostics, Ro, Dadi, Hannah, and Underdog. Dr. G.J.S. holds ownership in Sandstone Diagnostics Inc., developers of the Trak Male Fertility Testing System. In addition, Dr. G.J.S. has a patent pending related to Trak Male Fertility Testing System issued.

<https://pubmed.ncbi.nlm.nih.gov/33564831/>

SMART DEVICES/MOBILE PHONE IN PATIENTS WITH EPILEPSY? A SYSTEMATIC REVIEW

Ali A Asadi-Pooya, Nafiseh Mirzaei Damabi, Marzieh Rostaminejad, Mina Shahisavandi, Anahita Asadi-Pooya. Smart devices/mobile phone in patients with epilepsy? A systematic review. *Acta Neurol Scand.* 2021 Jun 28. doi: 10.1111/ane.13492.

Abstract

We systematically reviewed the existing literature on the safety of the use of smartphone, mobile phone/Internet, and Wi-Fi by people with epilepsy (PWE), according to the recommendations of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement. Scopus, MEDLINE, and Google Scholar from the inception to April 9, 2021 were searched. These key words were used "epilepsy" OR "seizure" AND "Mobile Phone" OR "Cell Phone" OR "Smartphone" OR "Wi-Fi" OR "Electromagnetic" OR "Radiation." The primary search yielded 7766 studies; 33 studies were related. In total, 19 manuscripts were based on animal/computational studies and 14 articles reported human investigations. Among animal studies, 10 articles suggested detrimental effects by electromagnetic fields (EMFs) on brain function/seizure activity, while nine studies negated this hypothesis. Among human studies, seven studies suggested detrimental effects by EMFs on brain function/seizure activity, while seven studies negated this hypothesis. None of the studies provided a good level of evidence. In one human study, all seven patients with epilepsy and abnormal EEG during the sham exposure, had an increase in the number of epileptic events with exposure to mobile phone radiation. In another study of the detrimental effects of smart

technology device overuse among school students, an association was found between reporting seizures and the hours of smart technology device use. While high- quality evidence on the safety of the use of smartphone, mobile phone/Internet, and Wi-Fi in PWE is lacking, prudent use of these technologies, including using wired hand-free sets or other exposure-reducing measures is recommended.

<https://pubmed.ncbi.nlm.nih.gov/34180044/>

LOW-SAR ANTENNA DESIGN AND IMPLEMENTATION FOR MOBILE PHONE APPLICATIONS

Bao Lu, Bo Pang, Wei Hu, Wen Jiang. Low-SAR Antenna Design and Implementation for Mobile Phone Applications. IEEE Access. 30 June 2021. doi: 10.1109/ACCESS.2021.3093720.

Abstract

This paper presents a novel approach for designing low-SAR (specific absorption rate) terminal antenna. First of all, the relationship between the magnetic field and the SAR value is deduced by using the electric field formula of SAR and Maxwell's equations. Then, by using the boundary conditions of the medium, the interaction mechanism between the radiated magnetic field of the antenna and the magnetic field inside human tissue is studied. Next, the relationship between the surface current and the radiated magnetic field of antenna is constructed. The approach of reducing the antenna radiated magnetic field is proposed by directly adjusting the antenna surface current, so that the SAR of human tissues is reduced. Finally, two low-SAR antennas are designed according to the proposed approach. The simulation and measurement results of the two low-SAR antennas show that the -6 dB bandwidth of both antennas is 2.4-2.49 GHz, which can meet the 2.4 G WLAN frequency band. At the same time, compared with the reference antenna, the peak values of 10 g average SAR of two antennas are reduced by more than 30%. In addition, the measured efficiencies of the two low-SAR antennas are higher than 40% in the operating frequency range. The validity of the proposed approach is proved by the simulation and measurement results of the antenna, which can be used to guide the SAR reduction design of mobile terminal antennas.

Open access paper: <https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9468878>

MAGNETIC INTERFERENCE ON CARDIAC IMPLANTABLE ELECTRONIC DEVICES FROM APPLE IPHONE MAGSAFE TECHNOLOGY

Fahd Nadeem, Arismendy Nunez Garcia, Cao Thach Tran, and Michael Wu. Magnetic Interference on Cardiac Implantable Electronic Devices From Apple iPhone MagSafe Technology. Journal of the American Heart Association. Jun 2, 2021. doi: 10.1161/JAHA.121.020818.

Abstract

Background Magnet wireless charging is being utilized increasingly in current generation smartphones. Apple's MagSafe is a proprietary wireless charging technology with an array of magnets that has the capacity to generate magnet fieldstrength >50 gauss (G). We hypothesize that there is clinically significant magnet interference caused by Apple's MagSafe technology on cardiac implantable electronic devices (CIED).

Methods and Results This study has an in vivo and an ex vivo component. The in vivo component consists of consecutive patients who presented to the electrophysiology

laboratory with previously implanted CIEDs. The iPhone 12 Pro Max was directly placed on the skin over the pocket of these patients and the effect was studied by device interrogation. For the ex vivo component of the study, CIEDs from major device companies were tested for magnetic interference caused by iPhone 12 Pro Max through unopened packages. We found that iPhone 12 Pro Max resulted in clinically identifiable magnet interference in 3/3 (100%) participants in vivo and in 8/11 (72.7%) devices ex vivo.

Conclusions Apple's iPhone 12 Pro Max MagSafe technology can cause magnet interference on CIEDs and has the potential to inhibit lifesaving therapy.
<https://www.ahajournals.org/doi/10.1161/JAHA.121.020818>

Excerpt

Magnet mode activation had been shown to occur in CIED's with exposure to a magnetic field as little as 10 G.⁹ The magnetic field strength of the iPhone 12 Pro Max can be greater than 50 G when in direct contact with the magnetometer. In our ex vivo study, we were able to trigger magnet reversion by placing the iPhone 12 Pro Max at up to 1.5 cm from certain CIED. The difference in magnet response to the iPhone 12 Pro Max among different devices is likely attributed to different hall-sensor magnet sensitivity as all of the devices were susceptible to the standard donut magnet. Boston Scientific Accolade MRI pacemaker for example requires a magnet stronger than 70 G to activate magnet mode.¹⁰

Our case series has several clinical implications. People often put their smartphones in a breast pocket over a device which can be in close proximity to CIEDs. This can lead to asynchronous pacing or disabling of anti-tachycardic therapies. Our study adds to the growing literature demonstrating EMI from magnets in several common technological products such as smart tablets, E-cigarettes, fitness watch wristbands, and wireless headphones.^{11, 12, 13, 14}

Our case series has several limitations. Our sample size is small and we tested on selected device types and the results of our study may not be generalizable. A large scale study should be performed to confirm our findings.

In conclusion, this report highlights the importance of public awareness regarding an interaction between CIEDs and a recently released smartphone model with magnetic charging capability. Although the Food and Drug Administration website states that cellphones do not pose a significant health risk for patients with these devices, they do acknowledge that certain precautions may be advisable.¹⁵ Based on the variability of interactions with respect to different smartphone models, patients are advised to consult with a heart rhythm specialist regarding recommendations specific to their smartphone and CIED.

GENERATIONAL RISK OF COLON AND RECTAL CANCER IN RECENT BIRTH COHORTS UNDER AGE 40 - THE HYPOTHETICAL ROLE OF RADIOFREQUENCY RADIATION FROM CELL PHONES

Devra L Davis, Aaron M. Pilarcik, Anthony B. Miller. Generational Risk of Colon and Rectal Cancer in Recent Birth Cohorts under Age 40 - the Hypothetical Role of Radiofrequency Radiation from Cell Phones. *Ann Gastroenterol Dig Dis*, 3(1): 09-16. 2020.

Abstract

To determine if there are shifts in patterns of cancer, rates of disease can be evaluated in terms of Generational Risk (GR), comparing those born recently with those born decades earlier. Using data from the U.S. Centers for Disease Control and Prevention (CDC), the U.S. Surveillance Epidemiology and End-Results (SEER) Program and Iranian cancer registries, increases in GR of colon and rectal cancer in those under age 50 are presented. For the U.S. those born in the 1990s have a doubled risk of colon cancer (GR=2) and a fourfold increase in rectal cancer (GR=4) by the time they reach age 24 compared to those born six decades ago. Experimental studies have determined that the colon and rectum of Sprague-Dawley rats are exquisitely sensitive to both ionizing and non- ionizing radiofrequency radiation (RFR), expressing significant differences in patterns of methylation of a number of well-identified proteins and other biomarkers predictive of cancer risk. Modeling of nonionizing exposures also indicates that absorption of RFR into the colon and rectum from cell phones stored in the pocket exceeds current test limits by up to 5-fold. French government tests of phones positioned next to the body report exposures to non-ionizing radiation that are up to 11 times more than current guidelines. Based on these findings, it is prudent to develop policies to reduce direct exposures to RFR from cell phones, as occurs when they are kept next to the body, and to promote advances in hardware and software that reduce direct exposures to RFR.

Open access paper: <https://www.somatopublications.com/increased-generational-risk-of-colon-and-rectal-cancer-in-recent-birth-cohorts-under-age-40-the-hypothetical-role-of-radiofrequency-radiation-from-cell-phones.pdf>

THE NON-THERMAL BIOLOGICAL EFFECTS AND MECHANISMS OF MICROWAVE EXPOSURE

X. Zhao, G. Dong, C. Wang. The non-thermal biological effects and mechanisms of microwave exposure. *Int J Radiation Research*. 19(3):475-486. July 2021. doi: 10.18869/acadpub.ijrr.19.2.475.

Abstract

The purpose of this article was to present a brief review of pertinent information regarding the effects of microwave radiation on biological systems. Researchers have been intrigued by the interaction of electromagnetic fields (EMFs) and various life processes since the 18th century. Microwaves refer to the oscillation of an EMF with a wavelength of 1 mm to 1 m, which penetrates matter to varying degrees. With the widespread and ever-increasing use of microwaves, such as cellular telephones and other wireless technologies, great attention and research has been paid to the potential adverse biological effects. It is well recognized that microwaves affect the biological functions of living organisms at both the cellular and molecular levels, and can lead to the appearance of toxicity, genotoxicity and transformation. However, until now no satisfactory mechanism has been proposed to explain the biological effects of these fields. Therefore, increasing attention should be focused on the biological

effects of microwaves in the future, especially since microwaves have extensive applications in various fields. <https://ijrr.com/article-1-3716-en.html>

Conclusions

During recent years there has been increasing public concern regarding the biological effects of microwave radiation emissions from wireless communications. The number of reports on the effects induced by microwave radiation in various cellular and molecular systems continues to increase, attracting public attention toward microwave radiation protection. However, until now no satisfactory mechanism has been proposed to explain the biological effects of these fields. The theories and hypotheses that explain some of the local physiological phenomena have limitations, making it difficult to formulate a specific diagnostic criteria and specific methods of prevention and treatment.

Currently, physical protection is the main measure of EMR protection, including protective clothing with metal wires. In addition, medical interventions through the use of antioxidants are also beneficial for protection against EMR, though specific drugs are lacking. Possible medications include melatonin (28), carnitine, caffeic acid phenethyl ester (CAPE) (59), vitamin C (60) and different kinds of Chinese medicines (61).

Whether mobile phones or other wireless technologies are harmful to humans is still controversial, but research increasingly suggests that they are unlikely to cause brain cancers in adults (62) or affect learning and memory processes (63, 64). In addition, some studies have even found that microwaves emitted from mobile phones can induce adaptive responses that make animals more resistant to ionizing radiation (65). That said, efforts should be made to minimize microwave radiation exposure by remaining remote from TV towers, radar stations and other strong radiation areas. Furthermore, while mobile phone use has not been linked to brain cancer, long- term use may be harmful to the eyes (66) and brain (67). Thus, mobile devices should be kept at a safe distance from the head and, if possible, earphones should be used.

Microwave radiation may also be emitted from microwave ovens, which are widely used for food preparation in daily life. A commonly used frequency in such ovens is 2450 MHz (68). In general, microwave ovens may be considered a safe form of cooking (68); however, improper use can damage the eyes or even result in scald injuries and burns (69). As a safety precaution, a safe distance should be maintained from microwave ovens when in use. Thus, some studies suggest that although current literature lacks definitive evidence associating male infertility with cell phone exposure, limitation of exposure to the possible harmful effects of cell phone, laptop, and microwave ovens is recommended (70).

In conclusion, although the biological effects of microwaves are controversial, future studies regarding the non- thermal effects and mechanisms of microwave radiation are likely to progress our understanding, as well as our comprehension of temporary biological effects that may have long-term impacts on microbial cells, animals and humans. An increased number of studies are paying attention to the biological effects of microwaves and they are expected to increase in number in the future since microwaves have extensive applications in various fields.

A REVIEW OF PROTOCOLS AND GUIDELINES ADDRESSING THE EXPOSURE OF OCCUPANTS TO ELECTROMAGNETIC FIELD RADIATION (EMFR) IN BUILDINGS

Shabnam Monadizadeh, Charles J. Kibert, Jiaxuan Li, Junghoon Woo, Ashish Asutosh, Samira Roostaie, Maryam Kouhirostami. A review of protocols and guidelines addressing the exposure of occupants to electromagnetic field radiation (EMFR) in buildings. *Journal of Green Building*. (2021) 16 (2): 55–81. doi: 10.3992/jgb.16.2.55.

Highlights

- A comprehensive review of increasing electromagnetic field radiation (EMFR) impacts on building occupant health
- Scientific evidence identifying adverse effects of EMFR
- Governments and public health agencies are creating regulations to reduce EMFR exposure
- National and international regulations with respect to the thresholds they set to protect human health
- Provides a robust foundation for researchers to use in additional studies of EMFR impacts in various built environment scenarios

Abstract

A significant share of the technology that has emerged over the past several decades produces electromagnetic field (EMFR) radiation. Communications devices, household appliances, industrial equipment, and medical equipment and devices all produce EMFR with a variety of frequencies, strengths, and ranges. Some EMFR, such as Extremely Low Frequency (ELF), Radio Frequency (RF), and Ionizing Range (IR) radiation have been shown to have harmful effects on human health. Depending on the frequency and strength of the radiation, EMFR can have health effects at the cellular level as well as at brain, nervous, and cardiovascular levels. Health authorities have enacted regulations locally and globally to set critical values to limit the adverse effects of EMFR. By introducing a more comprehensive field of EMFR study and practice, architects and designers can design for a safer electromagnetic (EM) indoor environment, and, as building and construction specialists, will be able to monitor and reduce EM radiation. This paper identifies the nature of EMFR in the built environment, the various EMFR sources, and its human health effects. It addresses European and US regulations for EMFR in buildings and provides a preliminary action plan. The challenges of developing measurement protocols for the various EMFR frequency ranges and determining the effects of EMFR on building occupants are discussed. This paper argues that a mature method for measuring EMFR in building environments and linking these measurements to human health impacts will foster occupant health and lead to the adequate development of safeguards for occupants of buildings in future research.

<https://doi.org/10.3992/jgb.16.2.55>

Dissertation (open access): <https://ufdc.ufl.edu/UFE0052666/00001>

EXPOSURE TO RADIOFREQUENCY ELECTROMAGNETIC FIELDS: COMPARISON OF EXPOSIMETERS WITH A NOVEL BODY-WORN DISTRIBUTED METER

Anke Huss, Stefan Dongus, Reza Aminzadeh, ArnoThielens, Matthias van den Bossche, Patrick Van Torre, René de Seze, Elisabeth Cardis, Marloes Eeftens, Wout Joseph, Roel Vermeulen, MartinRöösli. *Environment International*. 156, November 2021, 106711. doi: 10.1016/j.envint.2021.106711

Highlights

- Personal exposimeters are often used for RF-EMF measurements, but have shortcomings.
- We developed a novel on-body calibrated device with distributed sensors (BWDM).
- BWDM is designed to minimize body shielding and to use frequency specific antennas.
- We measured RF-EMF exposures in parallel with three types of devices.
- Exposimeters provide slightly lower exposures but rank averaged exposures accurately.

Abstract

Background Exposure to radiofrequency electromagnetic fields (RF-EMF) is often measured with personal exposimeters, but the accuracy of measurements can be hampered as carrying the devices on-body may result in body shielding. Further, the compact design may compromise the frequency selectivity of the sensor. The aim of this study was to compare measurements obtained using a multi-band body-worn distributed-exposimeter (BWDM) with two commercially available personal exposimeters (ExpoM-RF and EmeSpy 200) under real-life conditions.

Conclusion

Our device comparisons are informative for the interpretation of existing epidemiological research results and may help future studies regarding how to design improved exposure assessment strategies and how to interpret RF-EMF exposure values reported in the scientific literature. Overall, within the currently applied frequency bands, our study indicates that using one single exposimeter results in slightly lower exposures than measured by a BWDM, most likely due to body shielding, but rank exposure levels reliably. Body shielding could probably at least be in part mitigated by using two exposimeters, on opposite sides of the body (Bhatt et al., 2016). Given their easier handling and relatively compact size, exposimeters thus represent valuable tools to assess population exposure to RF-EMF.

Open access paper:

<https://www.sciencedirect.com/science/article/pii/S0160412021003366?via%3Dihub>

COMPLIANCE ASSESSMENT OF THE EPITHELIAL OR ABSORBED POWER DENSITY BELOW 10 GHZ USING SAR MEASUREMENT SYSTEMS

Theodoros Samaras, Andreas Christ, Niels Kuster. Compliance Assessment of the Epithelial or Absorbed Power Density Below 10 GHz Using SAR Measurement Systems. Bioelectromagnetics. 2021 Jun 15. doi: 10.1002/bem.22355.

Abstract

The introduction of new dosimetric quantities, in particular, epithelial or absorbed power density for frequencies above 6 GHz, in exposure guidelines and safety standards requires the development of new experimental assessment procedures for compliance testing. In this study, we propose to approximate the peak spatial-average absorbed power density (psSab) using the same measured data and algorithms that are used for determining the peak spatial-average specific absorption rate psSAR, which is currently limited to frequencies up to 10 GHz. The uncertainty component for the transformation of psSAR to psSab was evaluated as less than 0.55 dB (13.5%) for any source as close as 0.02λ from the tissue simulating media. The approach is easy to implement and allows determining compliance with the basic restrictions of the latest safety guidelines. In the next project, we will expand dosimetric probes, phantoms, and procedures for frequencies above 10 GHz.

<https://pubmed.ncbi.nlm.nih.gov/34130354/>

EFFECT OF MICROWAVE RADIATION ON EXPERIMENTAL TUMOR GROWTH AT DIFFERENT INTENSITY LEVELS

Vladimir F. Pyankov; Olga V. Kryukova; Alexey F. Kopylov; Gennady M. Aldonin; Yuri P. Salomatov. Engineering (ElConRus).26-29 Jan.2021. doi:10.1109/ElConRus51938.2021.9396077

Abstract

The radiophysical installation designed to study the effect of microwave radiation on biological objects, with an operating frequency of 915 MHz, was used to influence mice with Ehrlich ascites carcinoma. The laboratory animals were exposed to microwave EM radiation of different intensity. The experimental results obtained revealed the dependence of the tumor growth on the intensity of the microwave radiation exposure.

<https://ieeexplore.ieee.org/abstract/document/9396077>

Excerpts

It is shown that the EM radiation exerts a particularly strong impact on the rate of the EAC growth during the first stage of the tumor growth. The highest rate of tumor growth was detected in the animals of the control group, the next was the group exposed to the radiation of $100 \mu\text{W}/\text{cm}^2$, and the lowest rate was observed in the group exposed to $10 \mu\text{W}/\text{cm}^2$. The differences in the tumor growth rate are smoothed over upon reaching the number of 15 thousand cells. Thus, the death of animals with EAC which were exposed to radiation depends on the dose of the microwave radiation....

The data presented in the paper evidences that even in the case of low intensity the effect of microwave EM radiation on an organism with tumor can influence the tumor growth, shifting the stages of its development and, thus, this may influence the life expectancy of animals.

MODERATE EXERCISE TRAINING AS AN EFFECTIVE STRATEGY TO REDUCE THE HARMFUL EFFECTS OF CELL PHONE RADIATION ON WISTAR RAT'S SEMEN QUALITY

HA Akbari, AA Gaeini. Moderate exercise training as an effective strategy to reduce the harmful effects of cell phone radiation on Wistar rat's semen quality. *Int J Radiation Research*. 19(2):317-323. 2021. doi: 10.18869/acadpub.ijrr.19.2.317.

Abstract

Background: The purpose of this study was to evaluate the impact of moderate exercise training as an effective strategy to attenuate the harmful effects of electromagnetic radiation emitted from a cell phone on Wistar Rat's semen quality.

Materials and Methods: Twenty four male Wistar rats (10 weeks old) were randomly assigned to groups: control group, exercise group, radiation group, and radiation plus exercise group. The animals in radiation and radiation plus exercise groups were exposed to radiofrequency electromagnetic radiation of a cell phone 3 hours/day for 28 days. The animals in exercise and radiation plus exercise groups performed moderate exercise training six days a week for 28 days. At the same time, the control and exercise groups exposed to a mobile phone in switch off. Basic parameters of testes weight, sperm count, motility, progressivity, morphology, and viability assessed.

Results: Exposure to the cell phone for 28 days significantly reduced sperm count, progressivity, and normal morphology. Exercise alone caused a significant increase in sperm progressivity only. Radiation plus exercise caused a significant increase in sperm progressivity and morphology compared with the radiation group.

Conclusion: Moderate exercise training may attenuate the harmful effects of exposure to cell phone radiation and enhance sperm quality and the fertility status of men.

<https://ijrr.com/article-1-3646-en.html>

ACUTE EFFECTS OF 2.856 GHZ AND 1.5 GHZ MICROWAVES ON SPATIAL MEMORY ABILITIES AND CREB-RELATED PATHWAYS

Shengzhi Tan, Hui Wang, Xinping Xu, Li Zhao, Jing Zhang, Ji Dong, Binwei Yao, Haoyu Wang, Yanhui Hao, Hongmei Zhou, Yabing Gao, Ruiyun Peng. Acute effects of 2.856 GHz and 1.5 GHz microwaves on spatial memory abilities and CREB-related pathways. *Sci Rep*. 2021 Jun 11;11(1):12348. doi: 10.1038/s41598-021-91622-4.

Abstract

This study aimed to evaluate the acute effects of 2.856 GHz and 1.5 GHz microwaves on spatial memory and cAMP response element binding (CREB)-related pathways. A total of 120 male Wistar rats were divided into four groups: a control group (C); 2.856 GHz microwave exposure group (S group); 1.5 GHz microwave exposure group (L group); and 2.856 and 1.5 GHz cumulative exposure group (SL group). Decreases in spatial memory abilities, changes in EEG, structural injuries, and the downregulation of phosphorylated-Ak strain transforming (p-AKT), phosphorylated-calcium/calmodulin-dependent protein kinase II (p-CaMKII), phosphorylated extracellular signal regulated kinase (p-ERK) and p-CREB was observed 6 h

after microwave exposure. Significant differences in the expression of p-CaMKII were found between the S and L groups. The power amplitudes of the EEG waves (θ , δ), levels of structural injuries and the expression of p-AKT, p-CaMK II, p-CREB, and p-ERK1/2 were significantly different in the S and L groups compared to the SL group. Interaction effects between the 2.856 and 1.5 GHz microwaves were found in the EEG and p-CREB changes. Our findings indicated that 2.856 GHz and 1.5 GHz microwave exposure induced a decline in spatial memory, which might be related to p-AKT, p-CaMK II, p-CREB and p-ERK1/2.
<https://pubmed.ncbi.nlm.nih.gov/34117282/>

The power density in all three RF conditions was 10 milliwatts per square centimeter. Excerpt In terms of the molecular level, differences in the expression of p-CaMKII/CaMKII between 2.856 GHz and 1.5 GHz were found, which suggests the different sensitivities of the signal pathways to microwaves with different frequencies. Interaction effects of the two microwave frequencies were found for the changes in p- CREB/CREB. The accumulative exposure of the 2.856 GHz and 1.5 GHz microwaves aggravated the downregulation of p-CREB/CREB. Previous studies used combined exposure to communication microwaves (849 MHz and 1.95 GHz), but none of these studies analyzed the interaction effects^{53,54,55,56}.

The present study provides new insights into the biological effects of microwave radiation. Microwaves affect multiple metabolic pathways, and the frequency played an important role in the biological effects. The results related to frequency-dependent effects suggested that microwave safety standards should be based on the frequency. Past safety standards were primarily based on radiation power, and radiation frequency was defined only in a rough range⁵⁷. The interaction effects suggested that the biological effects caused by electromagnetic waves were much more complicated than previously thought. More attention should be given to this field.

ELECTROMAGNETIC WAVES FROM MOBILE PHONES MAY AFFECT RAT BRAIN DURING DEVELOPMENT

Dilek Akakin, Olgu Enis Tok, Damla Anil, Akin Akakin, Serap Sirvanci, Goksel Sener, Feriha Ercan. Electromagnetic Waves from Mobile Phones may Affect Rat Brain During Development.

Turk Neurosurg. 2021;31(3):412-421. doi: 10.5137/1019-5149.JTN.31665-20.2.

Abstract

Aim: To investigate the effects of electromagnetic waves (EMWs) from mobile phones (MPs) on rat brains of rats by morphological and biochemical analysis.

Material and methods: EMW was applied for two hours/day until birth in stand-by fetal and EMW fetal groups and postnatal 60th day in stand-by and EMW groups. The control group was not exposed to MP. On postnatal 60th day, brain malondialdehyde (MDA) and glutathione (GSH) levels were measured, and western blot analysis was performed to determine glial fibrillary acidic protein (GFAP) content. Hematoxylin and eosin staining and GFAP immunohistochemistry were applied. Trigeminal nerves were examined using the transmission electron microscope.

Results: In comparison to controls, rats exposed to MP in stand-by or talk modes had significantly increased neuronal damage in the cortex and hippocampus. Increased MDA levels in the EMW group and decreased GSH levels in the stand-by, EMW fetal and EMW groups were found compared with controls. Increased GFAP content in the EMW group and increased GFAP staining in the EMW fetal and EMW groups compared to controls were observed. EMW group had a significantly decreased number of myelinated axons than control animals.

Conclusion: The results of this study suggests that 1800 MHz EMWs (SAR=1.79 W/kg) exposure in the prenatal and early postnatal life may lead to trigeminal nerve damage in addition to oxidative stress-induced neuronal degeneration and astroglial activation in the rat brain. Effects seem to be mode related, being more detrimental in groups exposed to MP during talk mode.

Open access paper: http://www.turkishneurosurgery.org.tr/pdf/pdf_JTN_2485.pdf

RESIDENTIAL EXPOSURE TO ELECTROMAGNETIC FIELDS AND RISK OF AMYOTROPHIC LATERAL SCLEROSIS (ALS): A DOSE-RESPONSE META-ANALYSIS

Tommaso Filippini, Elizabeth E Hatch, Marco Vinceti. Residential exposure to electromagnetic fields and risk of amyotrophic lateral sclerosis: a dose-response meta-analysis. *Sci Rep.* 2021 Jun 7;11(1):11939. doi: 10.1038/s41598-021-91349-2.

Abstract

Amyotrophic lateral sclerosis (ALS) is neurodegenerative disease characterized by a fatal prognosis and still unknown etiology. Some environmental risk factors have been suggested, including exposure to magnetic fields. Studies have suggested positive associations in occupationally-exposed populations, but the link with residential exposure is still debated as is the shape of such relation. Due to recent availability of advanced biostatistical tools for dose-response meta-analysis, we carried out a systematic review in order to assess the dose-response association between ALS and residential exposure to magnetic fields. We performed an online literature searching through April 30, 2021. Studies were included if they assessed residential exposure to electromagnetic fields, based either on distance from overhead power lines or on magnetic field modelling techniques, and if they reported risk estimates for ALS. We identified six eligible studies, four using distance-based and one modelling-based exposure assessment, and one both methods. Both distance-based and particularly modelling-based exposure estimates appeared to be associated with a decreased ALS risk in the highest exposure category, although estimates were very imprecise (summary RRs 0.87, 95% CI 0.63-1.20, and 0.27, 95% CI 0.05-1.36).

Dose-response meta-analysis also showed little association between distance from power lines and ALS, with no evidence of any threshold. Overall, we found scant evidence of a positive association between residential magnetic fields exposure and ALS, although the available data were too limited to conduct a dose-response analysis for the modelled magnetic field estimates or to perform stratified analyses.

Conclusions

Overall, we found little association between exposure to magnetic fields and risk of ALS, using either distance from high-voltage overhead power lines or magnetic field modelling, although the available data were too limited to conduct a dose–response analysis for the modelled exposure studies or to perform further stratified analyses. Therefore, possible associations between magnetic fields exposure and ALS risk in selected subgroups and at very high exposure cannot be entirely ruled out.

Open access paper: <https://www.nature.com/articles/s41598-021-91349-2>

SURFACE ELECTROMAGNETIC WAVES AT GRADUAL INTERFACES BETWEEN LOSSY MEDIA

Igor I. Smolyaninov; Surface Electromagnetic Waves at Gradual Interfaces between Lossy Media. Progress in Electromagnetics Research. 170:177-186. 2021. doi:10.2528/PIER21043006.

Abstract

A low loss propagating electromagnetic wave is shown to exist at a gradual interface between two lossy conductive media. Such a surface wave may be guided by a seafloor-seawater interface, and it may be used in radio communication and imaging underwater. It should allow communication distances of the order of 500m at 10 kHz along a sandy seabed. Similar surface waves may also be guided by various tissue boundaries inside a human body. For example, such surface wave solutions may exist at planar interfaces between skull bones and grey matter inside a human head at 6 GHz.

Conclusion

I have demonstrated that a new kind of low loss propagating surface electromagnetic wave, an “interfacial wave”, may exist at a gradual interface between two lossy conductive media. The comparison of its properties with the more well-known surface electromagnetic wave solutions, such as surface plasmons and Zenneck waves is summarized in Table 1. Such a deep-subwavelength surface wave may be guided by a seafloor- seawater interface, and it may be used in radio communication and imaging underwater. Similar surface waves may also be guided by various tissue boundaries inside a human body. For example, such surface wave solutions may exist at planar interfaces between skull bones and grey matter inside a human head at 6 GHz. A possibility of deep sub-wavelength SEW cavities (or “hot spots”) has been revealed in numerical simulations of SEW-related effects in human tissues. Therefore, it will be important to reexamine EM radiation safety issues associated with the potential excitation and scattering of the newly discovered surface electromagnetic waves inside a human body.

<https://www.jpier.org/PIER/pier.php?paper=21043006>

EFFECT OF THE ELECTRIC FIELD AT 50 HZ AND VARIABLE INTENSITIES ON BIOCHEMICAL MARKERS IN THE HONEY BEE'S HEMOLYMPH

Paweł Migdał, Agnieszka Murawska, Paweł Bieńkowski, Aneta Strachecka, Adam Roman. Effect of the electric field at 50 Hz and variable intensities on biochemical markers in the honey bee's hemolymph. PLoS One. 2021 Jun 24;16(6):e0252858. doi: 10.1371/journal.pone.0252858.

Abstract

The amount of artificial electromagnetic fields of various parameters in the honey bee's environment increases globally. So far, it had been proven that exposure to an E-field at 50 Hz can cause changes in bee's behavior, alter the activity of proteases, and enzymatic antioxidants. Due to the potentially harmful effect of this factor on honey bees, we decided to investigate the activity of aspartate aminotransferase (AST), alanine aminotransferase (ALT), and alkaline phosphatase (ALP), and the concentration of albumin and creatinine in bee's hemolymph after exposure to 50 Hz E-field. Honey bee workers were placed in wooden cages (200 × 150 × 70 mm) and exposed to the 50 Hz E-field with the intensity of <1, 5.0, 11.5, 23.0, or 34.5 kV/m for 1, 3, 6, or 12h. A homogeneous 50 Hz E-field was generated in the form of a plate capacitor. Hemolymph samples for analysis were taken immediately after the end of exposure to the E-field from 100 bees from each group. According to our study, the activity of AST, ALT, and ALP in honey bees' hemolymph decreased after exposure to 50 Hz E-field with various intensities. The decrease in AST, ALT, and ALP activity intensified with prolonged exposure time. 50 Hz E-field may cause the impairment of crucial metabolic cycles in the honey bees' organism (such as the citric acid cycle, ATP synthesis, oxidative phosphorylation, β -oxidation). Moreover, exposure to E-Field altered the concentration of creatinine and albumin, which are important non-enzymatic antioxidants. Such changes may indicate a disturbance in protein metabolism and increased muscle activity.

<https://pubmed.ncbi.nlm.nih.gov/34166412/>

ROLE OF INSULIN/GLUCAGON RATIO AND CELL REDOX STATE IN THE HYPERGLYCAEMIA INDUCED BY EXPOSURE TO A 60-HZ MAGNETIC FIELD IN RATS

Gabriel Martiñón-Gutiérrez, María Luna-Castro, Rolando Hernández-Muñoz. Role of insulin/glucagon ratio and cell redox state in the hyperglycaemia induced by exposure to a 60-Hz magnetic field in rats. Sci Rep. 2021 Jun 3;11(1):11666. doi: 10.1038/s41598-021-91228-w.

Abstract

The exposure to extremely low-frequency electromagnetic fields (EMFs) could adversely affect the endocrine system and cellular proliferative response. Nonetheless, the use of 60-Hz EMFs in the form of magneto-therapy exerts beneficial actions on human health but can also induce hyperglycaemia. Therefore, the present study was aimed to search for metabolic responses of fed or fasted male rats to a single EMF exposure. We performed a 15 min-single exposure to 60-Hz (3.8 mT, intensity) EMF, and determined serum levels of glucose, lipids, and indicators of cellular redox state and energy parameters. A single exposure to a 60-Hz EMF induced hyperglycaemia in both animal groups, and an attenuated second serum insulin peak. The 60-Hz EMF also decreased free fatty acids and lactate serum levels, oppositely increasing pyruvate and acetoacetate levels.

Significant increases in blood glucose level and rat's glucose metabolism were related to a more oxidized cellular redox state and variations in insulin and glucagon secretion. The 60-Hz EMF's effects were not modified in animals previously subjected to chronic EMFs exposure (14 days). In conclusion, increased serum glucose levels and glucose metabolism induced by a single 60-Hz EMF exposure were closely related to the cellular redox state and the insulin/glucagon ratio.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8175349/>

EFFECTS OF LONG-TERM EXPOSURE OF INTERMEDIATE FREQUENCY MAGNETIC FIELDS (20 KHZ, 360 μ T) ON THE DEVELOPMENT, PATHOLOGICAL FINDINGS, AND BEHAVIOR OF FEMALE MICE

Alexander Lerchl, Karen Drees, Née Grote, Isabel Gronau, Dirk Fischer, Julia Bauch, Axel Hoppe. Effects of Long- Term Exposure of Intermediate Frequency Magnetic Fields (20 kHz, 360 μ T) on the Development, Pathological Findings, and Behavior of Female Mice. Bioelectromagnetics. 2021 May;42(4):309-316. doi: 10.1002/bem.22337.

Abstract

The use of magnetic fields in the intermediate-frequency (IF) range to wirelessly charge electric cars with power transfer in the kilowatt range has become increasingly widespread, leading to unavoidable stray fields in the microtesla range. Only a handful of studies have assessed the potential biological risks associated with exposure to such fields. We exposed female mice (n = 80 per group) to either 20 kHz, 360 μ T (rms), or sham in Helmholtz coils to conduct a blind design study. Exposure started at 3 months of age (24 h/day). Body mass was recorded every 1-2 weeks. At 10 months of age, three behavioral tests were performed on 24 animals per group. Three months later, the mice were sacrificed and organs (brain, liver, kidney, spleen, and lung) were removed and prepared for microscopic analysis. Our findings demonstrate no differences in the development of body mass and survival rates (96% and 89%, respectively). Similarly, no significant differences were observed in tumor incidence rates. When it comes to behavioral tests, the 8-arm maze results revealed no significant differences. In contrast, the Rotarod data were significantly ($P < 0.001$) different with longer retention times seen in the exposed mice. In the open field, the number of supported rears was significantly lower ($P < 0.01$), whereas the other endpoints did not show any differences. Overall, our data reveal no adverse effects of exposure to 20 kHz, 360 μ T on the development and tumor incidences, while the significant differences in the behavioral tests may indicate higher levels of alertness in mice.

<https://pubmed.ncbi.nlm.nih.gov/33822410/>

BIOLOGICAL EFFECTS OF CHRONIC EXPOSURE OF BLAPTICA DUBIA (AKA DUBIA ROACH) NYMPHS TO STATIC AND EXTREMELY LOW FREQUENCY MAGNETIC FIELDS

Larisa Ilijin, Marija Mrdaković, Dajana Todorović, Milena Vlahović, Anja Grčić, Aleksandra Filipović, Vesna Perić- Mataruga. An Acad Bras Cienc. 2021 Jun 4;93(2):e20190118. doi: 10.1590/0001- 3765202120190118.

Abstract

In this paper, we analyzed the effects of chronic exposure (5 months) to static magnetic field (110 mT; SMF) and extremely low frequency magnetic field (ELF MF; 10 mT, 50 Hz) on *Blaptica dubia* nymphs. We have examined acetylcholinesterase (AChE) activity and heat shock protein 70 (HSP70) level, two sensitive biomarkers of stress in terrestrial insects. Relative growth rate (RGR), as a life history trait, was estimated. AChE activity was determined spectrophotometrically and HSP70 levels were quantified using indirect non-competitive ELISA and Western blotting. Calculated RGR was significantly changed upon exposure to both types of ambiental MFs. The effects of chronic exposure of *B. dubia* nymphs to SMF and ELF MF (50 Hz) were observed as decreased activity of AChE. The increased level of HSP70 was present only after exposure to SMF. The strength of ELF MF was most likely below the energy level needed to induce the expression of this stress protein. Different patterns of the expression of two HSP70 isoforms, where isoform 2 was sensitive only to SMF, are most likely a possibly switch - off in the expression of constitutive and/or inducible HSP70 isoforms.

Open access paper: <https://www.scielo.br/j/aabc/a/Hbdwv4Fyjdjmxgff6t797yzK/?lang=en>

STATIC MAGNETIC STIMULATION INDUCES CHANGES IN THE OXIDATIVE STATUS AND CELL VIABILITY PARAMETERS IN A PRIMARY CULTURE MODEL OF ASTROCYTES

da Costa CC, Martins LAM, Koth AP, Ramos JMO, Guma FPCR, de Oliveira CM, Pedra NS, Fischer G, Helena ES, Gioda CR, Sanches PRS, Junior ASV, Soares MSP, Spanevello RM, Gamaro GD, de Souza ICC. Static Magnetic Stimulation Induces Changes in the Oxidative Status and Cell Viability Parameters in a Primary Culture Model of Astrocytes. Cell Biochem Biophys. 2021 Jun 27. doi: 10.1007/s12013-021-01015-7

Abstract

Astrocytes play an important role in the central nervous system function and may contribute to brain plasticity response during static magnetic fields (SMF) brain therapy. However, most studies evaluate SMF stimulation in brain plasticity while few studies evaluate the consequences of SMF at the cellular level. Thus, we here evaluate the effects of SMF at 305 mT (medium-intensity) in a primary culture of healthy/normal cortical astrocytes obtained from neonatal (1 to 2-day-old) Wistar rats. After reaching confluence, cells were daily subjected to SMF stimulation for 5 min, 15 min, 30 min, and 40 min during 7 consecutive days. Oxidative stress parameters, cell cycle, cell viability, and mitochondrial function were analyzed. The antioxidant capacity was reduced in groups stimulated for 5 and 40 min. Although no difference was observed in the enzymatic activity of superoxide dismutase and catalase or the total thiol content, lipid peroxidation was increased in all stimulated groups. The cell cycle was changed after 40 min of SMF stimulation while 15, 30, and 40 min led cells to death by necrosis.

Mitochondrial function was reduced after SMF stimulation, although imaging analysis did not reveal substantial changes in the mitochondrial network. Results mainly revealed that SMF compromised healthy astrocytes' oxidative status and viability. This finding reveals how important is to understand the SMF stimulation at the cellular level since this therapeutic approach has been largely used against neurological and psychiatric diseases.

<https://pubmed.ncbi.nlm.nih.gov/34176101/>

STATIC MAGNETIC FIELD (0.2-0.4 T) STIMULATES THE SELF-RENEWAL ABILITY OF OSTEOSARCOMA STEM CELLS THROUGH AUTOPHAGIC DEGRADATION OF FERRITIN

Bin Zhao, Tongyao Yu, Shenghang Wang, Jingmin Che, Liangfu Zhou, Peng Shang. Static Magnetic Field (0.2-0.4 T) Stimulates the Self-Renewal Ability of Osteosarcoma Stem Cells Through Autophagic Degradation of Ferritin. Bioelectromagnetics. 2021 Jun 3. doi: 10.1002/bem.22352.

Abstract

Static magnetic field (SMF) can alter cell fate decisions in many ways. However, the effects of SMF on cancer stem cells (CSCs) are little-known. In this particular study, we evaluate the biological effect of moderate- intensity SMF on osteosarcoma stem cells (OSCs) and try to clarify the underlying mechanisms of action. First, we demonstrated that prolonged exposure to SMF induced the proliferation and tumorsphere formation in K7M2 and MG63 OSCs. Moreover, SMF promoted the release of ferrous iron (Fe^{2+}) and provoked reactive oxygen species (ROS) in OSCs. Interestingly, SMF evidently triggered the autophagic degradation of ferritin, which is characterized by the activation of microtubule-associated protein 1 light chain 3 (LC3) and nuclear receptor co-activator 4 (NCOA4), and downregulation of ferritin heavy chain 1 (FTH1) in OSCs. Particularly, the colony-forming ability of K7M2 OSCs promoted by SMF was obviously abolished by using a small interfering RNA (siRNA) against NCOA4. Finally, treatment of the tumor-bearing mice with SMF did not affect the tumor volume or tumor mass, nor pulmonary metastasis of K7M2 OSCs, but the SMF-treated K7M2 OSCs caused a preference of pulmonary metastasis in a mouse model, which suggested that SMF might induce the metastatic characteristic of OSCs. Consequently, this paper demonstrates for the first time that the cumulative SMF exposure promoted the self-renewal ability of OSCs via autophagic degradation of ferritin, implying that ferritinophagy may be a potential molecular target for cancer. <https://pubmed.ncbi.nlm.nih.gov/34082485/>

Conclusions

Our study has provided, to our knowledge, the first evidence that moderate-intensity SMF stimulated the self- renewal ability of OSCs through autophagic degradation of ferritin (Fig. 7a). The long-term exposure to moderate-intensity SMF did not affect the tumor volume or tumor mass, nor the lung metastasis in tumor- bearing mice, but the OSCs exposed to SMF did cause a preference of pulmonary metastasis in tumor-bearing mice (Fig. 7b). Finally, it is important to state that the effects of SMF on OSCs here are not directly applicable to humans and should be interpreted with caution.

THE ROLE OF PULSED ELECTROMAGNETIC FIELDS ON THE RADICAL PAIR MECHANISM

Pablo Castello, Pablo Jimenez, Carlos F Martino. The Role of Pulsed Electromagnetic Fields on the Radical Pair Mechanism. Bioelectromagnetics. 2021 Jul 5. doi: 10.1002/bem.22358.

Abstract

In recent decades, the use of pulsed electromagnetic fields (PEMF) in therapeutics has been one of the main fields of activity in the bioelectromagnetics arena. Nevertheless, progress in this area has been hindered by the lack of consensus on a biophysical mechanism of interaction that can satisfactorily explain how low-level, non-thermal electromagnetic fields would be able to sufficiently affect chemistry as to elicit biological effects in living organisms. This specifically applies in cases where the induced electric fields are too small to generate a biological response of any consequence. A growing body of experimental observations that would explain the nature of these effects speaks strongly about the involvement of a theory known as the radical pair mechanism (RPM). This mechanism explains how a pair of reactive oxygen species with distinct chemical fate can be influenced by a low-level external magnetic field through Zeeman and hyperfine interactions. So far, a study of the effects of complex spatiotemporal signals within the context of the RPM has not been performed. Here, we present a computational investigation of such effects by utilizing a generic PEMF test signal and RPM models of different complexity. Surprisingly, our results show how substantially different chemical results can be obtained within ranges that depend on the specific orientation of the PEMF test signal with respect to the background static magnetic field, its waveform, and both of their amplitudes. These results provide a basis for explaining the distinctive biological relevance of PEMF signals on radical pair chemical reactions.

Conclusion

Experimental observations speak strongly for the involvement of the radical pair mechanism in biological systems. For this purpose, we computationally studied whether a pulse train waveform can change the quantum singlet yields in a radical pair reaction. For a simple radical pair model, we demonstrated that the suggested reaction can be influenced by PEMFs.

This conclusion does not rule out the possibility of induced electric field effects stemming from PEMFs. However, experimental evidence suggests controversial results with the use of PEMFs that cannot be explained by the accepted mechanism of action [Barnes and Greenebaum, 2018]. Our study establishes the role of PEMF as a diagnostic tool that may indicate the involvement of magneto-sensitive radical pair reactions in biological systems. Extending this tool to determine orientation and amplitude dependence in which the input PEMF waveforms affect the reaction products can reveal the chemical nature of the radical pairs involved. Finally, using the oscillating or PEMF input waveform as a diagnostic tool to modify singlet quantum yields can easily be transferred to finding the optimal control to maximize the singlet yield. At the most fundamental level, one could investigate how a radical reaction can be controlled by perturbing spin interconversion to maximize a cost functional, the quantum singlet yield, through the selection of optimal control functions, namely the magnetic waveform.

<https://pubmed.ncbi.nlm.nih.gov/34224591/>

MAGNETIC SENSITIVITY OF CRYPTOCHROME 4 FROM A MIGRATORY SONGBIRD

Xu J, Jarocha LE, Zollitsch T, Konowalczyk M, Henbest KB, Richert S, Golesworthy MJ, Schmidt J, Déjean V, Sowood DJC, Bassetto M, Luo J, Walton JR, Fleming J, Wei Y, Pitcher TL, Moise G, Herrmann M, Yin H, Wu H, Bartölke R, Käsehagen SJ, Horst S, Dautaj G, Murton PDF, Gehrckens AS, Chelliah Y, Takahashi JS, Koch KW, Weber S, Solov'yov IA, Xie C, Mackenzie SR, Timmel CR, Mouritsen H, Hore PJ. Magnetic sensitivity of cryptochrome 4 from a migratory songbird. *Nature*. 2021 Jun;594(7864):535-540. doi: 10.1038/s41586-021-03618-9.

Abstract

Night-migratory songbirds are remarkably proficient navigators¹. Flying alone and often over great distances, they use various directional cues including, crucially, a light-dependent magnetic compass^{2,3}. The mechanism of this compass has been suggested to rely on the quantum spin dynamics of photoinduced radical pairs in cryptochrome flavoproteins located in the retinas of the birds⁴⁻⁷. Here we show that the photochemistry of cryptochrome 4 (CRY4) from the night-migratory European robin (*Erithacus rubecula*) is magnetically sensitive in vitro, and more so than CRY4 from two non-migratory bird species, chicken (*Gallus gallus*) and pigeon (*Columba livia*). Site-specific mutations of ErCRY4 reveal the roles of four successive flavin-tryptophan radical pairs in generating magnetic field effects and in stabilizing potential signalling states in a way that could enable sensing and signalling functions to be independently optimized in night-migratory birds.

<https://pubmed.ncbi.nlm.nih.gov/34163056/>

THE MAGNETIC COMPASS OF BIRDS: THE ROLE OF CRYPTOCHROME

Roswitha Wiltschko, Christine Nießner, Wolfgang Wiltschko. The Magnetic Compass of Birds: The Role of Cryptochrome. *Front Physiol*. 2021 May 19;12:667000. doi: 10.3389/fphys.2021.667000.

Abstract

The geomagnetic field provides directional information for birds. The avian magnetic compass is an inclination compass that uses not the polarity of the magnetic field but the axial course of the field lines and their inclination in space. It works in a flexible functional window, and it requires short-wavelength light. These characteristics result from the underlying sensory mechanism based on radical pair processes in the eyes, with cryptochrome suggested as the receptor molecule. The chromophore of cryptochrome, flavin adenine dinucleotide (FAD), undergoes a photocycle, where radical pairs are formed during photo-reduction as well as during re-oxidation; behavioral data indicate that the latter is crucial for detecting magnetic directions. Five types of cryptochromes are found in the retina of birds: cryptochrome 1a (Cry1a), cryptochrome 1b, cryptochrome 2, cryptochrome 4a, and cryptochrome 4b. Because of its location in the outer segments of the ultraviolet cones with their clear oil droplets, Cry1a appears to be the most likely receptor molecule for magnetic compass information.

Outlook

More than 50 years after the discovery of the avian magnetic compass, we finally have a concrete idea about the primary mechanism leading to the detection of directions by the magnetic field in birds: a radical pair process, with cryptochrome playing a crucial role in this process. Yet there are still several conflicting findings that have to be resolved, and a number of open questions that have to be answered.

One of the most important questions concerns the mechanisms by which the information obtained by the radical pair is transformed into a biological signal. Cryptochrome is found in the outer segments of cones, i.e., in a cell type that is also important for color vision. Is the visual information from the opsin and information on magnetic directions from cryptochrome transmitted separately or together? One might expect the latter since the cone has only one known way of transmitting information. This would mean, however, the two types of information have somehow to be separated, either already in the eyes or later in higher centers in the brain.

There are speculations about possibilities (see, e.g., Bischof et al., 2011), but it will require many more considerations and experiments until we hopefully reach a complete understanding of how birds perceive the direction of the geomagnetic field.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8171495/>

UNRAVELLING THE ENIGMA OF BIRD MAGNETORECEPTION

Eric J Warrant. Unravelling the enigma of bird magnetoreception. *Nature*. 2021 Jun;594(7864):497-498. doi: 10.1038/d41586-021-01596-6. doi: 10.1038/d41586-021-01596-6

No abstract. <https://pubmed.ncbi.nlm.nih.gov/34163048/>

RANDOM EFFECTS IN MAGNETOBIOLGY AND A WAY TO SUMMARIZE THEM

Vladimir N Binhi. Random Effects in Magnetobiology and a Way to Summarize Them. *Bioelectromagnetics*. 2021 Jul 7. doi: 10.1002/bem.22359.

Abstract

In magnetobiology, it is difficult to reproduce the nonspecific (not associated with specialized receptors) biological effects of weak magnetic fields. This means that some important characteristic of the data may be missed in standard statistical processing, where the set of measurements to be averaged belongs to the same population so that the contribution of fluctuations decreases according to the Central Limit Theorem. It has been shown that a series of measurements of a nonspecific magnetic effect contains not only the usual scatter of data around the mean but also a significant random component in the mean itself. This random component indicates that measurements belong to different statistical populations, which requires special processing. This component, otherwise called heterogeneity, is an additional characteristic that is typically overlooked, and which reduces reproducibility. The current method for studying and summarizing highly heterogeneous data is the random-effect meta-analysis of absolute values, i.e., of magnitudes, rather than the values themselves.

However, this estimator-the average of absolute values-has a significant positive bias when it comes to the small effects that are characteristic of magnetobiology. To solve this problem, an improved estimator based on the folded normal distribution that gives several times less bias is proposed. We used this improved estimator to analyze the nonspecific effect of the hypomagnetic field in the Stroop test in 40 subjects and found a statistically significant meta-

effect with a standardized average of magnitudes of about 0.1. It has been shown that the proposed approach can also be applied to a single study.

<https://pubmed.ncbi.nlm.nih.gov/34233018/>

EFFECTS OF NON-IONIZING ELECTROMAGNETIC FIELDS ON FLORA AND FAUNA, PART 1. RISING AMBIENT EMF LEVELS IN THE ENVIRONMENT

B. Blake Levitt, Henry C. Lai, Albert M. Manville. Effects of non-ionizing electromagnetic fields on flora and fauna, part 1. Rising ambient EMF levels in the environment. *Rev Environ Health*. 2021 May 27. doi: 10.1515/reveh- 2021-0026.

Abstract

Ambient levels of electromagnetic fields (EMF) have risen sharply in the last 80 years, creating a novel energetic exposure that previously did not exist. Most recent decades have seen exponential increases in nearly all environments, including rural/remote areas and lower atmospheric regions. Because of unique physiologies, some species of flora and fauna are sensitive to exogenous EMF in ways that may surpass human reactivity.

There is limited, but comprehensive, baseline data in the U.S. from the 1980s against which to compare significant new surveys from different countries. This now provides broader and more precise data on potential transient and chronic exposures to wildlife and habitats. Biological effects have been seen broadly across all taxa and frequencies at vanishingly low intensities comparable to today's ambient exposures. Broad wildlife effects have been seen on orientation and migration, food finding, reproduction, mating, nest and den building, territorial maintenance and defense, and longevity and survivorship. Cyto- and geno-toxic effects have been observed. The above issues are explored in three consecutive parts: Part 1 questions today's ambient EMF capabilities to adversely affect wildlife, with more urgency regarding 5G technologies. Part 2 explores natural and man-made fields, animal magnetoreception mechanisms, and pertinent studies to all wildlife kingdoms. Part 3 examines current exposure standards, applicable laws, and future directions. It is time to recognize ambient EMF as a novel form of pollution and develop rules at regulatory agencies that designate air as 'habitat' so EMF can be regulated like other pollutants. Wildlife loss is often unseen and undocumented until tipping points are reached. Long-term chronic low-level EMF exposure standards, which do not now exist, should be set accordingly for wildlife, and environmental laws should be strictly enforced.

<https://pubmed.ncbi.nlm.nih.gov/34047144/>

Conclusion

Ambient background levels of EMF have risen sharply in the last four decades, creating a novel energetic exposure that previously did not exist at the Earth's surface, lower atmospheric levels, or underwater environments. Recent decades have seen exponential increases in nearly all environments, including remote regions. There is comprehensive but outdated baseline data from the 1980s against which to compare significant new surveys from other countries which found increasing RFR levels in urban, suburban and remote areas, primarily from cell infrastructure/phone/WiFi exposures. One indicative comparison of similar sites between 1980 and today found a 70-fold (7,000%) increase in ambient RFR [149]. The increased infrastructure required for 5G networks will widely infuse the environment with new atypical exposures, as are increasing satellite systems communicating with ground-based

civilian networks. The new information provides broader perspective with more precise data on both potential transient and chronic exposures to wildlife and habitats. Biological effects have been seen broadly across all taxa at vanishingly low intensities comparable to today's ambient exposures as examined in Part 2. The major question presented in Part 1 was whether increasing anthropogenic environmental EMF can cause biological effects in wildlife that may become more urgent with 5G technologies, in addition to concerns over potentially more lenient allowances being considered by major standards-setting committees at FCC and ICNIRP (examined in Part 3). There are unique signaling characteristics inherent to 5G transmission as currently designed of particular concern to non-human species. Background levels continue to rise but no one is studying cumulative effects to nonhuman species.

MILLIMETER (MM) WAVE AND MICROWAVE FREQUENCY RADIATION PRODUCE DEEPLY PENETRATING EFFECTS: THE BIOLOGY AND THE PHYSICS

Martin L Pall. Millimeter (MM) wave and microwave frequency radiation produce deeply penetrating effects: the biology and the physics. *Rev Environ Health*. 2021 May 26. doi: 10.1515/reveh-2020-0165.

Abstract

Millimeter wave (MM-wave) electromagnetic fields (EMFs) are predicted to not produce penetrating effects in the body. The electric but not magnetic part of MM-EMFs are almost completely absorbed within the outer 1 mm of the body. Rodents are reported to have penetrating MM-wave impacts on the brain, the myocardium, liver, kidney and bone marrow. MM-waves produce electromagnetic sensitivity-like changes in rodent, frog and skate tissues. In humans, MM-waves have penetrating effects including impacts on the brain, producing EEG changes and other neurological/neuropsychiatric changes, increases in apparent electromagnetic hypersensitivity and produce changes on ulcers and cardiac activity. This review focuses on several issues required to understand penetrating effects of MM-waves and microwaves: 1. Electronically generated EMFs are coherent, producing much higher electrical and magnetic forces than do natural incoherent EMFs. 2. The fixed relationship between electrical and magnetic fields found in EMFs in a vacuum or highly permeable medium such as air, predicted by Maxwell's equations, breaks down in other materials. Specifically, MM-wave electrical fields are almost completely absorbed in the outer 1 mm of the body due to the high dielectric constant of biological aqueous phases. However, the magnetic fields are very highly penetrating. 3. Time-varying magnetic fields have central roles in producing highly penetrating effects. The primary mechanism of EMF action is voltage-gated calcium channel (VGCC) activation with the EMFs acting via their forces on the voltage sensor, rather than by depolarization of the plasma membrane. Two distinct mechanisms, an indirect and a direct mechanism, are consistent with and predicted by the physics, to explain penetrating MM-wave VGCC activation via the voltage sensor. Time-varying coherent magnetic fields, as predicted by the Maxwell-Faraday version of Faraday's law of induction, can put forces on ions dissolved in aqueous phases deep within the body, regenerating coherent electric fields which activate the VGCC voltage sensor. In addition, time-varying magnetic fields can directly put forces on the 20 charges in the VGCC voltage sensor. There are three very important findings here which are rarely recognized in the EMF scientific literature: coherence of electronically generated EMFs; the key role of time-varying magnetic fields in generating highly penetrating effects; the key role of both modulating and pure EMF pulses in greatly increasing very short term high level time-variation of magnetic and electric

fields. It is probable that genuine safety guidelines must keep nanosecond timescale-variation of coherent electric and magnetic fields below some maximum level in order to produce genuine safety. These findings have important implications with regard to 5G radiation. <https://pubmed.ncbi.nlm.nih.gov/34043892>

ASPECTS ON THE INTERNATIONAL COMMISSION ON NON-IONIZING RADIATION PROTECTION (ICNIRP) 2020 GUIDELINES ON RADIOFREQUENCY RADIATION

Lennart Hardell, Mona Nilsson, Tarmo Koppel, Michael Carlberg. Aspects on the International Commission on Non-Ionizing Radiation Protection (ICNIRP) 2020 Guidelines on Radiofrequency Radiation. *J Cancer Sci Clin Ther.* 2021; 5(2): 250-285. doi: 10.26502/jcsct.5079117.

Abstract

The International Commission on Non-Ionizing Radiation Protection (ICNIRP) published 2020 updated guidelines on radiofrequency (RF) radiation in the frequency range 100 kHz to 300 GHz. Harmful effects on human health and the environment at levels below the guidelines are downplayed although evidence is steadily increasing.

Only thermal (heating) effects are acknowledged and therefore form the basis for the guidelines. Despite the increasing scientific evidence of non-thermal effects, the new ICNIRP guidelines are not lower compared with the previous levels. Expert groups from the WHO, the EU Commission and Sweden are to a large extent made up of members from ICNIRP, with no representative from the many scientists who are critical of the ICNIRP standpoint.

Excerpts

As a general rule ICNIRP, WHO, SCENIHR and SSM have for many years dismissed available studies showing harmful effects from non-thermal RF exposure and have based their conclusions mainly on studies showing no effects. Results showing risk are criticized, disregarded or not even cited while studies showing no risks are accepted as evidence of no risk in spite of severe methodological problems. Many statements by these agencies are misleading and not correct. They are easily rebutted by reading the relevant publications.... All these expert groups dominated by ICNIRP consequently reach similar conclusions that there are no health effects below ICNIRP guidelines. No representative from the scientific community that is of the opinion that there is increasing evidence of health risks below the ICNIRP guidelines, e.g. as expressed in the EMF Scientists Appeal [24], has ever been a member of the expert groups at the WHO, the EU, the SSM or ICNIRP. Certainly scientists who do not discount evidence of health effects from exposure to RF radiation that are observed at exposures below guideline levels should be represented....

ICNIRP is not representative of the scientific community since it does not include representatives from scientists that agree there is evidence of harmful effects at levels well below ICNIRPs limits although these scientists are in majority in the scientific community [24].

Conclusion

ICNIRP's conclusion [48] on cancer risks is: "In summary, no effects of radiofrequency EMFs on the induction or development of cancer have been substantiated." This conclusion is not correct and is contradicted by scientific evidence. Abundant and convincing evidence of

increased cancer risks and other negative health effects are today available. The ICNIRP 2020 guidelines allow exposure at levels known to be harmful. In the interest of public health, the ICNIRP 2020 guidelines should be immediately replaced by truly protective guidelines produced by independent scientists.

Open access paper:

<https://www.fortunejournals.com/abstract/aspects-on-the-international-commission-on-nonionizing-radiation-protection-icnirp-2020-guidelines-on-radiofrequency-radiation-2261.html>

THE MICROWAVE AUDITORY EFFECT

James C. Lin. The microwave auditory effect. IEEE Journal of Electromagnetics, RF, and Microwaves in Medicine and Biology, pp 1-13, Mar 1, 2021. Digital Object Identifier: 10.1109/JERM.2021.3062826.

Abstract

The microwave auditory effect has been widely recognized as one of the most interesting and significant biological phenomena from microwave exposure. The hearing of pulsed microwaves is a unique exception to sounds encountered in human auditory perception. The hearing of microwave pulses involves electromagnetic waves. This paper reviews the research on humans and animals leading to scientific documentation the absorption of a single microwave pulse impinging on the head may be perceived as an acoustic zip, click, or knocking sound. A train of microwave pulses may be sensed as a buzz, chirp, or tune by humans. It describes neurophysiological, psychophysical, and behavioral observations from laboratory studies involving humans and animals. Mechanistic investigations show that the microwave pulse, upon absorption by tissues in the head, launches a pressure wave that travels by bone conduction to the inner ear, where it activates the cochlear receptors via the same process involved for normal sound hearing. Depending on the impinging microwave pulse power, the level of induced sound pressure could be considerably above the threshold of perception to cause tissue injury. The microwave auditory effects and pressure waves could potentially render damage to brain tissues to cause lethal or nonlethal injuries.

<https://ieeexplore.ieee.org/document/9366412>

Take-Home Messages

- This paper describes neurophysiological, psychophysical, and behavioral observations from laboratory studies involving human and animal subjects.
- Absorption of a single microwave pulse impinging on the head may be perceived as an acoustic zip, click, or knocking sound.
- A train of microwave pulses may be sensed as an audible buzz, chirp, or tune by humans.
- Mechanistic studies show absorption of microwave pulses by soft tissues in the head launches a thermoelastic pressure wave that travels in the brain
- Depending on the power of the impinging microwave pulses, the level of induced sound pressure could be considerably above the threshold of auditory perception.
- The microwave auditory effects and associated pressure waves could potentially render damage to brain tissues to cause lethal or nonlethal injury.

VIII. SUMMARY

The microwave auditory effect occurs from miniscule but rapid (microsecond) rise of temperature (10⁻⁶ degrees C) in the brain from absorption of pulsed microwave radiation. The sudden rise in temperature creates thermoelastic expansion of the brain matter, which can launch a pressure wave that propagates through the head and is detected by the sensory hair cells in the cochlea. The nerve signal is then relayed to the central auditory system for perception and recognition.

The preceding sections document that an audible sound originates from within the head when human subjects are exposed to pulsed microwave radiation. The auditory detection of pulsed microwaves in laboratory animals has been confirmed both in behavioral and neurophysiological studies. The site of microwave-to-sound conversion is shown to be in the brain tissue. The primary mechanism of interaction is microwave pulse-induced thermoelastic expansion of brain matter.

Depending on the power of the impinging microwave pulses, the levels of induced sound pressure in the brain could be considerably above the threshold of auditory perception, so that they may approach or exceed levels of discomfort and cause brain tissue injury. A high-power microwave pulse-generated acoustic pressure wave initiated in the brain and reverberating inside the head could bolster the initial pressure, causing injury of brain matter. Thus, it is conceivable that the microwave auditory effect or the microwave pulse-induced pressure shock wave inside the head could become a potentially lethal or nonlethal weapon against animals and humans.

The unique character of microwave-induced acoustic wave in biological tissues has prompted the exploration of its potential for application in biomedical imaging [68,69]. The principle of operation of microwave thermoacoustic tomography (MTT) and some results have been available since the early 1980's [69,70]. It was conjectured that the potential contrast advantage of microwave imaging and the resolution advantage of ultrasonic imaging could combine to make MTT imaging of biological tissues a potentially useful dual modality for diagnostic imaging. For example, the wavelength in muscle for microwaves is 17.5 mm at 2450 MHz; for ultrasonic waves, the wavelength is a mere 0.5 mm at 3 MHz. The potential gain in spatial resolution is tremendous for tissue imaging compared with relying on using microwave radiation alone. The research initiated in the 1980's is being actively pursued in developing MTT imaging for medical diagnosis, especially for early detection of breast cancer. Indeed, currently, MTT is a subject of vigorous research both from a systems development perspective and as a dual imaging modality amenable to greater utility in a wide range of medical application scenarios [71–75].

MICROWAVE THERMOACOUSTIC TOMOGRAPHIC (MTT) IMAGING

James C. Lin. Microwave Thermoacoustic Tomographic (MTT) Imaging. *Physics in Medicine and Biology*. 66(10). May 12, 2021. <https://doi.org/10.1088/1361-6560/abf954>.

Abstract

Microwave thermoacoustic tomography (MTT) uses microwave-pulse-induced thermoelastic pressure waves to form planar or tomographic images. Since the generation and detection of thermoelastic pressure waves depends on dielectric permittivity, specific heat, thermal expansion, and acoustic properties of tissue, microwave thermoacoustic imaging possesses the characteristic features of a dual-modality imaging system. The unique attributes of the high contrast offered by microwave absorption and the fine spatial resolution furnished by ultrasound are being explored to provide a nonionizing and noninvasive imaging modality for characterization of tissues, especially for early detection of breast cancer. This paper reviews the research being conducted in developing MTT imaging for medical diagnosis. It discusses the science of thermoelastic wave generation and propagation in biological tissues, the design of prototype MTT systems, the reconstruction of tomographic images, and the application and accomplishment of prototype MTT systems in phantom models and experimental subjects.

<https://iopscience.iop.org/article/10.1088/1361-6560/abf954/pdf>

MOBILE PHONE USE AND TRENDS IN THE INCIDENCE OF CANCERS OF THE PAROTID AND OTHER SALIVARY GLANDS

Ken Karipidis, Rohan Mate, Masoumeh Sanagou, Chris Brzozek, David Urban, Mark Elwood. Mobile phone use and trends in the incidence of cancers of the parotid and other salivary glands. *Cancer Epidemiol*. 2021 May 18;73:101961. doi: 10.1016/j.canep.2021.101961.

Highlights

- Adult parotid gland cancer in Australia since 2006 has decreased in males and increased in females.
- The incidence of other salivary gland cancers has remained stable.
- Mobile phone use is unlikely related to the incidence of salivary gland cancers.
- The increase in female parotid gland cancer may be attributed to possible risk factors specific to this gender.

Abstract

Background: There has been a significant increase in the use of mobile phones over the last three decades and a possible association with head cancers has been suggested, including cancers of the parotid and other salivary glands. We examined the incidence time trends of parotid and other salivary gland cancers in Australia to ascertain the influence of increased mobile phone use.

Methods: Analyses of incidence time trends were carried out using Poisson regression to estimate the annual percentage change (APC) in the incidence of salivary gland cancers of all available national registration data from 1982 to 2016, as well as specific time periods (1982-1993, 1994-2005, 2006-2016) representing changes in the prevalence of mobile phone use.

Results: The incidence of parotid gland cancer was stable for the periods 1982-1993 and 1994-2005. During 2006-2016 there was a large decrease in parotid gland cancer for males (APC: -3.71, 95 %CI: -6.66 to -0.67) and a large increase in females (4.80, 1.77-7.91) for adults aged 20-59 years. The incidence for other salivary gland cancers was stable during all the periods.

Conclusions: The results do not indicate that mobile phone use increased the incidence of parotid or other salivary gland cancers. An increase in parotid gland cancer in females since 2006 may be attributed to other possible risk factors specific to this gender.

<https://pubmed.ncbi.nlm.nih.gov/34020314/>

Excerpts

Several epidemiological studies have generally not found an association between mobile phone use and parotid gland tumours [2]. However, a recent meta-analysis of three case-control studies reported a small increase in the risk [3]....

In conclusion, these results do not indicate that mobile phone use increased the incidence of parotid or other salivary gland cancers. An increase in cancer of the parotid gland in females since 2006 may be attributed to possible risk factors specific to this gender. Our study is an ecological observational investigation and its interpretation has to be cautious. These results highlight the need for further research on possible risk factors for parotid gland cancer. Better understanding of the aetiology of the disease may reveal opportunities for public health strategies to reduce the rising incidence in females.

My note: Although I agree that more research is needed, since the authors did not study mobile phone use, they have no basis for ruling this out as a possible risk factor for the increased incidence of parotid gland tumors observed in females.

CELL PHONE USE AND THE RISK OF GLIOMA: ARE CASE-CONTROL STUDY FINDINGS CONSISTENT WITH CANADIAN TIME TRENDS IN CANCER INCIDENCE?

Paul J Villeneuve, Franco Momoli, Marie-Élise Parent, Jack Siemiatycki, Michelle C Turner, Daniel Krewski. Environ Res. 2021 May 21;111283. doi: 10.1016/j.envres.2021.111283.

Highlights

- The possibility that the use of cell phones increases the risk of cancer remains controversial.
- We examined trends in cell phone use and incidence rates of glioma for Canada between 1992 and 2015.
- Age standardized incidence rates for glioma in Canada were relatively stable between 1992 and 2015.
- Canadian trends in glioma and cell phone use were not compatible with increased risks of glioma that have been reported in some previous case-control studies.

Abstract

Background: There remains controversy as to whether cell phones cause cancer. We evaluated whether temporal changes in cell phone use and the incidence of glioma in Canada were consistent with the hypothesis of an increased risk.

Design: We used data from the Canadian Cancer Registry to calculate annual incidence rates for glioma between 1992 and 2015. The annual number of new cell phone subscribers was determined using national industry statistics. The number of newly diagnosed gliomas was compared to the predicted number by applying risks from epidemiological studies to age-specific population estimates. Specifically, we calculated the "predicted" number of incident gliomas by determining the annual prevalence of cell phone users and years of use. These estimates were multiplied by the corresponding risk estimates to determine the predicted number of gliomas.

Results: The number of cellular subscriptions in Canada increased from nil in the early-1980s to approximately 29.5 million in 2015. In contrast, age-standardized glioma incidence rates remained stable between 1992 and 2015. The application of risk estimates from

- i) a recent pooled analyses of Swedish case-control studies,
- ii) the 13 country INTERPHONE study, and
- iii) the Canadian data from INTERPHONE overestimated the observed number of glioma cases diagnosed in Canada in 2015 by 49%, 85%, and 63%, respectively.

Interpretation: Predictions of glioma incidence counts using estimates of the relative risk of glioma due to cell phone use from case-control studies over-estimated the incidence rates of glioma in Canada. The absence of an elevation in incidence rates of glioma in conjunction with marked increases in cell phone use suggests that there may not be a causal link between cellphones and glioma. <https://pubmed.ncbi.nlm.nih.gov/34029549/>

ELECTROMAGNETIC RADIATION REDUCTION USING NOVEL METAMATERIAL FOR CELLULAR APPLICATIONS

Ahmed Mahfuz Tamim, Mohammad Rashed Iqbal Faruque, Mayeen Uddin Khandaker, Mohammad Tariqul Islam, David Andrew Bradley. Radiation Physics and Chemistry, 178,108976. 2021. <https://doi.org/10.1016/j.radphyschem.2020.108976>.

Highlights

- A novel ELC resonator-based metamaterial is developed to reduce the specific absorption rate (SAR) on the mobile phone.
- Peak SAR investigated for three categories of modern smartphone usages- voice calling, messaging, and video calling.
- The proposed metamaterial can decrease a significant amount of SAR.
- Thus, the metamaterial can be helpful to protect the human head from harmful radiation.

Abstract

Excessive exposure to radiation has an adverse impact on human health, as an increase in body temperature may damage human organs or tissues, including the brain, eyes, and skin. Hence, this study assessed the effect of overexposure of radiation on the human head by analysing specific absorption rate (SAR) and reduction of SAR through the use of novel metamaterial (MTM). The SAR reduction was performed for GSM 900 MHz and 1800 MHz bands. A high-frequency electromagnetic simulator was employed throughout this study. The SAR investigation was performed on the head model for three categories of usage, namely voice calling, messaging, and video calling. This study looked into the impact on SAR of various

free space distances between mobile phone and head model. A novel electric field driven LC (ELC) resonator-based MTM was used to reduce SAR. Based on the properties of the tissue, the rate of tissue absorption escalated with an increase in radiated power, especially when the distance between head and mobile phone decreased. The study outcomes signified that MTM could decrease a significant amount of SAR. This is beneficial to protect the human body from harmful radiation, wherein the distance from the device in biological effect should be maintained.

Excerpts

Even if the tissue temperature is too little even to consider causing any harm, it is associated with electric fields that may harm the biological cells by an unknown non-thermal mechanism. Table 1 lists some biological criticalness due to absorbed radiation, as reported by the World Health Organisation (WHO) (Staebler, 2017)....

In this era of 4G, 5G, and 6G, temperature and SAR values of the brain, the eyes, and the skin tissues have been greatly affected by the induced heating of 4G or 5G mobile phone radiation (Christopher et al., 2020b). Based on several techniques of SAR reduction of mobile phones (Hossain et al., 2014; Rosaline and Raghavan, 2016; Lee and Lee, 2017; Kwak et al., 2011; Faruque and Islam, 2013), using metamaterials as the middle layer appears to be an effective approach. Metamaterials (MTMs) refer to artificial metal-dielectric composite structure materials that gain its electrical properties from its structure, instead of inheriting them directly from the materials it is composed of. This material has some unique properties that are absent in naturally formed substances. These MTMs are assembled from individual elements that are significantly smaller in typical size than the electromagnetic waves working wavelength (Smith et al., 2004; Kadic et al., 2013). The recent MTM studies not only focus on the varied wideband operations (Hasan et al., 2016; Islam et al., 2019), but also look into more superior applications, including MTM antenna (Hasan et al., 2018), cloaking (Islam et al., 2016), MTM absorber (Hasan et al., 2017a), and filter (Alam et al., 2019) etc....

Conclusion

This paper presents a detailed numerical investigation of the absorbed electromagnetic radiation by the head from a modern smartphone for three usages, namely voice calling, messaging, and video calling. For SAR calculation, a novel PIFA [planar inverted-F] antenna was operated at GSM 900 MHz and 1800 MHz bands. Next, an ELC resonator-based MTM was developed and placed in the middle layer between the radiation patch and the substrate of the PIFA antenna to reduce the radiation exposure of the antenna, wherein the SAR decreased for all configurations stated in this study. It was observed that the radiation emitted from voice calling, messaging, and video calling applications had affected the inner ear, chin, and nose the most, respectively. The highest radiation absorbed by the head is when one uses the mobile phone for voice calling application. The absorbed radiation increased the temperature substantially, thus harming the biological tissues of a human. The distances between the smartphone and the different organs of the head for varying usages had been investigated as well. This highlights the fact that safety precautions should be adopted to stay safe from modern smartphone radiation.
<https://www.sciencedirect.com/science/article/pii/S0969806X20302826>

EFFECTS OF ELECTROMAGNETIC FIELDS ON NEURONAL ION CHANNELS: A SYSTEMATIC REVIEW

Federico Bertagna, Rebecca Lewis, S Ravi P Silva, Johnjoe McFadden, Kamalan Jeevaratnam. Effects of electromagnetic fields on neuronal ion channels: a systematic review. *Ann N Y Acad Sci.* 2021 May 4. doi: 10.1111/nyas.14597.

Abstract

Many aspects of chemistry and biology are mediated by electromagnetic field (EMF) interactions. The central nervous system (CNS) is particularly sensitive to EMF stimuli. Studies have explored the direct effect of different EMFs on the electrical properties of neurons in the last two decades, particularly focusing on the role of voltage-gated ion channels (VGCs). This work aims to systematically review published evidence in the last two decades detailing the effects of EMFs on neuronal ion channels as per the PRISM guidelines. Following a predetermined exclusion and inclusion criteria, 22 papers were included after searches on three online databases. Changes in calcium homeostasis, attributable to the voltage-gated calcium channels, were found to be the most commonly reported result of EMF exposure. EMF effects on the neuronal landscape appear to be diverse and greatly dependent on parameters, such as the field's frequency, exposure time, and intrinsic properties of the irradiated tissue, such as the expression of VGCs. Here, we systematically clarify how neuronal ion channels are particularly affected and differentially modulated by EMFs at multiple levels, such as gating dynamics, ion conductance, concentration in the membrane, and gene and protein expression. Ion channels represent a major transducer for EMF-related effects on the CNS. <https://pubmed.ncbi.nlm.nih.gov/33945157/>

Excerpts

Different effects for different fields - An extensive literature exists reporting a myriad of EMF exposure-related effects on many biological processes, ranging from cell differentiation, survival, and changes in gene expression^{96, 97} to effects on cell membranes and signal transduction pathways.³⁰ However, many other studies indicated the absence of significant effects elicited by these fields.^{59, 61} A possible explanation for the different effects reported could be related to the fact that the way in which EMFs interact with the body depends on what combination of frequencies are used and the related wavelengths.

It is well known that the effects of exposure to EMFs differ significantly based on the exposure intensities and the exposure time,^{1, 98} and because of this any reasonable comparison must be made between groups having the same experimental conditions. In this study, we found different types of EMFs employed, although the two most commonly represented categories were ELF-EMFs and RF-EMFs, in line with the well-documented biological relevance of these fields.^{94, 99, 100} It is important to point out that the effects exerted by these two types of exposures are not equal due to the intrinsic electrical properties of the neuronal membrane. For instance, electrical phenomena involving a redistribution of charges in the membrane subsequent to EMF exposure, such as counterion polarization and phospholipid reorientation, are not likely to occur in RF-EMF exposure, due to the high inertia of charged particles at this high frequency.¹⁰¹

Moreover, pulsed EMFs are often reported to be more active relative to static EMFs, which are characterized by a continuous electromagnetic wave to which the cell could be more

adapted,¹⁰² and they could affect the gating properties of VGCs since these proteins are intrinsically sensitive to minimal electrical variations. ¹⁰⁰ Likewise, the effect of SMFs could similarly influence VGCs through a deformation of the membrane involving a reorientation of the phospholipid bilayer, as suggested by Rosen's study. ¹⁰³ Indeed, both of the studies reviewed here that focused on SMF effects reported effects on VGCs, specifically on the gating dynamics of VGCCs⁴⁷ and the inactivation dynamics of VGPCs. ⁴⁶

Lastly, it is worth mentioning that the frequency-related impact of the various type of EMFs has not been totally clarified, and theories exist suggesting that only specific frequencies would relevantly interact with the cell. ^{104- 106} However, although many different types of fields have been used, the frequencies used were similar (specifically 50 Hz for ELF-EMFs and 835 and 900 Hz for RF-EMFs).

Conclusion

The studies reviewed here show VGCs as an important transducer of the effect of EMFs in neurons, and the central role played by these proteins in the regulation of important biological processes, central in the regulation of brain physiology, sheds a light on the influence that modern exposure to EMFs could have on human health. While a diverse range of biological systems were used, cell lines were the preferred option, and VGCCs were the most studied ion channels, in line with their central role in the regulation of many physiological processes in neurons. However, many other VGCs have been shown to be affected by EMFs and the results are often conflicting. In spite of the controversy, this systematic review reports significant correlation between EMFs and multiple changes in the electrophysiological properties of diverse neuronal tissues, and these results, if interpreted well, could pave the way to a new understanding of the relationship between electromagnetic stimulation and brain functions. In conclusion, we systematically demonstrate how the complex effects of EMFs in neuronal ion channels are exerted at multiple levels and how their significance in the alteration of neuronal functions is strictly dependent on different parameters relative to the type of field used and the studied cell or tissue. Improved experimental reproducibility will be key to any advances in this field, and the development of new experimental procedures capable of measuring the small but profound way in which certain types of EMF exposure seem to affect our brain might help us to establish whether it is harmful and its therapeutic potential. We hope this work will help in improving our knowledge about the molecular dynamics of neuronal VGCs, which will be key both for any progress in the treatment of neurodegenerative diseases and for an advancement in the general understanding of the relationship between technological progress and cellular dynamics.

EXPOSURE OF SOUTH KOREAN POPULATION TO 5G MOBILE PHONE NETWORKS (3.4-3.8 GHZ)

Brahim Selmaoui, Paul Mazet, Pierre-Baptiste Petit, Kihwea Kim, Donggeun Choi, René de Seze. 2021 May 16. doi: 10.1002/bem.22345.

Abstract

As industrialized countries race to install and deploy 5G networks, some countries have taken the lead and already have operational 5G networks in place. South Korea is among these. In this study, we measured exposure to electromagnetic fields in South Korea to evaluate the

relative contribution of 5G as compared with other frequencies such as 2G, 3G, and 4G. Results show that the emission of 5G contributes about 15% to total telecommunications emissions. The highest levels were observed in the vicinity of 5G antennas and remain below the limits set by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). <https://pubmed.ncbi.nlm.nih.gov/33998007>

Excerpts

The measurements were carried out at the end of November 2019 in collaboration with the Korean institute in charge of telecommunications regulation—the National Radio Research Agency (NRRA)—6 months after the network was opened to public customers. This collaboration provided early insight into the organization of the 5G service, distinctions between the antennas of the various Korean operators, and information on the distribution of antennas in the country....

The average levels of 5G TDD LB varied with both the type of environment and the area and ranged from 20 to 560 mV/m with the ExpoM-RF dosimeter and from 20 to 70 mV/m with the EME Spy 200 dosimeter....

The maximum recorded EMF for 5G was 130 mV/m....

It appears that the contribution of 5G-NR to the overall EMF exposure is small in comparison with other frequencies. Indeed, the measurements carried out in Seoul and in different geographical areas of Naju showed that this contribution is less than 15% of the total exposure in rural and urban zones, and results were similar outside the city. The 5G exposure during the urban trip (the trip inside the city) contributed about 15% of the total exposure.

Our results show that 5G exposure levels were under the exposure limit set by the ICNIRP [International Commission on Non-Ionizing Radiation Protection [ICNIRP], 1998, 2020]. Indeed, for 5G, the threshold is 39 V/m at 700 MHz and 61 V/m at 3.5 and 26 GHz. Our data show that the maximum exposure measured on the rooftop in the vicinity of the base station was still below the threshold exposure limit set by ICNIRP, and at moving positions in the nearby street, it was far below this level (below 0.08 V/m). Our study is in line with other reports [Agence National des Fréquences [ANFR], 2020] that were published online....

Measurements over 24 h at a fixed point showed that the EMF emission from the base station is usually stable over time, at around 5 mV/m, except during rush hours, which are concentrated in the morning (around 8–9 am), where exposure increases up to 130 mV/m, and in the evening (around 6–8 pm), which shows only a two- fold increase compared to the median day and nighttime levels....

Measurements with a field meter in the vicinity of the 5G-NR base station were found to be 12 V/m (15 m from the antenna) with its baseline power. This value increased to 21 V/m when the antenna was at its maximum power at the same distance....

In our study, the results with the two different dosimeters, ExpoM-RF and EME Spy 200, present some variability. This is mainly attributed to the different sensitivity thresholds of

each device when the ambient exposure is close to the measurement threshold. However, at significant exposure levels, the measured values converge....

This study, in line with other reports [ANFR, 2020; Ofcom report, 2021; Telstra report, 2021] published online, is a preliminary study because EMF measurements were carried out only 6 months after the opening of the network to the Korean public. It is likely that the 5G network was not being used to its maximum and the number of subscribers was relatively low, due to the tendency of young people to use Wi-Fi spots for free. It is necessary to continue conducting EMF measurements to monitor the overall trends in the long term.

Nevertheless, it appears from this study that exposure levels from 5G were in a low range. Moreover, the fact that the radio signal beam mainly focuses on the end-users will contribute to reducing exposure in unnecessary areas. This work is part of a measurement campaign in South Korea, whose final report was published in 2019 [French Expertise Report, 2019].

IN SITU ASSESSMENT OF 5G NR MASSIVE MIMO BASE STATION EXPOSURE IN A COMMERCIAL NETWORK IN BERN, SWITZERLAND

Sam Aerts, Kenneth Deprez, Davide Colombi, Matthias Van den Bossche, Leen Verloock, Luc Martens, Christer Törnevik, Wout Joseph. In Situ Assessment of 5G NR Massive MIMO Base Station Exposure in a Commercial Network in Bern, Switzerland. *Appl. Sci.* 11(8): 3592. 2021. <https://doi.org/10.3390/app11083592>.

Abstract

This paper describes the assessment of radiofrequency (RF) electromagnetic field (EMF) exposure from fifth generation (5G) new radio (NR) base stations in a commercial NR network in Bern, Switzerland. During the measurement campaign, four base station sites were investigated and the exposure induced by the NR massive multiple-input-multiple-output (MaMIMO) antennas was assessed at 22 positions, at distances from the base station between 30 m and 410 m. The NR base stations operated at 3.6 GHz and used codebook-based beamforming. While the actual field levels without inducing downlink traffic were very low (<0.05 V/m) due to a low traffic load and low antenna input powers of up to 8 W, setting up a maximum downlink traffic stream towards user equipment resulted in a time-averaged exposure level of up to 0.4 V/m, whereas the maximum extrapolated exposure level reached 0.6 V/m. Extrapolated to an antenna input power of 200 W, values of 4.3 V/m and 4.9 V/m, respectively, were obtained, which amount to 0.5–0.6% of the reference level recommended by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). In Bern, it was found that the impact of the NR network on the total environmental RF exposure was very limited; with maximum downlink, it contributed 2% on average. Finally, it was also concluded that extrapolation to the maximum exposure level can be done without prior knowledge of the radiation patterns, directly based on the measurement of the Physical Downlink Shared Channel (PDSCH) resource elements.

Conclusions

To the authors' knowledge, this study provides the first assessment of the range of actual and maximum exposure levels in a 5G NR commercial network. It was found that the impact of the investigated network on the total environmental RF-EMF exposure was small, only a few percent of the total RF-EMF exposure even in the case of 100% induced traffic. Moreover, an

extrapolation method was demonstrated for which no prior information from the network provider or radio equipment manufacturer is necessary, validated using the actual antenna radiation patterns.

In the (near) future, as 5G technologies evolves (e.g., when introducing reciprocity-based beamforming and other advanced MaMIMO techniques), some aspects of the presented methodology, such as the position of the UE relative to the base station and measurement probe, the averaging time to assess E_{avg} , and the use of the UE to stimulate the maximum exposure scenario may also have to be amended. In addition, we are confident that the described procedure is valid also at frequencies in Frequency Range 2 (FR2), above 24 GHz ('mmWaves'), provided that the measurement settings are adjusted to account for wider channel bandwidths as well as larger SCS. However, a comprehensive in situ validation study remains essential.

Open access paper: <https://www.mdpi.com/2076-3417/11/8/3592>

INDIVIDUAL EXPOSURE TO ENVIRONMENTAL RADIOFREQUENCY ELECTROMAGNETIC FIELDS IN HOSPITALIZED PRETERM NEONATES

Dimitri Besset, Brahim Selmaoui, Stéphane Delanaud, René de Seze, André Leke, Erwan Stéphan-Blanchard. *Bioelectromagnetics*. 2021 May 16. doi: 10.1002/bem.22349.

No abstract. <https://pubmed.ncbi.nlm.nih.gov/33998013/>

Excerpts

RF-EMF levels in the different frequency bands over the 3-week recording period for all infants are shown in Figure 1. For the median level, the total exposure (0.03 ± 0.01 V/m) was mainly due to the FM/TV (0.01 ± 0.01 V/m), mobile phone (UL) (0.01 ± 0.00 V/m), and DECT (0.01 ± 0.01 V/m). With regard to the P99 level, the frequency bands' relative contribution to total exposure (0.24 ± 0.11 V/m) differed; the mobile phone UL signal (0.23 ± 0.11 V/m) was higher than the DECT signal (0.044 ± 0.020 V/m). For both the median and P99 levels, all other bands were below the exposimeter's limit of detection....

The present study is the first to (i) show that preterm neonates are chronically exposed to low environmental RF-EMF levels while in the NICU; (ii) describe the exposure's characteristics; and (iii) provide objective, quantitative exposure data. On average, the RF-EMF levels to which preterm neonates were exposed were much lower than the currently recommended 50 V/m threshold [International Commission on Non-Ionizing Radiation Protection ICNIRP, 2020] but higher than those found in the general environment of the unit [Besset et al., 2020]. DECT, mobile phone (UL), and FM/TV were the main contributors to this chronic exposure. Even though RF-EMF levels were recorded every 2 min (meaning that many peak and minimal values were probably missed), 94.1% of the infants were nevertheless transiently exposed to levels above 1 V/m ($0.045 \pm 0.074\%$ of all RF-EMF values, for around 41 s per day). These peak exposures were mainly due to mobile phone UL signals. Exposure to DECT and mobile phone frequency bands suggests that neonate exposure is mainly due to human activities around them (e.g., parent attentiveness, medical care) and explains why individual exposure levels were higher than those of the NICU general environment.

These exposure profiles do not place preterm neonates at risk. However, some attention to the RF-EMF induced by human activities around the infants is needed because of their high sensitivity to the environment, physiological limitations, and increased body absorption of RF-EMF energy [Peyman, 2011] during a critical maturation period. Due to the small sample size and the specificity of the hospital environment in this study, further exposure investigations are required to confirm the present data.

ARTIFICIAL EMG (ELECTROMYOGRAM) BY WLAN EXPOSURE

Lebrecht von Klitzing. Artificial EMG (electromyogram) by WLAN exposure. Journal of Biostatistics and Biometric Applications. 6(1): 2021.

Abstract

WLAN (wireless local area network) is used as an important worldwide communication-technique. By this, always there is an electromagnetic field exposure. In contrast to the ICNIRP-safety guidelines, whereby no bioeffect is possible by these low-energetic electromagnetic fields, we found artificial signals in the nervous system in dependence on WLAN-exposure.

Open access paper: <http://www.annexpublishers.com/articles/JBIA/6101-Artificial-EMG-by-WLAN-Exposure.pdf>

NEW ICEMAN PROJECT SEEKS ANSWERS TO FIGHTER PILOT DISORIENTATION [HEALTH MATTERS]

James C. Lin. New ICEMAN Project Seeks Answers to Fighter Pilot Disorientation [Health Matters], IEEE Microwave Magazine. 2021; 22 (4):13-15. DOI: 10.1109/MMM.2020.3048210.

Excerpt

In summary, the ambient RF–EM field levels in a typical fighter/attack aircraft’s cockpit are now unclear. Quantitative surveys and measurements are necessary to allow the proper assessment of the RF–EM field’s potential effects on pilots’ brain activity, neurophysiology, and behavioral responses. Noticeably, fighter cockpits are subjected to strong impinging RF and/or microwave radar pulses under some operational conditions.

There are two pulsed microwave-induced auditory responses in humans and mammals when the head is exposed to high-power microwave pulses that could impact a pilot’s cognitive performance and response. Both are related to microwave pulse-induced acoustic pressure waves in the head: the microwave auditory effect and the acoustically induced startle reflex and motor reaction from a sudden, unexpected, intense auditory stimulus. The startle reaction from a sudden, unexpected auditory stimulus may cause the pilot to experience SD [spatial disorientation], during which one’s perception of aircraft position, motion, altitude, or attitude does not correspond to actuality.

Note that this supposition is derived from theoretic treatment along with available but limited experimental evidence. The kinds of confirmational studies that would be useful are neurophysiological and psychophysical investigations of pulsed microwave-exposed animals, including observations of the subjects’ behavioral and performance responses.

<https://ieeexplore.ieee.org/document/9366591>

METHODOLOGY FOR DETERMINING THE THRESHOLD DISTANCE FOR ESTIMATING THE MAIN EM EXPOSURE CONTRIBUTION IN WLAN

Marta Fernández, David Guerra. Methodology for determining the threshold distance for estimating the main EM exposure contribution in WLAN. Engineering Science and Technology, an International Journal, 2021. <https://doi.org/10.1016/j.jestch.2021.05.001>.

Abstract

The location of radiation sources in wireless networks is a key factor to characterize their contribution to electromagnetic exposure levels in order to deploy future networks that account for minimizing electromagnetic field levels. In relation to wireless local area networks, considering that mobile communication devices comply with the SAR (Specific Absorption Rate) limits imposed by the international standardization organizations for preserving human health, the interest is nowadays focused on the signal levels coming from the WiFi access points. This paper presents a methodology to determine the threshold distance at which the field strength levels from the AP are negligible in comparison with the radiation generated by a user equipment. The theoretical concepts, which can be applied to other technologies, were implemented by means of simulations and experimental measurements. For the simulations, actual WiFi antennas were modelled. Experimental measurements completed the results obtained in simulations, resulting in a greater number of real situations. Results showed that the threshold distance depends on the WiFi standard employed by the devices connected to the network.

Conclusions

The contribution each of the EM [electromagnetic] sources to the total exposure in wireless networks is an essential parameter for characterizing EM exposure and develop network deployments that minimize EMfield levels. This manuscript presents a methodology for determining the distance at which the EM field levels transmitted by an AP [access point] can be negligible in comparison with those transmitted by a UE [user equipment] in order to simplify the assessment of the total exposure. This can also give information about the positions in which measurements should be taken in future measurement campaigns. In the proposed methodology, the threshold distances were calculated for a specific distance of the UE. In this work the threshold distance was determined by means of a rigorous methodology assuming the worst case scenario, that is, full activity of the WiFi network, 90th percentiles, and a distance of 20 cm between the UE and the reception point. This distance allows us to perform measurements in the far field region and, it can represent scenarios in which a smartphone is placed on a desk at 20 cm from the user, or scenarios in which the exposure is due to the UE of another person. Moreover, it is consistent with the definition of portable devices, since these devices are defined as transmitting devices designed to be used so that the radiating structures of the devices are within 20 cm of the body of the user [11]. Thus, in the selected conditions the results provide the largest threshold distance when the own user or the person next to him/her is exposed to radiation coming from the UE. However, for people who are at greater distances from the exposure source, the threshold distances would be longer. This way, under free space propagation conditions, threshold distances ranging from 1.64 m and 4.99 m were obtained for the cases of hotspots transmitting 160 mW with bandwidths of 80 MHz and 40 MHz, respectively. These distances can increase in multipath environments as demonstrated in Section 5.4, where a maximum threshold distance of 5.92 m was obtained. In all the cases, if the UE was placed closer to the human

body, the threshold distance would be lower. This conclusion may lead to a misconception that needs to be clarified. The results point out that the higher the UE power is, the lower threshold distance to the AP is allowed. However, regarding safety as the main scope of this type of research, such threshold situations characterized by maximum power UEs and minimum distances to APs should be avoided by all means. That is, the higher the threshold distance is, the better the exposure scenario will be.

Experimental results showed that the threshold distance depends on the bandwidth employed by the communication link, which is related to the WiFi standard employed by the AP and UE. Furthermore, measurements in the whole 5 GHz WiFi band were carried out in a university environment and it was proved that WiFi signals coming from further APs have lower influence than at 2.4 GHz due to the higher attenuations suffered at higher frequencies. Finally, the threshold distances were also calculated in multipath environments. The proposed methodology ensures that WiFi APs will not be a matter of concern if the user is located at larger distances than the threshold distance. On the contrary, measurements of the radiation coming from the hotspot would be needed at shorter distances, in order to check the compliance with the reference levels. Finally, the duration of signal transmission may be of great interest when assessing EMF exposure, especially in places where people spend many hours such as schools, universities or offices. In this regard, WiFi exposure levels at 2.4 GHz were evaluated by means of 24-hour measurements in a university environment in [14]. As can be observed in Fig. 4 of [14], WiFi signals transmission increased in working days compared to weekends. But even during working hours, WiFi activity and therefore AP and UE transmissions varies from one location to another inside the same building.

<https://www.sciencedirect.com/science/article/pii/S2215098621001099>

EFFECTS OF 1.5 AND 4.3 GHZ MICROWAVE RADIATION ON COGNITIVE FUNCTION AND HIPPOCAMPAL TISSUE STRUCTURE IN WISTAR RATS

Ruiqing Zhu, Hui Wang, Xinping Xu, Li Zhao, Jing Zhang, Ji Dong, Binwei Yao, Haoyu Wang, Hongmei Zhou, Yabing Gao, Ruiyun Peng. Sci Rep. 2021 May 12;11(1):10061. doi: 10.1038/s41598-021-89348-4.

Abstract

Previous studies have shown that single-frequency microwave radiation can lead to cognitive decline in rats. However, few studies have focused on the combined effects of irradiation with different frequencies of microwaves. Our research aimed to investigate the effects of 1.5 GHz and 4.3 GHz microwave radiation, singly and in combination, on cognitive function and hippocampal tissue structure in rats. A total of 140 male Wistar rats were randomly divided into 4 groups: the S group (sham radiation group), L10 group (10 mW/cm² 1.5 GHz group), C10 group (10 mW/cm² 4.3 GHz band group) and LC10 group (10 mW/cm² 1.5 and 4.3 GHz multi-frequency radiation group). For 1-28 days after microwave radiation, we analyzed the average escape latency for the Morris water maze task, electroencephalograms, change in hippocampal tissue structure and ultrastructure, content of the Nissl body in the hippocampus, and activities of lactate dehydrogenase and succinate dehydrogenase. Compared to the S group, all exposure groups showed varying degrees of learning and memory decline and hippocampal structural damage. The results showed that 1.5 GHz and 4.3 GHz microwave radiation was able to induce cognitive impairment and hippocampal

tissue damage in rats and combined radiation with both frequencies caused more serious injuries, but none of these damaging effects varied with microwave frequency.

Excerpts

The microwave pulses were delivered at 200 pps, with a pulse width of 500 ns. The peak field power densities tested with a calibrated detector and an oscilloscope for the exposure groups was 100 W/cm². The average field power densities were calculated to be 10 mW/cm² (Fig. 1A and Supplementary Fig. 1A)....

[According to Table 1, the SAR values were 3.7 W/kg for the L10 group and 3.3 W/kg for the C10 group.] Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8115682/>

PROTECTIVE EFFECTS OF SELENIUM ON ELECTROMAGNETIC FIELD-INDUCED APOPTOSIS, AROMATASE P450 ACTIVITY, AND LEPTIN RECEPTOR EXPRESSION IN RAT TESTIS

Sareh Khoshbakht, Fatemeh Motejaded, Sareh Karimi, Narjes Jalilvand, Alireza Ebrahimzadeh-Bideskan. Iran J Basic Med Sci. 2021 Mar;24(3):322-330. doi:10.22038/ijbms.2021 45358

Abstract

Objectives: Electromagnetic field (EMF) emitted by mobiles may affect the male reproductive system. Selenium, as an antioxidant, may protect against electromagnetic field-induced tissue damage. This study aimed to investigate the effects of selenium on rat testis exposed to electromagnetic fields.

Materials and methods: Twenty-four male Wistar rats were divided into four groups, namely EM group (2100 MHz), EM/SE group (2100 MHz + selenium (0.2 mg/kg), SE group (selenium 0.2 mg/kg), CONT (control group). Serum LH, FSH, testosterone, leptin and aromatase levels, testis weight and volume index, sperm parameters (count and abnormal percent), seminiferous tubule diameters, germinal epithelia thickness, immunoreactivity of leptin receptor and caspase-3 (for apoptotic cells in germinal epithelium) were investigated.

Results: Our results showed that serum LH, FSH, GnRH, testosterone level, sperm count, germinal epithelium thickness, and seminiferous tubule diameter were significantly declined in the EM group compared with the CONT group ($P<0.05$). However, in the EM group, the serum leptin level, sperm abnormality, aromatase enzyme level, apoptotic cells, and leptin receptor were increased compared with the CONT group ($P<0.05$). Furthermore, an increase in sperm count, germinal epithelium thickness, seminiferous diameters, serum LH, FSH, and GnRH, and testosterone levels, and a significant decrease in sperm abnormality, leptin receptor and apoptotic cells in the EM/SE group compared with the EM group were also observed ($P<0.05$).

Conclusion: This study showed that electromagnetic radiation may have detrimental impacts on the male reproductive system, which can be prevented by use of selenium.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8087852/>

CHANGES IN GROWTH KINETIC PARAMETERS, MORPHOLOGY AND MITOTIC ACTIVITY OF YEASTS CANDIDA GUILLIERMONDII EXPOSED TO THE LOW-INTENSITY WAVES OF 51.8-GHZ FREQUENCY

Seda Marutyan, Syuzan Marutyan, Liparit Navasardyan, Karlen Hovnanyan, Armen Trchounian. Changes in growth kinetic parameters, morphology and mitotic activity of yeasts *Candida guilliermondii* exposed to the low- intensity waves of 51.8-GHz frequency. Arch Microbiol. 2021 May 3. doi: 10.1007/s00203-021-02336-0.

Abstract

Under the influence of electromagnetic waves of millimeter range with the frequency of 51.8 GHz, changes in the morphology, growth parameters and mitotic activity of yeasts *C. guilliermondii* NP-4 are revealed. Filamentous and giant cells appeared in a population of exposed yeasts. The sigmoid shape of the growth curve remained but the lag phase duration was increased by 2 h in comparison with non-exposed yeasts; accordingly, the log and stationary phases followed 2 h later. The specific growth rate in the log growth phase and colony- forming ability of exposed yeasts was decreased. It is suggested that yeasts have some response mechanisms to 51.8-GHz frequency electromagnetic waves. The results can be used to understand the response mechanisms of microorganisms to non-ionizing radiation, as well as to develop approaches to protect living organisms from it. The effect of electromagnetic waves of 51.8-GHz frequency to suppress yeasts can be applied in biotechnology and medicine. <https://pubmed.ncbi.nlm.nih.gov/33938972/>

Excerpt

EHF EMR ($\nu = 51.8$ GHz) was used in the amplitude modulation mode with a frequency of 1 Hz (signal frequency stability was 0.05%). A low-intensity EMR (flow power of 0.06 mW cm^{-2}) was applied (Hovnanyan et al. 2017; Soghomonyan and Trchounian 2018; Torgomyan et al. 2012)

Summarizing a comparative study of the effects of low-intensity electromagnetic waves with a frequency of 51.8 GHz on the yeast *C. guilliermondii* NP-4, it can be concluded that electromagnetic waves of the frequency of 51.8 GHz lead to changes in the morphology of yeast cells, to decreases in their growth, mitotic activity and colony-forming ability. These results increase interest in further studying the effects of EMR of other EHF on yeasts and other cells.

Based on the fact that yeasts are a convenient model for studying higher eukaryotes, it is appropriate to assume that electromagnetic waves with a frequency of 51.8 GHz will have the same overwhelming effect on the growth of higher eukaryotes. From this point of view, based on our results, it can be assumed that abuse of cellular communications is dangerous not only for microorganisms, but also for higher eukaryotes, in particular for human health.

The results can be used to elucidate the response mechanisms of living organisms to radiation stress, as well as to develop approaches for protecting living organisms from non-ionizing radiation. In addition, they can be useful in biotechnology and medicine to suppress yeast, when pathogenic species cause various diseases.

COMPARISON OF EFFECTS OF HIGH- AND LOW-FREQUENCY ELECTROMAGNETIC FIELDS ON PROLIFERATION AND DIFFERENTIATION OF NEURAL STEM CELLS

Wenfang Bai, Meihui Li, Weicheng Xu, Mingsheng Zhang. Comparison of effects of high- and low-frequency electromagnetic fields on proliferation and differentiation of neural stem cells. *Neurosci Lett*. 2021 Jan 10;741:135463. doi: 10.1016/j.neulet.2020.135463.

Highlights

- Transcranial magnetic stimulation (TMS) has been widely applied to diagnose and treat intractable brain diseases.
- Both 50 Hz LF-EMF and HF-EMF can promote the proliferation of NSCs in vitro.
- LF-EMF can accelerate NSCs to differentiate into neurons.

Abstract

To compare the effects of high- (HF-EMF) and low-frequency electromagnetic fields (LF-EMF) on the proliferation and differentiation of neural stem cells (NSCs). NSCs were obtained from SD rat hippocampus and cultured in suspension and adherent differentiation media. NSCs were exposed to LF-EMF (5 m T, 50 Hz, 30 min daily), HF-EMF (maximum magnetic induction 2.5 T, 40 % MO, 50 Hz, 10 min daily) and no electromagnetic field. At 3 d, cell viability and quantity of NSCs in suspension were detected by CCK-8 assay and cell counting plate.

Immunofluorescence staining and qRT-PCR were performed to detect the percentage of Tuj-1 and GFAP-positive NSCs and the expression of Tuj-1 and GFAP mRNA. The P3 NSCs were positive with Nestin and induced NSCs expressed Tuj-1, GFAP and oligodendrocyte markers (MBP). CCK-8 assay and cell counting showed that the OD value and quantity of LF-EMF group were significantly higher than those in other two groups (both $P < 0.05$).

Compared with the control group, the OD value and quantity were significantly higher in the HF-EMF group ($P < 0.05$). Immunofluorescence staining and qRT-PCR revealed that the percentage of Tuj-1 positive cells and the expression of Tuj-1 mRNA of NSCs exposed to LF-EMF were the highest (both $P < 0.05$). The proportion of GFAP- positive NSCs and the expression of GFAP mRNA did not significantly differ among three groups (all $P > 0.05$).

Both 50 Hz LF-EMF and HF-EMF can promote the proliferation of NSCs in vitro and LF-EMF can accelerate NSCs to differentiate into neurons.

<https://pubmed.ncbi.nlm.nih.gov/33129846/>

BIOMOLECULAR RESPONSE TO HOUR-LONG ULTRALOW FIELD MICROWAVE RADIATION: AN EFFECTIVE COARSE-GRAINED MODEL SIMULATION

Anang Kumar Singh, P S Burada, Anushree Roy. *Phys Rev E*. 2021 Apr;103(4-1):042416. doi: 10.1103/PhysRevE.103.042416.

Abstract

Various electronic devices, which we commonly use, radiate microwaves. Such external perturbation influences the functionality of biomolecules. In an ultralow field, the cumulative response of a molecule is expected only over a time scale of hours. To study the structural dynamics of biomolecules over hours, we adopt a simple methodology for constructing the

coarse-grained structure of the protein molecule and solve the Langevin equation under different working potentials. In this approach, each amino acid residue of a biomolecule is mapped onto a number of beads, a few for the backbone, and few for the side chain, depending on the complexity of its chemical structure. We choose the force field in such a way that the dynamics of the protein molecule in the presence of ultralow radiation field of microvolt/nm could be followed over the time frame of 2h. We apply the model to describe a biomolecule, hen egg white lysozyme, and simulate its structural evolution under ultralow strength electromagnetic radiation. The simulation revealed the finer structural details, like the extent of exposure of bioactive residues and the state of the secondary structures of the molecule, further confirmed from spectroscopic measurements [details are available in Phys. Rev. E 97, 052416 (2018)10.1103/PhysRevE.97.052416 and briefly described here]. Though tested for a specific system, the model is quite general. We believe that it harnesses the potential in studying the structural dynamics of any biopolymer under external perturbation over an extended time scale. <https://pubmed.ncbi.nlm.nih.gov/34005990/>

Summary

This article aims to address the possible effect of an hour-long ultralow radiation field on a biosystem using the CG modeling technique. We obtain an optimized CG structure of a biopolymer, by considering the complexity of the chemical structure of its constituent amino acid residues. The chosen force field could reveal the dynamics of the molecule for hours. We demonstrate the efficacy of the model to study the overall topology and few finer structural details (e.g., dynamics of specific residues and secondary structures) of the HEWL molecule over an extended duration (~2 h) under ultralow field radiation (~10–7V/nm). It is computationally expensive to carry out all-atom quantum mechanical simulations over the above-mentioned time frame. Surely, the given model finds limitations (e.g., the explicit solvent effect has been neglected) and requires a higher level of optimization. Nonetheless, we believe that the approach, which we followed, is quite general and may pave a way to follow the structural dynamics of other biopolymers under an external perturbation for hours, if the solvent effect can be appreciably taken into account by the random noise in the beads' motion. It will also be interesting to test other more elegant CG models, mentioned in Sec. I of this article, to probe the effect of hour-long ultralow field radiation on biomolecules. From a biology viewpoint, we demonstrate the adverse effect of the radiation field on biomolecules, even when its strength is superlow.

EXPOSURE TO EXTREMELY LOW-FREQUENCY MAGNETIC FIELDS AND CHILDHOOD CANCER: A SYSTEMATIC REVIEW AND META- ANALYSIS

GyeongAe Seomun, Juneyoung Lee, Jinkyung Park. Exposure to extremely low-frequency magnetic fields and childhood cancer: A systematic review and meta-analysis. PLoS ONE 16(5): e0251628. May 14, 2021. doi:10.1371/journal.pone.0251628.

Abstract

Background Extremely low frequency magnetic fields (ELF-MFs) are classified as a possible carcinogenic factor (Group 2B). This study assessed the association between ELF-MFs and childhood cancer through a systematic review and meta-analysis.

Methods Three databases were searched in January 2020. We conducted a meta-analysis for the association between the ELF-MFs exposure level and childhood cancer.

Results A total of 33 studies were identified. Thirty studies with 186,223 participants were included in the meta-analysis. Children exposed to 0.2-, 0.3-, and 0.4- μ T ELF-MFs had a 1.26 (95% confidence interval [CI] 1.06–1.49), 1.22 (95% CI 0.93–1.61), and 1.72 (95% CI 1.25–2.35) times higher odds of childhood leukemia. In childhood brain tumors, children exposed to 0.2- μ T had a 0.95 (95% CI 0.59–1.56) times higher odds, and those exposed to 0.4- μ T ELF-MFs had a 1.25 (95% CI 0.93–1.61). Children exposed to 0.2- and 0.4- μ T ELF-MFs had a 1.10 (95% CI 0.70–1.75) and 2.01 (95% CI 0.89–4.52) times higher odds of any childhood cancers.

Conclusions Significant associations were observed between exposure to ELF-MFs and childhood leukemia. Furthermore, a possible dose-response effect was also observed.

Excerpts

The debate on the effect of electromagnetic fields (EMFs) on the human body still continues, and several studies have investigated the effect of magnetic fields that are not well shielded by objects [1–3]. The question of whether exposure to extremely low-frequency magnetic fields (ELF-MFs) from power transmission and distribution or the use of electrical appliances is associated with an increased risk of childhood cancer has engendered scientific debate [4–6]. In 2001, the ELF-MFs were classified by the International Agency for Research on Cancer (IARC) as possibly carcinogenic (Group 2B), based on the limited clinical evidence, inadequate experimental support, and the lack of plausible mechanisms at the exposure levels that were observed in epidemiological studies [7, 8]. This classification was endorsed by the subsequent weight of evidence assessments carried out by the World Health Organization (WHO) [8]. Subsequently, clinical evidence emerged from epidemiological studies on the etiology of childhood leukemia that indicated a weak association with ELF-MFs [9–12]....

Conclusions In this large pooled analysis of more than 36,000 children diagnosed with childhood leukemia, statistically significant associations were observed between exposure to ELF-MF and childhood leukemia. Furthermore, the intensity of the association between exposure to ELF-MFs and childhood leukemia was high, as indicated by the dose-response effect.

The risk of ELF-MFs, which have been classified as a possibly carcinogenic (Group 2B) factor based on limited evidence in humans, can be ascertained through precise evidence from the integrated results of this study.

Open access paper:

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0251628>

EXTREMELY LOW FREQUENCY ELECTROMAGNETIC EXPOSURE ASSESSMENT IN SCHOOLS: A STATISTICAL ANALYSIS OF URBAN AND SEMI-URBAN AREAS

Y Kiouvrekis, A Alexias, V Softa, M Alkhorayef, A Sulieman, C Tyrakis, C Kappas. Radiat Prot Dosimetry. 2021 May 28; doi: 10.1093/rpd/ncab076.

Abstract

The aim of this research paper is to estimate the mean value of extremely low frequency (ELF) exposure in schools in Greece. Detailed ELF measurements were conducted and analyzed by

the Weighted Peak Method (WPM), which estimates the overall contribution of electromagnetic waves from 1 Hz to 400 kHz, including their phases. A sample of 243 schools was sampled to calculate. The mean value of ELF magnetic fields (MFs) measured in these two groups comprising 243 schools was the principal focus of interest. ELF MF measurements taken in schools situated both far from and near ELF sources had mean and maximum values well below the current International Commission on Non-Ionizing Radiation Protection (ICNIRP) standard. The mean value of ELF MFs from all sources within the sampled schools in Greece was 0.21 μT . A statistically significant difference between the mean MFs measured in the two groups of schools was found, but the MFs in both groups were much lower than the ICNIRP standard. <https://pubmed.ncbi.nlm.nih.gov/34047347>

Excerpts

The median value of the MFs measured in the ELF range was 0.174 μT , which is 0.174% of the 2010 ICNRP standard (100 μT). In Figure 1, we present a typical example, and Table 1 shows the relative contribution of the harmonics....

Mean value is 0.23 μT for urban areas and 0.14 μT for semi-urban areas with s.d. of 0.23 and 0.13, respectively; the median is 0.18 for the urban area category and 0.11 for the semi-urban category, while the 25 and 75% percentiles are 0.12 μT and 0.25 μT for urban areas and 0.04 μT and 0.18 μT for the semi-urban areas, respectively (Figure 4). Similarly, concerning the WPM variable, the mean value is 1.33% for urban areas and 1.20% for semi-urban areas with s.d. of 0.18 and 0.13%, respectively; the median is 1.30% for the urban area category and 1.19% for the semi-urban category, while the 25 and 75% percentiles are 1.25 and 1.36% for urban areas and 1.10 and 1.26% for semi-urban areas, respectively (Figure 4).

Conclusions

ELF measurements (Max Peak, WPM) in schools situated both far from and near ELF sources allow for a comprehensive assessment of exposure in schools. Furthermore, statistical analysis shows that, on one hand, there is a statistically significant difference between schools in areas described as urban and semi-urban.

Despite the fact that there is a statistically significant difference between the levels of MFs measured in schools two separate areas, it is negligible with regard to the current standard by virtue of it being a difference by several orders of magnitude.

What is noteworthy is that the WPM values (6–500 Hz range) of the urban and semi-urban schools were found to be approximately an order of magnitude higher than that recorded in WPM frequency range (1 Hz–400 kHz). This is because in semi-urban and urban areas, the mode frequency is 50 Hz due to the power grid. While in terms of the MAX PEAK value, it also makes sense to be different and there is a statistically significant difference, as we are referring to an instantaneous value of each measurement point, which may have been affected but without affecting the final value of the point.

IDENTIFICATION OF ENVIRONMENTAL AND EXPERIMENTAL FACTORS INFLUENCING HUMAN PERCEPTION OF DC AND AC ELECTRIC FIELDS

Kathrin Jankowiak, Sarah Driessen, Andrea Kaifie, Simon Kimpeler, Thomas Krampert, Thomas Kraus, Dominik Stunder, Michael Kursawe. Identification of Environmental and Experimental Factors Influencing Human Perception of DC and AC Electric Fields. Bioelectromagnetics. 2021 May 11. doi: 10.1002/bem.22347.

Abstract

As part of the energy transition in Germany, high-voltage direct current (HVDC) lines producing DC electric fields (EF) are in planning. Since the human perception of DC EF was rarely investigated in the past, we aimed to identify environmental and experimental factors influencing the human perception of direct current (DC) EF, alternating current (AC) EF, and the co-exposure of DC EF and AC EF (hybrid EF) under whole-body exposure.

Additionally, first estimates of DC EF and AC EF perception thresholds as well as differences in human perception of DC EF and AC EF concerning the type of sensation experienced and the affected body part were evaluated. A highly sophisticated exposure lab was built to expose participants to various EF strengths and ask for their assessment concerning the presence of an EF. To estimate the individual perception thresholds of 11 participants, the signal detection theory as well as the single-interval-adjustment matrix procedure were applied. Relative humidity could be identified as an environmental factor influencing the perception of AC EF and DC EF in different ways. An appropriate ramp slope and an exposure duration for future studies could be elaborated. Additionally, perception thresholds were lower under hybrid EF exposure than under DC EF or AC EF exposure alone. Cutaneous sensations evoked under DC EF and AC EF exposure were individually different and attributed to various parts of the body. Several environmental and experimental factors influencing the human perception of EF could be identified and provide an essential basis for a large-scale study. <https://pubmed.ncbi.nlm.nih.gov/33973657/>

COMPARISON OF ELF-EMFS STIMULATION WITH CURRENT STIMULATION ON THE REGULATION OF LTP OF SC-CA1 SYNAPSES IN YOUNG RAT HIPPOCAMPUS

Yu Zheng, Wenjun Zhao, Xiaoxu Ma, Lei Dong, Lei Tian, Mei Zhou. Comparison of ELF-EMFs stimulation with current stimulation on the regulation of LTP of SC-CA1 synapses in young rat hippocampus. Int J Radiat Biol. 2021 May 10;1-21. doi: 10.1080/09553002.2021.1928781.

Abstract

Background: Long-term potentiation (LTP) is an important functional indicator for studying synaptic plasticity. Extremely low frequency electromagnetic fields (ELF-EMFs) are a physical means to regulate LTP. Time-varying magnetic fields induce induced currents. It is unknown which of the two parameters is the key factor when LTP is regulated by the magnetic field. **New Method:** A method is proposed for calculating the current value induced by ELF-EMFs. Then, a comparison of ELF-EMFs stimulation (100 Hz/2 mT and 200 Hz/2 mT) with current stimulation (0.1 μ A and 0.2 μ A) on the regulation of theta-burst or high-frequency stimulation (TBS/HFS)-LTP was performed.

Results: The LTP after ELF-EMFs and μA current regulation was significantly reduced compared with the corresponding value in the control group. The regulatory effect of 100 Hz/2 mT ELF-EMFs on LTP was stronger than 200 Hz/2 mT. However, the regulatory level of the currents (0.1 μA , 0.2 μA) was similar; the effect of 0.1 μA current was similar to that of 100 Hz/2 mT ELF-EMFs, while 0.2 μA had a stronger regulatory effect than that of 200 Hz/2 mT on HFS-LTP. Comparison with Existing Methods: Most of the existing methods were used to calculate the induced current in human models, while a calculation model of the induced current in the rat brain slice induced by ELF-EMFs was established in this work, which is more accurate than the existing methods.

Conclusions: This work demonstrated that both current and ELF EMFs stimulation reduced LTP. However, their effect was not exactly the same, indicating that the regulatory effect of ELF-EMFs on LTP is not entirely deriving from the induced current, since its magnetic mechanism might have played a certain role. <https://pubmed.ncbi.nlm.nih.gov/33970763/>

MAGNETIC FIELD-INDUCED Ca^{2+} INTAKE BY MESENCHYMAL STEM CELLS IS MEDIATED BY INTRACELLULAR Zn^{2+} AND ACCOMPANIED BY A Zn^{2+} INFLUX

Alp Özgün, Bora Garipcan. *Biochim Biophys Acta Mol Cell Res.* 2021 May 22;119062. doi: 10.1016/j.bbamcr.2021.119062.

Highlights

- Magnetic field-induced oxidative stress is not derived from mitochondrial activity.
- Ca^{2+} intake under magnetic field exposure is accompanied by Zn^{2+} intake.
- Magnetic field-evoked cation intake depends on availability of intracellular Zn^{2+} .
- 2-APB-sensitive pathways are major gateways for cation influx induced by magnetic fields.

Abstract

Chronic exposure to magnetic fields (MFs) has a diverse range of effects on biological systems but definitive molecular mechanisms of the interaction remain largely unknown. One of the most frequently reported effects of MF exposure is an elevated concentration of intracellular Ca^{2+} through disputed pathways. Other prominent effects include increased oxidative stress and upregulation of neural markers through EGFR activation in stem cells. Further characterization of cascades triggered by MF exposure is hindered by the phenotype diversity of biological models used in the literature. In an attempt to reveal more mechanistic data in this field, we combined the most commonly used biological model and MF parameters with the most commonly reported effects of MFs. Based on clues from the pathways previously defined as sensitive to MFs (EGFR and Zn^{2+} -binding enzymes), the roles of different types of channels (voltage gated Ca^{2+} channels, NMDA receptors, TRP channels) were inquired in the effects of 50 Hz MFs on bone marrow-derived mesenchymal stem cells. We report that, an influx of Zn^{2+} accompanies MF-induced Ca^{2+} intake, which is only attenuated by the broad-range inhibitor of TRP channels and store-operated Ca^{2+} entry (SOCE), 2-Aminoethoxydiphenyl borate (2-APB) among other blockers (memantine, nifedipine, ethosuximide and gabapentin). Interestingly, cation influx completely disappears when intracellular Zn^{2+} is chelated. Our results rule out voltage gated Ca^{2+} channels as a gateway to MF-induced Ca^{2+} intake and suggest Zn^{2+} -related channels as a new focus in the field. <https://pubmed.ncbi.nlm.nih.gov/34033861/>

RARE-EARTH MAGNETS INFLUENCE MOVEMENT PATTERNS OF THE MAGNETICALLY SENSITIVE NUDIBRANCH TRITONIA EXSULANS IN ITS NATURAL HABITAT

Russell C Wyeth, Theora Holden, Hamed Jalala, James A Murray. Rare-Earth Magnets Influence Movement Patterns of the Magnetically Sensitive Nudibranch *Tritonia exsulans* in Its Natural Habitat. *Biol Bull.* 2021 Apr;240(2):105-117. doi: 10.1086/713663.

Abstract

The nudibranch *Tritonia exsulans* (previously *Tritonia diomedea*) is known to have behaviors and neurons that can be modified by perturbations of the Earth's magnetic field. There is no definitive evidence for how this magnetic sense is used in nature. Using an exploratory approach, we tested for possible effects of magnetic perturbations based on underwater video of crawling patterns in the slugs' natural habitat, with magnets of varying strength deployed on the substrate. For analysis, we used a paired comparison of tracks of animals between segments 25-50 cm distant from the magnets and segments of the same tracks 0-25 cm from the magnets, to determine whether any differences depended on the strength of the magnet. Most track measurements (length, displacement, velocity, and tortuosity) showed no such differences. However, effects were observed for the changes in track headings between successive points. These results showed that tracks had relatively higher heading variability when they moved closer to stronger magnets. We suggest that this supports a hypothesis that *T. exsulans* continuously uses a magnetic sense to help maintain straight-line navigation. Further specific testing of the hypothesis is now needed to verify this new possibility for how animals can benefit from a compass sense. <https://pubmed.ncbi.nlm.nih.gov/33939940/>

MANMADE ELECTROMAGNETIC FIELDS AND OXIDATIVE STRESS—BIOLOGICAL EFFECTS AND CONSEQUENCES FOR HEALTH

David Schuermann, Meike Mevissen. Manmade Electromagnetic Fields and Oxidative Stress—Biological Effects and Consequences for Health. *Int. J. Mol. Sci.* 2021, 22(7), 3772; <https://doi.org/10.3390/ijms22073772>

(This article belongs to the Special Issue Molecular Mechanisms of Genotoxicity)

Abstract

Concomitant with the ever-expanding use of electrical appliances and mobile communication systems, public and occupational exposure to electromagnetic fields (EMF) in the extremely-low-frequency and radiofrequency range has become a widely debated environmental risk factor for health. Radiofrequency (RF) EMF and extremely-low-frequency (ELF) MF have been classified as possibly carcinogenic to humans (Group 2B) by the International Agency for Research on Cancer (IARC). The production of reactive oxygen species (ROS), potentially leading to cellular or systemic oxidative stress, was frequently found to be influenced by EMF exposure in animals and cells. In this review, we summarize key experimental findings on oxidative stress related to EMF exposure from animal and cell studies of the last decade. The observations are discussed in the context of molecular mechanisms and functionalities relevant to health such as neurological function, genome stability, immune response, and reproduction. Most animal and many cell studies showed increased oxidative stress caused by RF-EMF and ELF-MF. In order to estimate the risk for human health by manmade exposure, experimental studies in humans and epidemiological studies need to be considered as well.

Conclusions

The majority of recent animal studies on increased ROS production and oxidative stress caused by EMF were aimed at investigations of the nervous system and reproduction. Analogously, in cell studies, neurons or neuron-like cells were most frequently used. Animal studies on oxidative stress and possible impairment of reproduction at different stages (sperm maturation, very early stages of pregnancy such as implantation, and effects in newborns and after a few weeks of EMF exposure to the mother animals during pregnancy) follow in second place. These animal studies were supported by some cell studies, mainly in mouse cell lines of the male reproductive system and in sperm. Overall, more cells than animal studies were published, using, in addition to the above-mentioned cell types of the nervous and reproductive system, immune and cancer cells, as well as isolated cells from the skin and epithelia. For this report, animal and cell studies were included, according to their quality and research question, in order to give an informative overview of the available studies; however, this is not a systematic review.

In summary, indications for increased oxidative stress caused by RF-EMF and ELF-MF were reported in the majority of the animal studies and in more than half of the cell studies. Investigations in Wistar and Sprague-Dawley rats provided consistent evidence for oxidative stress occurring after RF-EMF exposure in the brain and testes and some indication of oxidative stress in the heart. Observations in Sprague-Dawley rats also seem to provide consistent evidence for oxidative stress in the liver and kidneys. In mice, oxidative stress induced by RF-EMF was predominantly demonstrated in the brain and testes, as well as in liver, kidneys, and ovaries. These observations were made with a variety of cell types, exposure times, and dosages (SAR or field strengths), within the range of the regulatory limits and recommendations. Certainly, some studies were subject to methodological uncertainties or weaknesses or are not very comprehensive regarding exposure time, dose, number, and quantitative analysis of the biomarkers used, to name a few. A trend is emerging, which becomes clear even when taking these methodological weaknesses into account, i.e., that EMF exposure, even in the low dose range, may well lead to changes in cellular oxidative balance. Organisms and cells are able to react to oxidative stress, and many observations after EMF exposure point to an adaptation after a recovery phase.

Adverse conditions, such as diseases (diabetes, neurodegenerative diseases), compromise the body's defense mechanisms, including antioxidant protection mechanisms, and individuals with such pre-existing conditions are more likely to experience health effects. The studies show that very young or old individuals can react less efficiently to oxidative stress, which of course also applies to other stressors that cause oxidative stress. Further investigations under standardized conditions are necessary to better understand and confirm these phenomena and observations.

Open access paper: <https://www.mdpi.com/1422-0067/22/7/3772>

SCIENCE, POLITICS, AND GROUPTHINK [HEALTH MATTERS]

James C. Lin. Science, Politics, and Groupthink [Health Matters]. IEEE Microwave Magazine. 22(5):24-26. Apr 1, 2021. DOI: 10.1109/MMM.2021.3056975.

Abstract

Discusses how the COVID-19 health pandemic worldwide was complicated by not only health and medical concerns, but the inclusion of politics, conspiracy theories, and social media. <https://ieeexplore.ieee.org/document/9393739>

In his latest column Professor Emeritus Lin criticizes the ICNIRP, the non-profit organization which the WHO relies upon for non-ionizing electromagnetic fields (EMF) exposure guidelines that it promotes worldwide. As you may know, this column is important not only because Professor Lin is one of the most respected EMF scientists in the world, he is the first scientist who has served on the ICNIRP Commission (2004 - 2016; chair of the committee on Physics & Engineering, 2008-2012; chair of the Radio Frequency group, 2012-2015) to challenge the credibility of ICNIRP's EMF exposure guidelines.

Excerpts

"Recently, a privately constituted group, with self-appointed membership, published a set of guidelines for limiting exposure to RF electromagnetic fields in the 100-kHz and 300-GHz frequency range [7]. The proposed guidelines were primarily based on the tissue-heating potentials of RF radiation to elevate animal body temperatures to greater than 1° C. While recognizing that the two aforementioned studies used large numbers of animals, best laboratory practice, and animals exposed for the entirety of their lives, the private group preferred to quibble with alleged "chance differences" between treatment conditions and the fact that the measured animal body core temperature changes reached 1° C, implying that a 1° C body core temperature rise is carcinogenic, ignoring the RF exposure. The group then pronounced that, when considered either in isolation or within the context of other animal carcinogenicity research, these findings do not provide evidence that RF radiation is carcinogenic.

Furthermore, the group noted that, even though many epidemiological studies of RF radiation associated with mobile phone use and cancer risk had been performed, studies on brain tumors, acoustic neuroma, meningioma, and parotid gland tumors had not provided evidence of an increased cancer risk. It suggested that, although somewhat elevated odds ratios were observed, inconsistencies and limitations, including recall or selection bias, precluded these results from being considered for setting exposure guidelines. The simultaneous penchant to dismiss and criticize positive results and the fondness for and eager acceptance of negative findings are palpable and concerning.

In contrast, the IARC's evaluation of the same epidemiological studies ended up officially classifying RF radiation as possibly carcinogenic to humans [2], [3].

An understandable question that comes to mind is this: How can there be such divergent evaluations and conclusions of the same scientific studies? Humans are not always rational or as transparent as advertised, and scientists are not impervious to conflicts of interest and can

be driven by egocentric motivations. Humans frequently make choices and decisions that defy clear logic.

Science has never been devoid of politics, believe it or not "

"Cellular mobile communication and associated wireless technologies have proven, beyond any debate, their direct benefit to humans. However, as for the verdict on the health and safety of billions of people who are exposed to unnecessary levels of RF radiation over extended lengths of time or even over their lifetimes, the jury is still out. When confronted with such divergent assessments of science, the ALARA—as low as reasonably achievable—practice and principle should be followed for RF health and safety."

[7] "ICNIRP guidelines for limiting exposure to electromagnetic fields (100 kHz to 300 GHz)," Health Phys., vol. 118, no. 5, pp. 483–524, 2020.

BACKGROUND DOCUMENT TO THE ADVISORY REPORT 5G AND HEALTH (HEALTH COUNCIL OF THE NETHERLANDS)

Health Council of the Netherlands. Background document to the advisory report 5G and health. Background document to 5G and health. The Hague: Health Council of the Netherlands, 2020; publication no. 2020/16Ae

Introduction

In this background document to the advisory report 5G and health, drafted by the Electromagnetic Fields Committee of the Health Council of the Netherlands, chapter 2 contains the search strategies used by the committee for the different topics. In chapter 3 the in- and exclusion criteria of the WHO are presented. In chapter 4 the committee gives an overview of the relevant publications on diseases and conditions, and in chapter 5 of the relevant publications on biological processes.

Open access background document report:

<https://www.healthcouncil.nl/binaries/healthcouncil/documents/advisory-reports/2020/09/02/5g-and-health/Background-document-to-the-advisory-report-5G-and-health.pdf>

Open access advisory report:

<https://www.healthcouncil.nl/binaries/healthcouncil/documents/advisory-reports/2020/09/02/5g-and-health/Advisory-report-5G-and-health.pdf>

IEEE C95.3-2021 - IEEE APPROVED DRAFT RECOMMENDED PRACTICE FOR MEASUREMENTS AND COMPUTATIONS OF EMF WITH RESPECT TO HUMAN EXPOSURE TO SUCH FIELDS, 0 HZ-300 GHZ

Institute of Electrical and Electronics Engineers. IEEE C95.3-2021 - IEEE Approved Draft Recommended Practice for Measurements and Computations of Electric, Magnetic and Electromagnetic Fields with Respect to Human Exposure to Such Fields, 0 Hz-300 GHz. 2021.

Standard Details

Best practices are described for the development, validation, and application of methods for the computation and measurement of relevant metrics characterizing human exposure to electric and/or magnetic fields (collectively, electromagnetic fields) over the frequency range of 0 Hz to 300 GHz. This recommended practice is a replacement for IEEE Std C95.3™-2002 and IEEE Std C95.3.1™-2010, extensively revising the contents from those and harmonizing with IEEE Std C95.1™-2019. Detailed methodology is not described; rather, requirements for best practice are expressed through guidance and references to other documents and standards. Examples are included to clarify the guidance. This recommended practice is intended for professional users who are familiar with basic electromagnetic field theory and practice and for persons involved in specifying or implementing critical hazard assessments or surveys such as those described in IEEE Std C95.7™-2014.

https://standards.ieee.org/standard/C95_3-2021.html

RADIATION PROTECTION STANDARD FOR LIMITING EXPOSURE TO RADIOFREQUENCY FIELDS - 100 KHZ TO 300 GHZ (ARPANSA)

Australian Radiation Protection and Nuclear Safety Agency (ARPANSA). Radiation Protection Standard for Limiting Exposure to Radiofrequency Fields - 100 kHz to 300 GHz. 2021, Radiation Protection Series S-1 (Rev. 1)

Open access report: https://www.arpansa.gov.au/sites/default/files/rps_s-1.pdf

THE EFFECT OF GSM ELECTROMAGNETIC FIELD EXPOSURE ON THE WAKING ELECTROENCEPHALOGRAM: METHODOLOGICAL INFLUENCES

Anna Dalecki, Adam Verrender, Sarah P Loughran, Rodney J Croft. The Effect of GSM Electromagnetic Field Exposure on the Waking Electroencephalogram: Methodological Influences. Bioelectromagnetics. 12 April 2021. <https://doi.org/10.1002/bem.22338>.

Abstract

Although there is consistent evidence that exposure to radiofrequency electromagnetic fields (RF-EMF) increases the spontaneous resting alpha spectral power of the electroencephalogram (EEG), the reliability of this evidence is uncertain as some studies have also failed to observe this effect. The present study aimed to determine whether the effect of RF-EMF exposure on EEG alpha power depends on whether EEG is derived from eyes open or closed conditions and assessed earlier (<5-min) versus later (>25-min) in the exposure interval.

Thirty-six adults participated in three experimental sessions, each involving one exposure: "Sham," "Low," and "High" RF-EMF corresponding to peak spatial specific absorption rates averaged over 10 g of 0, 1, and 2 W/kg, respectively. Resting EEG was recorded at baseline

(no exposure), during, and after exposure. Alpha power increase was found to be greater for the eyes open than eyes closed EEG during both the High ($P = 0.04$) and Low ($P = 0.04$) RF-EMF exposures. There was also a trend toward it being larger at the end, versus the start of the "High" 30-min exposure ($P < 0.01$; eyes open condition). This suggests that the use of eyes closed conditions, and insufficient RF-EMF exposure durations, are likely explanations for the failure of some studies to detect an RF-EMF exposure-related increase in alpha power, as such methodological choices decrease signal-to-noise ratios and increase type II error. <https://pubmed.ncbi.nlm.nih.gov/33847008/>

SIMULATED MOBILE COMMUNICATION FREQUENCIES (3.5 GHZ) EMITTED BY A SIGNAL GENERATOR AFFECTS THE SLEEP OF DROSOPHILA MELANOGASTER

Yahong Wang, Hongying Zhang, Ziyang Zhang, Boqun Sun, Chao Tang, Lu Zhang, Zhihao Jiang, Bo Ding, Yanyan Liao, Peng Cai. Simulated mobile communication frequencies (3.5 GHz) emitted by a signal generator affects the sleep of *Drosophila melanogaster*. *Environ Pollut.* 2021 Apr 5;283:117087. doi: 10.1016/j.envpol.2021.117087.

Abstract

With the rapid development of science and technology, 5G technology will be widely used, and biosafety concerns about the effects of 5G radiofrequency radiation on health have been raised. *Drosophila melanogaster* was selected as the model organism for our study, in which a 3.5 GHz radiofrequency radiation (RF-EMR) environment was simulated at intensities of 0.1 W/m², 1 W/m², and 10 W/m². The activity of parent male and offspring (F1) male flies was measured using a *Drosophila* activity monitoring system under short-term and long-term 3.5 GHz RF-EMR exposure. Core genes associated with heat stress, the circadian clock and neurotransmitters were detected by QRT-PCR technology, and the contents of GABA and glutamate were detected by UPLC-MS. The results show that short-term RF-EMR exposure increased the activity level and reduced the sleep duration while long-term RF-EMR exposure reduced the activity level and increased the sleep duration of F1 male flies. Under long-term RF-EMR, the expression of heat stress response-related hsp22, hsp26 and hsp70 genes was increased, the expression of circadian clock-related per, cyc, clk, cry, and tim genes was altered, the content of GABA and glutamate was reduced, and the expression levels of synthesis, transport and receptor genes were altered. In conclusion, long-term RF-EMR exposure enhances the heat stress response of offspring flies and then affects the expression of circadian clock and neurotransmitter genes, which leads to decreased activity, prolonged sleep duration, and improved sleep quality.

Highlights

- This study was simulating the most representative 3.5 GHz radio frequency radiation band of 5G in environment.
- This study was exploring the impact on the activity and sleep of organism under long-term RF-EMR.
- This study provided a reference for the hidden health effects brought by the development of science and technology.

<https://pubmed.ncbi.nlm.nih.gov/33894629/>

THE ADVERSE EFFECT OF MOBILE PHONE RADIATIONS ON DORSAL ROOT GANGLION OF ALBINO RATS

Faisal Taufiq, Mohammed Bhilal Babu, Aqeel Ahmad, Mohammed Eajaz, Ahmed Shariff, Noureldaim Elnoman Elbadawi, Semmal Syed Meerasa. The Adverse Effect of Mobile Phone Radiations on Dorsal Root Ganglion of Albino Rats. Journal of Pharmaceutical Research International, 19 March 2021. Pages 54-60. DOI: 10.9734/jpri/2021/v33i1331266.

Abstract

Objectives: To assess the effect of Mobile Phone Radio Frequency Electromagnetic Radiation (RF-EMR) on the histological structure of dorsal root ganglia in albino rats.

Methods: Twenty-four albino rats divided into one control and three experimental groups were studied for four weeks. The experimental groups were exposed to three different levels of RF-EMR through complete missed calls of 80,120 and160 calls per day respectively, using a GSM mobile phone of 0.9 GHz to1.8 GHz in silent mode. The dorsal root ganglia of the sacrificed Rats were examined under light and transmission electron microscope (TEM).

Results: Dorsal root ganglions of exposed rats showed considerable histological changes like reduction in cell size, condensation of cytoplasm, peripherally located heterochromatin nucleus, loss of nucleolus and densely packed myelinated nerve fibers. No such changes were observed in control rats.

Conclusion: Dorsal root ganglionic cells showed enduring and continuous changes when exposed to RF-EMR. The severity of histomorphological changes was dose-dependent, which increased constantly with radiation dosage increment. It might be fair to conclude that degenerative changes in the Dorsal Root Ganglion of the spinal cord, could be attributed to the long-term exposure to RF-EMR.

Open access paper: <https://www.journaljpri.com/index.php/JPRI/article/view/31266>

EFFECTS OF MOBILE PHONE EMISSIONS ON HUMAN RED BLOOD CELLS

Aniket Chowdhury, Yashveer Singh, Uttam Das, Deepak Waghmare, Raktim Dasgupta, Shovan Kumar Majumder. Effects of mobile phone emissions on human red blood cells. J Biophotonics. 2021 Apr 19;e202100047. doi: 10.1002/jbio.202100047.

Abstract

Raman spectroscopy was performed on GSM 900 MHz and 1800 MHz mobile phone signal exposed red blood cells (RBCs). The observed changes in the Raman spectra of mobile signal exposed RBCs compared to unexposed control suggest reduced hemoglobin-oxygen affinity for the exposed cells. The possible mechanism may involve activation of the voltage gated membrane Ca²⁺ channels by the mobile phone emissions resulting in an increase in the levels of adenosine triphosphate (ATP) and 2,3-diphosphoglycerate (2,3-DPG) in cells via altered metabolic activities. Further studies carried out with fluorescent Ca²⁺ indicator confirmed increased intracellular Ca²⁺ level in the exposed cells. Since intracellular ATP level influences the shape and mechanics of RBCs, exposed cells were studied using diffraction phase

microscopy and optical tweezers. Detectable changes in shape and mechanical properties were observed due to mobile signal exposure. <https://pubmed.ncbi.nlm.nih.gov/33871929/>

Excerpts

The RBC samples were exposed to mobile phone signal by keeping a commercial mobile handset (Nokia 105) in close proximity (~ few mm) of the glass cover slip. The handset was kept on call mode throughout the exposure time.

In the present study the SAR for the handset used is 1.501 W/kg which is less than the permissible limit of 1.6 W/kg prescribed in most parts of the world. The power density level from the typical GSM handsets is ~ 10⁻⁴ mW/cm² [33] and our mobile handset has no additional features like Global Positioning System (GPS), WiFi, Bluetooth etc. that may unwantedly add to the effect of basic GSM signals. ... thermal effects on blood samples due to low level microwave radiation from the handset should be negligible.

Further, an estimate of the power density at a distance of 100 meters (R) from a mobile network tower, typically emitting (P_t) ~ 100W, shows the radiation density is about 10⁻⁴ mW/cm². Such radiation exposure from network towers for people residing nearby will be of continuous nature [34] and closely resembles the pattern and level of exposure considered in present experiments. Such radiation exposure from network towers for people residing nearby will be of continuous nature [34] and closely resembles the pattern and level of exposure considered in present experiment.

ANTENNA/HUMAN BODY COUPLING IN 5G MILLIMETER-WAVE BANDS: DO AGE AND CLOTHING MATTER?

G. Sacco, D. Nikolayev, R. Sauleau, M. Zhadobov. Antenna/Human Body Coupling in 5G Millimeter-Wave Bands: Do Age and Clothing Matter? IEEE Journal of Microwaves, doi: 10.1109/JMW.2021.3063256.

Abstract

With the fast development of 5th generation (5G) mobile networks and prominence of the personal area networks and human-centered communications, people of all ages are increasingly exposed in the upper part of the microwave spectrum. In some exposure scenarios, presence of a textile between the radiating source and skin can affect the power absorption. This study investigates, for the first time, the effect of ageing and impact of textile on the power deposition in a skin-equivalent model under near-field exposure induced by multi-beam radiating structures at 26 GHz and 60 GHz. An array of four Yagi antennas has been used as a representative example of 5G reconfigurable antennas. The maximum increase of the averaged absorbed power density with respect to the averaged value for adults is observed at 70 year (8.8% at 26 GHz and 6.9% at 60 GHz). The strongest decrease is for 5-years-old children (-4.5% at 26 GHz and -3.7% at 60 GHz). In presence of a textile, the absorbed power density can increase or decrease depending on the textile properties (thickness and permittivity) and on the thickness of the air gap between textile and skin. With cotton and wool (considered as representative textile materials) the maximum increase of the averaged absorbed power density is about 40% compared to the bare skin.

Excerpts

"At microwave frequencies, the difference in exposure levels between children and adults was investigated [3]– [5]. It was demonstrated that, up to 5.6 GHz, the whole-body-average specific absorption rate (SAR) in children can go beyond the exposure limits [6], [7] by 40%–45% whereas remaining below these limits for adults (given the same incident field) [5], [8]. A more recent study [9] investigated the whole-body average SAR using the child models specified by the Commission on Radiological Protection (ICRP) instead of the scaled adult phantoms."

"This study deals with the electromagnetic exposure under near-field conditions considering typical reconfigurable antennas at 26 GHz and 60 GHz. For the first time, the analysis is performed taking into account biological tissue permittivity variations with age and presence of a textile in proximity or in contact with skin."

<https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9393791>

ARTIFICIAL EMG (ELECTROMYOGRAM) BY WLAN-EXPOSURE

L. von Klitzing. Artificial EMG by WLAN-Exposure. J Biostat Biometric App 6(1):101. 2021.

Abstract

WLAN (wireless local area network) is used as an important worldwide communication-technique. By this, always there is an electromagnetic field exposure. In contrast to the ICNIRP-safety guidelines, whereby no bioeffect is possible by these low-energetic electromagnetic fields, we found artificial signals in the nervous system in dependence on WLAN- exposure.

Open access letter: <http://www.annexpublishers.com/articles/JBIA/6101-Artificial-EMG-by-WLAN-Exposure.pdf>

30 HZ, COULD IT BE PART OF A WINDOW FREQUENCY FOR CELLULAR RESPONSE? (REVIEW)

Olga García-Minguillán, Ceferino Maestú. 30 Hz, Could It Be Part of a Window Frequency for Cellular Response? Int J Mol Sci. 2021 Mar 31;22(7):3642. doi: 10.3390/ijms22073642

Abstract

Many exogenous and endogenous risk factors have been proposed as precursors of brain tumors, including the exposure to non-ionizing electromagnetic fields. Nevertheless, there is still a debate among the scientific community about the hazard of the effects produced by non-ionizing radiation (NIR) because conflicting results have been found (number of articles reviewed >50). For that reason, to provide new evidence on the possible effects produced by exposure to NIR, we performed different studies with several combinations of extremely low frequencies, times, and field intensities in tumoral and non-tumoral cells. The results of our studies showed that cell viability was frequency dependent in glioblastoma cells. In fact, our results revealed that a frequency of 30 Hz-or even other frequencies close to 30 Hz-could constitute a window frequency determinant of the cellular response in tumoral and non-tumoral cells.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8036499/>

RESPONSE TO “CANCEROGENIC EFFECTS OF RADIOFREQUENCY RADIATION: A STATISTICAL REAPPRAISAL”

Fiorella Belpoggi, Laura Falcioni, Simona Panzacchi, Daria Sgargi, Daniele Mandrioli. Response to “Carcinogenic effects of radiofrequency radiation: A statistical reappraisal.” *Environmental Research*. 197, 2021,111067. <https://doi.org/10.1016/j.envres.2021.111067>.

No abstract. <https://www.sciencedirect.com/science/article/pii/S0013935121003613>

ACTIVATION OF MATRIX METALLOPROTEINASES AND FOXO3A IN HACAT KERATINOCYTES BY RADIOFREQUENCY ELECTROMAGNETIC FIELD EXPOSURE

Ju Hwan Kim, Dong-Jun Kang, Jun-Sang Bae, Jai Hyuen Lee, Sangbong Jeon, Hyung-Do Choi, Nam Kim, Hyung-Gun Kim, Hak Rim Kim. Activation of matrix metalloproteinases and FoxO3a in HaCaT keratinocytes by radiofrequency electromagnetic field exposure. *Sci Rep*. 2021 Apr 7;11(1):7680. doi: 10.1038/s41598-021-87263-2.

Abstract

As the skin is the largest body organ and critically serves as a barrier, it is frequently exposed and could be physiologically affected by radiofrequency electromagnetic field (RF-EMF) exposure. In this study, we found that 1760 MHz RF-EMF (4.0 W/kg specific absorption rate for 2 h/day during 4 days) exposure could induce intracellular reactive oxygen species (ROS) production in HaCaT human keratinocytes using 2',7'-dichlorofluorescein diacetate fluorescent probe analysis. However, cell growth and viability were unaffected by RF-EMF exposure. Since oxidative stress in the skin greatly influences the skin-aging process, we analyzed the skin senescence-related factors activated by ROS generation. Matrix metalloproteinases 1, 3, and 7 (MMP1, MMP3, and MMP7), the main skin wrinkle-related proteins, were significantly increased in HaCaT cells after RF-EMF exposure. Additionally, the gelatinolytic activities of secreted MMP2 and MMP9 were also increased by RF-EMF exposure. FoxO3a (Ser318/321) and ERK1/2 (Thr 202/Tyr 204) phosphorylation levels were significantly increased by RF-EMF exposure. However, Bcl2 and Bax expression levels were not significantly changed, indicating that the apoptotic pathway was not activated in keratinocytes following RF-EMF exposure. In summary, our findings show that exposure to 1760 MHz RF-EMF induces ROS generation, leading to MMP activation and FoxO3a and ERK1/2 phosphorylation. These data suggest that RF-EMF exposure induces cellular senescence of skin cells through ROS induction in HaCaT human keratinocytes.

Conclusion

In conclusion, exposure to 1760 MHz RF-EMF at 4 W/kg SAR induced intracellular ROS generation, which then stimulated MMPs (MMP1, 2, 3, 7, and 9) and activated the ERK1/2 (phospho-ERK1/2) and FoxO3a (phospho-FoxO3a) signaling pathways in HaCaT cells. Our results suggest that these changes induced by RF-EMF exposure would contribute to skin-aging processes.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8027011/>

1800 MHZ RADIOFREQUENCY ELECTROMAGNETIC FIELD IMPAIRS NEURITE OUTGROWTH THROUGH INHIBITING EPHA5 SIGNALING

Chunhai Chen, Qinglong Ma, Ping Deng, Min Lin, Peng Gao, Mindi He, Yonghui Lu, Huifeng Pi, Zhixin He, Chao Zhou, Yanwen Zhang, Zhengping Yu, Lei Zhang. 1800 MHz Radiofrequency Electromagnetic Field Impairs Neurite Outgrowth Through Inhibiting EPHA5 Signaling. *Front Cell Dev Biol.* 2021 Apr 12. 9:657623. doi:10.3389/fcell.2021.657623.

Abstract

The increasing intensity of environmental radiofrequency electromagnetic fields (RF-EMF) has increased public concern about its health effects. Of particular concern are the influences of RF-EMF exposure on the development of the brain. The mechanisms of how RF-EMF acts on the developing brain are not fully understood. Here, based on high-throughput RNA sequencing techniques, we revealed that transcripts related to neurite development were significantly influenced by 1800 MHz RF-EMF exposure during neuronal differentiation. Exposure to RF-EMF remarkably decreased the total length of neurite and the number of branch points in neural stem cells-derived neurons and retinoic acid-induced Neuro-2A cells. The expression of Eph receptors 5 (EPHA5), which is required for neurite outgrowth, was inhibited remarkably after RF-EMF exposure. Enhancing EPHA5 signaling rescued the inhibitory effects of RF-EMF on neurite outgrowth. Besides, we identified that cAMP-response element-binding protein (CREB) and RhoA were critical downstream factors of EPHA5 signaling in mediating the inhibitory effects of RF-EMF on neurite outgrowth. Together, our finding revealed that RF-EMF exposure impaired neurite outgrowth through EPHA5 signaling. This finding explored the effects and key mechanisms of how RF-EMF exposure impaired neurite outgrowth and also provided a new clue to understanding the influences of RF-EMF on brain development.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8075058/>

MATERNAL USE OF INDUCTION HEATING COOKERS DURING PREGNANCY AND BIRTH OUTCOMES: THE KYUSHU OKINAWA MATERNAL AND CHILD HEALTH STUDY

Akiko Tokinobu, Keiko Tanaka, Masashi Arakawa, Yoshihiro Miyake. Maternal Use of Induction Heating Cookers During Pregnancy and Birth Outcomes: The Kyushu Okinawa Maternal and Child Health Study. *Bioelectromagnetics.* 2021 Apr 12. doi: 10.1002/bem.22339.

Abstract

The effects of exposure to intermediate-frequency electromagnetic fields (IF-EMFs) during pregnancy on birth outcomes are uncertain. We investigated the association between the use of induction heating (IH) cookers, which are major sources of IF-EMFs, during pregnancy and preterm birth (PTB), low birth weight (LBW), small- for-gestational-age (SGA), and birth weight, using data from a prebirth cohort study in Japan. Study participants were 1,565 mothers with singleton pregnancies and the babies born from these pregnancies. We collected the data presented here using self-administered questionnaires. An adjustment was made for maternal age, region of residence, number of children, family structure, maternal education, maternal employment, maternal alcohol intake, smoking during pregnancy, maternal body mass index, baby's sex, and gestational age at birth. IH cooker use during pregnancy was independently associated with a reduced risk of PTB: the adjusted odds ratio

was 0.28 (95% confidence interval: 0.07-0.78). IH cooker use during pregnancy was not associated with LBW, SGA, or birth weight. This is the first study to show that IH cooker use during pregnancy is independently inversely associated with PTB.
<https://pubmed.ncbi.nlm.nih.gov/33846994/>

EFFECTS OF LONG-TERM EXPOSURE OF INTERMEDIATE FREQUENCY MAGNETIC FIELDS (20 KHZ, 360 μ T) ON THE DEVELOPMENT, PATHOLOGICAL FINDINGS, AND BEHAVIOR OF FEMALE MICE

Alexander Lerchl, Karen Drees Née Grote, Isabel Gronau, Dirk Fischer, Julia Bauch, Axel Hoppe. Effects of Long- Term Exposure of Intermediate Frequency Magnetic Fields (20 kHz, 360 μ T) on the Development, Pathological Findings, and Behavior of Female Mice. Bioelectromagnetics. 2021 Apr 6. doi: 10.1002/bem.22337.

Abstract

The use of magnetic fields in the intermediate-frequency (IF) range to wirelessly charge electric cars with power transfer in the kilowatt range has become increasingly widespread, leading to unavoidable stray fields in the microtesla range. Only a handful of studies have assessed the potential biological risks associated with exposure to such fields. We exposed female mice (n = 80 per group) to either 20 kHz, 360 μ T (rms), or sham in Helmholtz coils to conduct a blind design study. Exposure started at 3 months of age (24 h/day). Body mass was recorded every 1-2 weeks. At 10 months of age, three behavioral tests were performed on 24 animals per group. Three months later, the mice were sacrificed and organs (brain, liver, kidney, spleen, and lung) were removed and prepared for microscopic analysis. Our findings demonstrate no differences in the development of body mass and survival rates (96% and 89%, respectively). Similarly, no significant differences were observed in tumor incidence rates. When it comes to behavioral tests, the 8-arm maze results revealed no significant differences. In contrast, the Rotarod data were significantly ($P < 0.001$) different with longer retention times seen in the exposed mice. In the open field, the number of supported rears was significantly lower ($P < 0.01$), whereas the other endpoints did not show any differences. Overall, our data reveal no adverse effects of exposure to 20 kHz, 360 μ T on the development and tumor incidences, while the significant differences in the behavioral tests may indicate higher levels of alertness in mice.

Excerpts

Our results indicate that exposure to IF-MF does not increase the risk of pathological findings. However, in this study, only brain, liver, kidney, spleen, and lung tissue were analyzed for neoplastic lesions; the OECD guidelines recommend that some 40 tissues should be analyzed. Another potential limitation of our study is the relatively short exposure period of 10 months (OECD recommends 24 months of treatment). Yet another, but unavoidable, limitation was that for technical reasons the exposure could not be performed during the behavioral experiments. This is because interference of the strong magnetic field with the equipment (CCD camera or Rotarod) would inevitably have led to the person performing the experiment knowing whether or not the animals were being exposed....

It is also pertinent to understand the limitations of animal models as predictors of human biology. Therefore, the comparability and applicability of the results to humans need to be verified. Furthermore, this study only examined female mice and it is well-established that

sex plays a crucial role in terms of cancer incidence, prognosis, and mortality [Zhu et al., 2019]. Therefore, further long-term studies that include male animals are needed to investigate the biological impact of IF-MF exposure....

Taken together, the data presented in this study indicate some behavioral effects in the exposed animals, while no differences were observed in growth and tumorigenesis between exposed and control mice. The potentially increased stress levels must be regarded as mild since they did not cause a drop in body mass, which is a very sensitive parameter for stress in rodents [Jeong et al., 2013]. Therefore, the overall results do not indicate that chronic exposure to 20 kHz magnetic fields at 360 μ T poses a health risk. The overall conclusion of this study as well as of data from the literature is that there is a lack of evidence that IF-MF exposure contributes to any significant behavioral changes or influences growth or malignancy in the female mice model.

My note: According to Table 1, the exposed mice had a total of 17 pathological findings vs. 11 in the control mice.

CLASSIFICATION OF THE LAPTOP'S EXTREMELY LOW FREQUENCY MAGNETIC FIELD WITH ALTERNATING CURRENT AND BATTERY

Dejan Tanikić, Alessia Amelio. Classification of the Laptop's Extremely Low Frequency Magnetic Field With Alternating Current and Battery. 2021 20th International Symposium INFOTEH-JAHORINA(INFOTEH).17-19March2021.DOI:10.1109/INFOTEH51037.2021.9400677

Abstract

The magnetic field exposure of the laptop users represents an important task to be researched. This paper proposes an artificial neural network model for the prediction of the extremely low-frequency magnetic field produced by laptops. It is based on measurement data of 10 laptops and a 3-input vector representing typical laptop characteristics. The novelty of the proposed model is that it is created in order to predict the level of the magnetic field emission produced by a laptop when it is powered by both alternating current and battery. Then, the obtained data are classified by the self-organizing map method to describe the safe or dangerous level of a laptop. At the end, a few suggestions are proposed for a safer use of the laptops. <https://ieeexplore.ieee.org/document/9400677>

Excerpts

The magnetic field is measured at 18 measurement points in the laptop's neighbourhood. These points are at the top (T1...T9) and at the bottom (B1...B9) of the laptop [6], [10].

Also, we measured the magnetic field 10 cm away from the proposed measurement points, i.e. outside the laptops. The obtained levels of the magnetic field were around or below 0.01 μ T. Hence, they can be considered as a background magnetic field.

The general conclusion is that the value of the magnetic field is almost 2 times higher when the laptops are AC powered. The predicted values of the magnetic field of the 10 laptops were used to divide the laptops in 5 (4) clusters, which represent the grade of the magnetic field of each laptop. Furthermore, if we compare the ELF

magnetic field emission and the classes extracted by the clustering process, we can obtain the results given in Tables 3–4 for AC and battery powered laptops, respectively.

This paper proposed a new ANN model for estimating the ELF magnetic field emission based on the measurement of 10 laptops. The measurement of the magnetic field was performed at the top and at the bottom of each laptop which was AC and battery powered. Then, from the ANN predicted data, a SOM based clustering was performed. The obtained classification was used to clearly define a level ranking of the laptops according to their ELF magnetic field emission. In the end, important suggestions for a safer use of the laptops were proposed. In the future, we are planning to extend the analysis to the tablet devices

HONEY BEE PROTEOLYTIC SYSTEM AND BEHAVIOR PARAMETERS UNDER THE INFLUENCE OF AN ELECTRIC FIELD AT 50 HZ AND VARIABLE INTENSITIES FOR A LONG EXPOSURE TIME

Paweł Migdał, Agnieszka Murawska, Aneta Strachecka, Paweł Bieńkowski, Adam Roman. Honey Bee Proteolytic System and Behavior Parameters under the Influence of an Electric Field at 50 Hz and Variable Intensities for a Long Exposure Time. *Animals (Basel)*. 2021 Mar 18;11(3):863. doi: 10.3390/ani11030863.

Abstract

The effect of an artificial electromagnetic field on organisms is a subject of extensive public debate and growing numbers of studies. Our study aimed to show the effect of an electromagnetic field at 50 Hz and variable intensities on honey bee proteolytic systems and behavior parameters after 12 h of exposure. Newly emerged worker bees were put into cages and exposed to a 50 Hz E-field with an intensity of 5.0 kV/m, 11.5 kV/m, 23.0 kV/m, or 34.5 kV/m. After 12 h of exposure, hemolymph samples were taken for protease analysis, and the bees were recorded for behavioral analysis. Six behaviors were chosen for observation: walking, flying, self-grooming, contact between individuals, stillness, and wing movement. Bees in the control group demonstrated the highest number of all behavior occurrences, except flying, and had the lowest protease activity. Bees in the experimental groups showed a lower number of occurrences of walking, self-grooming, and contacts between individuals than the control bees and had significantly higher protease activity than the control bees (except that of alkaline proteases in the 23.0 kV/m group).

Simple Summary

The amount of electromagnetic field (EMF) in the environment emitted by electrical and electronic devices, mobile phone masts, or power lines is constantly increasing. Honey bees can be exposed to the EMF in the environment, and the influence of this factor on bees is still under consideration. Studying the impact of EMF on honey bees can give valuable information about whether it poses a threat to them. The honey bee is an important pollinator, playing a significant role in maintaining biodiversity and food production. Our research showed that a 50 Hz electric field at various intensities reduced the number of occurrences of walking, contacts between individuals, and self-grooming, and increased the activity of proteases, which are involved in the immune system response.

Conclusions

The amount of artificial electromagnetic field in the environment is constantly increasing, thus the honey bee is exposed to this factor. In our study, bees in the control group demonstrated the highest number of all behavior occurrences, except flying, and had the lowest activity of all types of proteases. Bees in the experimental groups showed a lower number of walking, self-grooming, and contact between individual occurrences than control bees and had higher protease activity than control bees. Our results show that an E-field is potential harmful factor to the honey bee. However, we do not know if the changes in behavior and protease activity of the honey bee after E-field exposure persist and for how long. It would be important to investigate behavior parameters and biochemical markers at different time intervals after the end of exposure to an E-field. It can be helpful to determine the interaction between the biochemical marker activity and behavioral parameters. Such an observation could provide a better understanding of the immune response of the honey bee exposing to environmental stressors.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8003097/>

EXTREMELY LOW-FREQUENCY ELECTROMAGNETIC FIELDS ENTRAIN LOCUST WINGBEATS

Sebastian Shepherd, Christopher W Jackson, Suleiman M Sharkh, Hitoshi Aonuma, Eugenio E Oliveira, Philip L Newland. Extremely Low-Frequency Electromagnetic Fields Entrain Locust Wingbeats. Bioelectromagnetics. 2021 Apr 6. doi: 10.1002/bem.22336.

Abstract

Extremely low-frequency electromagnetic fields (ELF EMFs) have been shown to impact the behavior and physiology of insects. Recent studies have highlighted the need for more research to determine more specifically how they affect flying insects. Here, we ask how locust flight is affected by acute exposure to 50 Hz EMFs. We analyzed the flights of individual locusts tethered between a pair of copper wire coils generating EMFs of various frequency using high-speed video recording. The mean wingbeat frequency of tethered locusts was 18.92 ± 0.27 Hz. We found that acute exposure to 50 Hz EMFs significantly increased absolute change in wingbeat frequency in a field strength-dependent manner, with greater field strengths causing greater changes in wingbeat frequency. The effect of EMFs on wingbeat frequency depended on the initial wingbeat frequency of a locust, with locusts flying at a frequency lower than 20 Hz increasing their wingbeat frequency, while locusts flying with a wingbeat frequency higher than 20 Hz decreasing their wingbeat frequency. During the application of 50 Hz EMF, the wingbeat frequency was entrained to a 2:5 ratio (two wingbeat cycles to five EMF cycles) of the applied EMF. We then applied a range of ELF EMFs that were close to normal wingbeat frequency and found that locusts entrained to the exact frequency of the applied EMF. These results show that exposure to ELF EMFs lead to small but significant changes in wingbeat frequency in locusts. We discuss the biological implications of the coordination of insect flight in response to electromagnetic stimuli. <https://pubmed.ncbi.nlm.nih.gov/33822398/>

CURRENT PROGRESS ON THE EFFECT OF MOBILE PHONE RADIATION ON SPERM QUALITY: AN UPDATED SYSTEMATIC REVIEW AND META-ANALYSIS OF HUMAN AND ANIMAL STUDIES

Gang Yu, Zhiming Bai, Song Chao, Qing Cheng, Gang Wang, Zeping Tang, Sixing Yang. Current progress on the effect of mobile phone radiation on sperm quality: an updated systematic review and meta-analysis of human and animal studies. *Environmental Pollution*. 30 March 2021. <https://doi.org/10.1016/j.envpol.2021.116952>.

Highlights

- Mobile phone use was related to sperm quality decline of men in some areas.
- Mobile phone RF-EMR directly impaired mature sperm of men in vitro.
- Mobile phone RF-EMR affected some parameters of sperm quality in experiment animals.
- Experiment conditions affected pooled results of animal experiments.
- More studies should be conducted to investigate this issue in new era.

Abstract

Potential suppression of fertility due to mobile phone radiation remains a focus of researchers. We conducted meta-analyses on the effects of mobile phone radiation on sperm quality using recent evidence and propose some perspectives on this issue. Using the MEDLINE/PubMed, Embase, WOS, CENTRAL, and ClinicalTrials.gov databases, we retrieved and screened studies published before December 2020 on the effects of mobile phone use/mobile phone RF-EMR on sperm quality. Thirty-nine studies were included. Data quality and general information of the studies were evaluated and recorded. Sperm quality data (density, motility, viability, morphology, and DFI) were compiled for further analyses, and we conducted subgroup, sensitivity, and publication bias analyses. The pooled results of human cross-sectional studies did not support an association of mobile phone use and a decline in sperm quality. Different study areas contributed to the heterogeneity of the studies. In East Europe and West Asia, mobile phone use was correlated with a decline in sperm density and motility. Mobile phone RF-EMR exposure could decrease the motility and viability of mature human sperm in vitro. The pooled results of animal studies showed that mobile phone RF-EMR exposure could suppress sperm motility and viability. Furthermore, it reduced sperm density in mice, in rats older than 10 weeks, and in rats restrained during exposure. Differences regarding age, modeling method, exposure device, and exposure time contributed to the heterogeneity of animal studies. Previous studies have extensively investigated and demonstrated the adverse effects of mobile phone radiation on sperm. In the future, new standardized criteria should be applied to evaluate potential effects of mobile phone RF-EMR dosages. Further sperm-related parameters at the functional and molecular levels as well as changes in biological characteristics of germ cells should be evaluated. Moreover, the impact of mobile phone RF-EMR on individual organs should also be examined.

Conclusion

The results of our meta-analysis indicated that in East Europe and West Asia, mobile phone use is associated with a decline in human sperm density and motility. Mobile phone RF-EMR can reduce motility and viability of mature human sperm in vitro, and it can also reduce sperm motility and viability in male animals and decrease sperm density of sexually mature restrained rats. Some important factors that affect the results of animal experiments are study setup and radiation device as well as age and exposure time. Our study is an extension

of previous studies and has scientific value for future studies on effects of mobile phone RF-EMR associated with sperm quality.

<https://www.sciencedirect.com/science/article/abs/pii/S0269749121005340>

PROTOCOL FOR PERSONAL RF-EMF EXPOSURE MEASUREMENT STUDIES IN 5TH GENERATION TELECOMMUNICATION NETWORKS

Maarten Velghe, Sam Aerts, Luc Martens, Wout Joseph, Arno Thielens. Protocol for personal RF-EMF exposure measurement studies in 5th generation telecommunication networks. *Environ Health*. 2021 Apr 1;20(1):36. doi: 10.1186/s12940-021-00719-w.

Abstract

Background: The general population is exposed to Radio-Frequency Electromagnetic Fields (RF-EMFs) used by telecommunication networks. Previous studies developed methods to assess this exposure. These methods will be inadequate to accurately assess exposure in 5G technologies or other wireless technologies using adaptive antennas. This is due to the fact that 5G NR (new radio) base stations will focus actively on connected users, resulting in a high spatio-temporal variations in the RF-EMFs. This increases the measurement uncertainty in personal measurements of RF-EMF exposure. Furthermore, a user's exposure from base stations will be dependent on the amount of data usage, adding a new component to the auto-induced exposure, which is often omitted in current studies.

Goals: The objective of this paper is to develop a general study protocol for future personal RF-EMF exposure research adapted to 5G technologies. This protocol will include the assessment of auto-induced exposure of both a user's own devices and the networks' base stations.

Method: This study draws from lessons learned from previous RF-EMF exposure research and current knowledge on 5G technologies, including studies simulating 5G NR base stations and measurements around 5G NR test sites.

Results: To account for auto-induced exposure, an activity-based approach is introduced. In survey studies, an RF-EMF sensor is fixed on the participants' mobile device(s). Based on the measured power density, GPS data and movement and proximity sensors, different activities can be clustered and the exposure during each activity is evaluated. In microenvironmental measurements, a trained researcher performs measurements in predefined microenvironments with a mobile device equipped with the RF-EMF sensor. The mobile device is programmed to repeat a sequence of data transmission scenarios (different amounts of uplink and downlink data transmissions). Based on simulations, the amount of exposure induced in the body when the user device is at a certain location relative to the body, can be evaluated.

Conclusion: Our protocol addresses the main challenges to personal exposure measurement introduced by 5G NR. A systematic method to evaluate a user's auto-induced exposure is introduced.

<https://pubmed.ncbi.nlm.nih.gov/33794922/>

NONUNIFORM EXPOSURE TO THE CORNEA FROM MILLIMETER WAVES

Kenneth R Foster, Ilkka Laakso, Steven Chalfin. Health Phys. 2021 May 1;120(5):525-531. doi: 10.1097/HP.0000000000001376.

Abstract

This study examines the nonuniform exposure to the cornea from incident millimeter waves at 94-100 GHz. Two previous studies measured temperature increases in the rhesus cornea exposed to brief (1-6 s) pulses of high- fluence millimeter waves (94 GHz), one of which also estimated thresholds for corneal damage (reported as ED50, the dose resulting in a visible lesion 50% of the time). Both studies noted large variations in the temperature increase across the surface of the cornea due to wave interference effects. This study examines this variability using high-resolution simulations of mm-wave absorption and temperature increase in the human cornea from exposures to plane wave energy at 100 GHz. Calculations are based on an earlier study. The simulations show that the peak temperature increases in the cornea from short exposures (up to 10 s) to high- intensity mm-wave pulses are 1.7-2.8 times the median increase depending on the polarization of the incident energy. A simple one-dimensional "baseline" model provides a good estimate of the median temperature increase in the cornea. Two different estimates are presented for the thresholds for producing thermal lesions, expressed in terms of the minimum fluence of incident 100 GHz pulses. The first estimate is based on thresholds for thermal damage from pulsed infrared energy, and the second is based on a thermal damage model. The mm- wave pulses presently considered far exceed current IEEE or ICNIRP exposure limits but may be produced by some nonlethal weapons systems. Interference effects due to wave reflections from structures in and near the eye result in highly localized variations in energy absorbed in the cornea and surrounding facial tissues and are important to consider in a hazard analysis for exposures to intense pulsed millimeter waves. <https://pubmed.ncbi.nlm.nih.gov/33760769/>

Discussion

Both IEEE C95.1-2019 and ICNIRP (2020) specifically limit the fluence of brief, high intensity mm-wave pulses to avoid excessive temperature rises. The limits discussed below refer to power densities measured outside the body, which are defined in ICNIRP 2020 as "reference levels" and in IEEE as "exposure reference levels." For 2-s pulses of mm-waves, IEEE C95.1-2019 (IEEE 2019) has fluence limits of 0.28 and 1.4 kJ m⁻² for persons in unrestricted or restricted environments (equivalent to general public and occupational limits). The corresponding limits in ICNIRP (2020) (for 100 GHz pulses) are 1.7 and 8.6 kJ m⁻² for general public and occupational exposures, respectively. All of the exposures in Table 2 far exceed these limits. Exposures summarized in Table 2 would be acutely painful (Beuerman and Talenian 1979) and would trigger blink reflexes in most (but perhaps not all) awake subjects that would protect them from corneal damage. However, interference effects will also cause nonuniform absorption patterns away from the cornea, which is apparent in Fig. 1. In their corneal damage studies on rabbits, Kojima et al. (2009) noted significant thermal damage to the eyelids as well, to the extent that the eyelids had to be held open with tape during exposure to obtain reproducible corneal damage data. When analyzing potential hazards from the Active Denial nonlethal weapons system or other sources of intense mm-wave exposures, local enhancements in exposure due to interference effects over an appropriate range of subjects and exposure parameters would need to be considered.

PROTECTION OF THE POPULATION HEALTH FROM ELECTROMAGNETIC HAZARDS - CHALLENGES RESULTING FROM THE IMPLEMENTATION OF THE 5G NETWORK PLANNED IN POLAND

Marek Zmyślony, Paweł Bieńkowski, Alicja Bortkiewicz, Jolanta Karpowicz, Jarosław Kieliszek, Piotr Politański, Konrad Rydzyński. [Protection of the population health from electromagnetic hazards - challenges resulting from the implementation of the 5G network planned in Poland] [Article in Polish]. Med Pr. 2020 Jan 20;71(1):105-113. doi: 10.13075/mp.5893.00867.

Abstract in English

There is an ongoing discussion about electromagnetic hazards in the context of the new wireless communication technology - the fifth generation (5G) standard. Concerns about safety and health hazards resulting from the influence of the electromagnetic field (EMF) emitted by the designed 5G antennas have been raised. In Poland, the level of the population's exposure to EMF is limited to 7 V/m for frequencies above 300 MHz. This limitation results from taking into account the protective measures related not only to direct thermal hazards, but also to diversified indirect and long-term threats. Many countries have not established legal requirements in this frequency range, or they have introduced regulations based on recommendations regarding protection against direct thermal risks only (Council Recommendation 1999/519/EC). For such protection, the permissible levels of electric field intensity are 20-60 V/m (depending on the frequency). This work has been created through an interdisciplinary collaboration of engineers, biologists and doctors, who have been for many years professionally dealing with the protection of the biosphere against the negative effects of EMF. It presents the state of knowledge on the biological and health effects of the EMF emitted by mobile phone devices (including millimeter waves which are planned to be used in the 5G network). A comparison of the EU recommendations and the provisions on public protection being in force in Poland was made against this background. The results of research conducted to date on the biological effects of the EMF radiofrequency emitted by mobile telecommunication devices, operating with the frequencies up to 6 GHz, do not allow drawing any firm conclusions; however, the research evidence is strong enough for the World Health Organization to classify EMF as an environmental factor potentially carcinogenic to humans. At the moment, there is a shortage of adequate scientific data to assess the health effects of exposure to electromagnetic millimeter waves, which are planned to be used in the designed 5G devices. Nevertheless, due to the fact that there are data indicating the existence of biophysical mechanisms of the EMF influence that may lead to adverse health effects, it seems necessary to use the precautionary principle and the ALARA principle when creating environmental requirements for the construction and exploitation of the infrastructure of the planned 5G system.<http://medpr.imp.lodz.pl/Ochrona-zdrowia-ludnosci-przed-zagrozeniami-elektromagnetycznymi-wyzwania-wynikajace,114069,0,2.html>

See also:

Bieńkowski, Paweł, Marek Zmyślony, Jolanta Karpowicz, Piotr Politański, Alicja Bortkiewicz, Jarosław Kieliszek, and Konrad Rydzyński. [Conditionings of population exposure to electromagnetic fields associated with the rational use of 5G radiocommunication networks in Poland] [Article in Polish]. Medycyna Pracy 71 no. 2 (2020): 245-253. doi:10.13075/mp.5893.00920. <http://medpr.imp.lodz.pl/Uwarunkowania-ekspozycji-ludnosci-na-pole-elektromagnetyczne-zwiazane-z-uzytkowaniem,114070,0,2.html>

CELLULAR PHONE USER'S AGE OR THE DURATION OF CALLS MODERATE AUTONOMIC NERVOUS SYSTEM? A META-ANALYSIS

Styliani A Geronikolou, Ölle Johansson, George Chrousos, Christina Kanaka-Gantenbein, Dennis Cokkinos. Cellular Phone User's Age or the Duration of Calls Moderate Autonomic Nervous System? A Meta- Analysis. Adv Exp Med Biol. 2020;1194:475-488. doi: 10.1007/978-3-030-32622-7_46.

Abstract

Background: The European Health Risk Assessment Network on Electromagnetic Fields Exposure (EFHRAN) reported in 2012: "Children and adolescents represent the first generation of Europeans to be exposed to diffuse EMF since their conception and birth, thus, they are expected to be more sensitive to these fields." On the other hand, the body's physiologic processes are regulated by the autonomic nervous system (ANS) in a way that warrants further elucidation.

Objective: Age and duration of exposure are investigated for modifying the variance of the reported effects of mobile/cellular phone call (CPC) on ANS indexed, herein, by the heart rate variability (HRV).

Method: Five studies targeted to 124 healthy subjects (aged 15.3-28.4 years (24.3 ± 5.2), who have been recorded in supine position before and/or sham versus real exposure (to a CPC lasting 5-32 minutes), are included in this meta-analysis. Age and duration of a CPC are evaluated as predictors in two separate meta- regressions.

Results: The meta-analysis identified a heterogeneity $I^2 = 63.2\%$ for all outcomes and $I^2 = 65.2\%$ for sympathovagal balance. Thus, we performed meta-regressions: for the sympathovagal balance rather than the combined parameters effect, the goodness of fit model presents significance only for age - the residual sum of squares compared to chi-square distribution (QR) is 4.24 for age ($p = 0.12$) - while, for minutes of exposure, QR = 8.2805 ($p = 0.016$).

Conclusions: The sympathovagal balance - indispensable for health/homeostasis maintenance is strongly predicted by age. Minutes of exposure did not affect overall HRV or sympathovagal balance. The results endorse/validate the EFHRAN 2012 suggestion for future research targeting to youngsters. <https://pubmed.ncbi.nlm.nih.gov/32468564/>

Excerpt

... the effect of age is higher in younger ages (adolescence) and decreases as age increases (Figs. 2a and 3a). This result supports the caution already expressed that the risk for children and adolescents from radiofrequencies exposure needs to be extensively investigated (Akselrod et al. 1981; Hjortskov et al. 2004). Our finding may be either explained in terms of development or in terms of accumulated exposure of each individual to electromagnetic fields even since conception (exposome). Indeed, potential health effects from this technology expansion in pediatric and adolescent populations are suggested by the European Health Risk Assessment Network on Electromagnetic Fields Exposure in its 2012 report (Nam et al. 2011; Morgan et al. 2014a): "These groups represent the first generation of Europeans

to be exposed to diffuse EMF fields since conception and birth, thus, are expected to be more sensitive to these fields.”

Conclusively, our analysis suggests that (a) minutes of exposure (minutes of speaking on the mobile phone) do not affect the autonomic nervous system of the heart or its sympathovagal balance, (b) age synergizes with other variables (exposome, SAR) so as to explain combined HRV parameter effects, and (c) the sympathovagal balance is strongly predicted by age, implying that adolescents’ autonomic nervous system seems to be more vulnerable to the CPC exposure. Additionally, risk assessment analysis of the additive effects of continuous exposure (exposome) is strongly suggested.

WHAT IS THE RADIATION BEFORE 5G? A CORRELATION STUDY BETWEEN MEASUREMENTS IN SITU AND IN REAL TIME AND EPIDEMIOLOGICAL INDICATORS IN VALLECAS, MADRID

Isabel López, Nazario Félix, Marco Rivera, Adrián Alonso, Ceferino Maestú. What is the radiation before 5G? A correlation study between measurements in situ and in real time and epidemiological indicators in Vallecas, Madrid. *Environ Res.* 2021 Mar;194:110734. doi: 10.1016/j.envres.2021.110734.

Abstract

Background: Exposure of the general population to electromagnetic radiation emitted by mobile phone base stations is one of the greater concerns of residents affected by the proximity of these structures due to the possible relationship between radiated levels and health indicators.

Objectives: This study aimed to find a possible relationship between some health indicators and electromagnetic radiation measurements.

Methods: A total of 268 surveys, own design, were completed by residents of a Madrid neighborhood surrounded by nine telephone antennas, and 105 measurements of electromagnetic radiation were taken with a spectrum analyzer and an isotropic antenna, in situ and in real - time, both outside and inside the houses.

Results: It was shown statistically significant p - values in headaches presence ($p = 0.010$), nightmares ($p = 0.001$), headache intensity ($p < 0.001$), dizziness frequency ($p = 0.011$), instability episodes frequency ($p = 0.026$), number of hours that one person sleeps per day ($p < 0.001$) and three of nine parameters studied from tiredness. Concerning cancer, there are 5.6% of cancer cases in the study population, a percentage 10 times higher than that of the total Spanish population.

Discussion: People who are exposed to higher radiation values present more severe headaches, dizziness and nightmares. Moreover, they sleep fewer hours.

<https://pubmed.ncbi.nlm.nih.gov/33434609/>

Highlights

- People who are exposed to higher radiation values present more severe headaches, dizziness and nightmares.
- The methodology for obtaining electromagnetic radiation measurements should be reviewed.
- The population continues to receive radiation peaks in distances greater than 200 m, no one is free from exposure.

Conclusion

In conclusion, the data obtained shows that there is a relationship between the power density of radiation that a person receives at home every day and the presence of headaches, as well as the presence of sleep disorders.

People who receive higher doses of radiation sleep less hours and have nightmares at night. In addition, these people suffer from headaches with greater intensity and are more prone to dizziness. In this study, indicators like fainting episodes, presence of tachycardias or instability cannot be related. No conclusive results were found for fatigue, since, out of nine parameters studied, only a statistically significant relationship was found in three of them. The study of how electromagnetic fields affect health, should not only be done in relation to cancer, but also health indicators related to day to day. The methodology for obtaining electromagnetic radiation measurements should be reviewed, the averaged radiation measurements that are described in the CENELEC standard are not the most appropriate, they should be carried out in a narrow band and with maximum peak measurements.

The measured intensity depends fundamentally on the direction of the fundamental radiation beam and not so much on the distance to the antenna. In the beam direction, differences are found in the presence of pathologies with respect to distances, when these are greater than 200 m. Even at this distance, the population continues to receive radiation peaks, so that no one is free from exposure to these radiation sources.

The need for this study is related to the situation before 5G in terms of electromagnetic radiation rates. This study may be compared with the new radiant procedures that will be adopted in a short time.

ELECTROMAGNETIC POLLUTION ALERT: MICROWAVE RADIATION AND ABSORPTION IN HUMAN ORGANS AND TISSUES

Nantakan Wongkasem. Electromagnetic pollution alert: Microwave radiation and absorption in human organs and tissues. *Electromagn Biol Med.* 2021 Feb 10;1-18. doi: 10.1080/15368378.2021.1874976.

Abstract

Electromagnetic radiation from communication and electronic devices, networks, systems and base stations has drawn concern due to excessive global usage with increasing power and operating frequency level. Numerous previous researches only focus on how the radiation from certain frequency ranges of particular devices could harm specific human organs and tissues, resulting in distinct symptoms. In this research, electromagnetic propagation and properties in 14 human organs and tissues were analyzed and investigated based on the

organs and tissues' electromagnetic and mechanical parameters, and chemical composition. Counting the organs and tissues as electromagnetic materials, their permittivity and conductivity, computed by a 4-Cole-Cole mode, directly respective to the operating frequency, are interrelated to wave behavior and hence influence the organs' response. Tests were conducted in 1 GHz to 105 GHz system settings, covering most microwave frequency uses: 2.4 GHz of 4G-LTE, Wi-Fi, Bluetooth, ZigBee and the 5G ranges: 28 GHz of 5G-mmW and 95 GHz of 5G-IoT. Trial human organs and tissues were placed in the wave propagation direction of 2.4 GHz and 28 GHz dipole antennas, and a waveguide port operating from 95 to 105 GHz. The quantitative data on the effects of 5G penetration and dissipation within human tissues are presented. The absorbance in all organs and tissues is significantly higher as frequency increases. As the wave enters the organ-tissue model, the wavelength is shortened due to the high organ-tissue permittivity. Skin-Bone-Brain layer simulation results demonstrate that both electric and magnetic fields vanish before passing the brain layer at all three focal frequencies of 2.4 GHz, 28 GHz and 100 GHz.

<https://pubmed.ncbi.nlm.nih.gov/33566706/>

Excerpts

At a cellular tower or a cell site, an effective-radiated power (ERP) of up to 500 W per radio channel or transmitter, depending on the tower height, was permitted by the Federal Communications Commission (FCC) (Haripriya 2020b). The majority of cellular or Personal Communications Services (PCS) cell sites in urban and suburban areas operate at an ERP of 100 W per channel or less. An ERP of 100 W corresponds to an actual- radiated power of 5–10 W, depending on the type of antenna used (Stutzman and Thiele 2012). As with all forms of EM energy, the power density from a cellular or PCS transmitter decays proportionally to $1/R^2$ as the distance from the cell site increases. Therefore, for a common 40 W of the power density of about 10 mW/m² at 100 m from the antenna tower, the power density will be 1 mW/m², as the distance increases to 300 m away from the base station and the device will receive 1μW, based on the effective antenna area of 0.001 m². Having several mobile phones active in the same area, the total EM radiation can get close to 2 W, the same level as the maximum power radiation of 5 mW/cm² at 2 inches from the microwave oven surface permitted by federal regulations (Haripriya 2020c). Besides the aforementioned cellular emission, there is also EM radiation from wireless networks, smart and wearable devices, unceasingly scattering around, whether or not you are connected to their networks....

Conclusion

Eight human organ and tissue models (blood, brain, breast, eye bulb, heart, kidney, liver and lung), while tested with electromagnetic excitation from 2.4 GHz and 28 GHz dipole antennas, and 95–105 GHz waveguide ports, show that the wavelength of the E- and H-fields propagating inside the models is shortened or the wave frequency is accelerating, approximately 5–8 times. The lowest propagation frequency is found in the lung model, where the permittivity and conductivity are lowest among those of the eight studied models. The E-field intensity boosts significantly as the EM source frequency increases, approximately 10 times between 2.4 GHz and 28 GHz. The E-field intensity (from 95 GHz to 105 GHz radiation) rises from 3.89% (in the brain) to 4.86% (in the liver). The field intensity dramatically decreases as the wave propagates passing the model, indicating high wave attenuation from the material absorption.

The EM absorbance is found to be greater than 60% in all 14 organ and tissue models. At 100 GHz, the bone (83.01%), urinary bladder wall (82.95%), lung (80.02%), liver (68.18%), brain (66.18%) and eye bulb (64.46%) present the highest absorbance level. The absorbance increases significantly as the frequency increases, for instance, 43.78% at 2.4 GHz, 51.95% at 28 GHz and 66.16% at 100 GHz in the brain model. The transmittance in all models is below e-20 level. At all three operating frequencies: 2.4 GHz, 28 GHz and 100 GHz, the E- and H-fields vanish before passing through the brain layer of the Skin-Bone-Brain layered model. The overall layer absorbance is greatly elevated at all 3 frequencies: 69.18% at 2.4 GHz, 65.69% at 28 GHz and 71.49% at 100 GHz. The reflectance is around 30%, while the transmittance is significantly low. The scattering behavior and multiple absorbing possibility is observed. The absorbance is inversely proportional to the permittivity and conductivity of human organs and tissues. There is no clear correlation between the EM absorbance and heat transfer and generation rate. The models confirm that the higher the frequency, the greater the attenuation. Penetration is therefore reduced with increasing frequency. However, on the other hand, the absorption level increases with frequency. The results quantitatively validate the classic effects of the increase in frequency in terms of penetration and dissipation in human tissues and organs.

Further study of the EM radiation and propagation, either on specific complete human parts, for example, an eye bulb with cornea, anterior chamber, pupil, iris, lens, ciliary and vitreous body, macular, retina, choroid and optic nerve, or the whole human body, will definitely provide more insight and correlation of the EM radiation and its effects amid all human organs

EMPIRICAL AND MODELING APPROACH FOR ENVIRONMENTAL INDOOR RF-EMF ASSESSMENT IN COMPLEX HIGH-NODE DENSITY SCENARIOS: PUBLIC SHOPPING MALLS CASE STUDY

Mikel Celaya-Echarri, Leyre Azpilicueta, Victoria Ramos, Peio Lopez-Iturri, Francisco Falcone. Empirical and Modeling Approach for Environmental Indoor RF-EMF Assessment in Complex High-Node Density Scenarios: Public Shopping Malls Case Study. IEEE Access. 22 March 2021.

Abstract

This work provides an intensive and comprehensive in-depth study from an empirical and modeling approach of the environmental radiofrequency electromagnetic fields (RF-EMF) radiation exposure in public shopping malls, as an example of an indoor high-node user density context aware environment, where multiple wireless communication systems coexist. For that purpose, current personal mobile communications (2G-5G FR 1) as well as Wi-Fi services (IEEE 802.11n/ac) have been precisely analyzed in order to provide clear RF-EMF assessment insight and to verify compliance with established regulation limits. In this sense, a complete measurements campaign has been performed in different countries, with frequency-selective exposimeters (PEMs), providing real empirical datasets for statistical analysis and allowing discussion and comparison regarding current health effects and safety issues between some of the most common RF-EMF exposure safety standards: ICNIRP 2020 (Spain), IEEE 2019 (Mexico) and a more restrictive regulation (Poland). In addition, environmental RF-EMF exposure assessment simulation results, in terms of spatial E-field characterization and Cumulative Distribution Function (CDF) probabilities, have been provided for challenging incremental high-node user dense scenarios in worst case conditions, by means of a deterministic in-house 3D Ray-Launching (3D-RL) RF-EMF safety

simulation technique, showing good agreement with the experimental measurements. Finally, discussion highlighting the contribution and effects of the coexistence of multiple heterogeneous networks and services for the environmental RF-EMF radiation exposure assessment has been included, showing that for all measured results and simulated cases, the obtained E-Field levels are well below the exposure limits established in the internationally accepted standards and guidelines. In consequence, the obtained results and the presented methodology could become a starting point to establish the RF-EMF assessment basis of future complex heterogeneous 5G FR 2 developments on the millimeter wave (mmWave) frequency range, where massive high- node user density networks are expected.

Excerpt

It must be clearly remarked that, in the current state of knowledge, the results of scientific and epidemiological research have shown that in case of exposures at E-field levels below of far below the ICNIRP limits, there is no evidence of a relationship between exposure to radiofrequency fields produced by wireless technologies and potential adverse health effects. However, since the trend of 5G technology in towards the use of higher frequency ranges, it would be of great importance to carry out studies aimed at characterizing the effects of mmWave exposures, for which the information from scientific literature is not so complete and exhaustive as for the sub 6 GHz frequencies. This is a sort of knowledge gap that must be filled up to permit a specific and targeted surveillance held by the authorized health agencies and possibly an update of the guidelines/standards currently in use.

The proposed simulation methodology can be a useful and suitable technique to satisfactorily assess and verify in advance environmental RF-EMF exposure recommendations and limits to implement safe, efficient and reliable current and future wireless deployments for complex high-node density heterogeneous environments.

Open access paper: <https://ieeexplore.ieee.org/stamp/stamp.jsp?tp=&arnumber=9382976>

HEMATOBIOCHEMICAL AND HISTOPATHOLOGICAL ALTERATIONS OF KIDNEY AND TESTIS DUE TO EXPOSURE OF 4G CELL PHONE RADIATION IN MICE

Imam Hasan, Tanjina Amin, Md. Rafiqul Alam, Mohammad Rafiqul Islam. Available online 17 February 2021. <https://doi.org/10.1016/j.sjbs.2021.02.028>.

Abstract

The radiofrequency electromagnetic radiation emitted by smart phones on biological systems has wide media coverage and public concern in recent years. The aim of this study was to explore the effects of fourth- generation cell phone radiation exposure on hematological (Total leukocyte count, Total erythrocyte count, and hemoglobin %), biochemical (Serum creatinine) parameters, and histopathological changes in the kidney and testis of Swiss albino mice. A total of 30 male Swiss albino mice weighing 45–65 g was randomly divided into three groups (n = 10). The first group A was the control group, the second group B, was exposed to 40 minutes of mobile phone radiation daily, the third group C was exposed to 60 minutes of radiation daily from two 2400 Megahertz fourth-generation connected mobile phones for 60 days, respectively. The electromagnetic radiation frequency radiometer measured the frequency of electromagnetic radiation emitted from cell phones. The specific absorption rate was calculated as 0.087 W/kg. The control group was kept under similar conditions, but the electromagnetic field was not given for the same period. All the mice were sacrificed at the

end of the experiment. The blood samples were collected for hematobiochemical study, and then kidney and testis tissues were collected for histopathological study. Results of the study showed that the body weight and total erythrocyte count values were significantly ($p < 0.05$) decreased while total leukocyte count, hemoglobin %, and serum creatinine values were significantly ($p < 0.05$) increased in both the radiation exposure groups relative to the control group. Histopathological observation showed the kidney of 60 minutes exposed mice interstitial inflammation that causes marked mononuclear cellular infiltration compared to the 40 minutes and control mice. Compared to control mice, histopathological examinations of testicular tissue from the exposed mice, showed irregular in shapes and non-uniform sizes and fewer spermatogenic cells layer that leads to the larger lumen in the seminiferous tubules. It is concluded that fourth-generation cell phone radiation exposure may affect blood hemostasis and inflammation of mice's kidney and testis tissue. Based on these studies, it is important to increase public consciousness of potential adverse effects of mobile phone radiofrequency electromagnetic radiation exposure.

Open access paper: <https://www.sciencedirect.com/science/article/pii/S1319562X21001017>

THE EFFECT OF 4.5 G (LTE ADVANCED-PRO NETWORK) MOBILE PHONE RADIATION ON THE OPTIC NERVE

Erkin Özdemir, Ülkü Çömelekoglu, Evren Degirmenci, Gülsen Bayrak, Metin Yildirim, Tolgay Ergenoglu, Banu Coşkun Yılmaz, Begüm Korunur Engiz, Serap Yalin, Dilan Deniz Koyuncu, Erkan Ozbay. The effect of 4.5 G (LTE Advanced-Pro network) mobile phone radiation on the optic nerve. *Cutan Ocul Toxicol.* 2021 Mar 3;1-27. doi: 10.1080/15569527.2021.1895825.

Abstract

Purpose: Rapid development in mobile phone technologies increase the average mobile phone usage duration. This increase also triggers exposure to radiofrequency radiation (RF), which is a risk factor for the health. In this study, it was aimed to investigate the effect of mobile phone working with LTE-Advanced Pro (4.5G) mobile network on the optic nerve, which is responsible for the transmission of visual information.

Material and methods: Thirty-two rats divided into two groups as control (no RF, sham exposure) and experimental (RF exposure using a mobile phone with LTE-Advanced Pro network; 2 hours/day, 6 weeks). The visual evoked potential (VEP) was recorded and determined amplitudes and latencies of VEP waves. Optic nerve malondialdehyde level, catalase and superoxide dismutase activities were determined. Furthermore, ultrastructural and morphometric changes of optic nerve were evaluated.

Results: In VEP recordings, the mean VEP amplitudes of experimental group were significantly lower than control group. In ultrastructural evaluation, myelinated nerve fibers and glial cells were observed in normal histologic appearance both in sham and experimental group. However, by performing morphometric analysis, in the experimental group, axonal diameter and myelin thickness were shown to be lower and the G-ratio was higher than in the sham group. In the experimental group, malondialdehyde level was significantly higher and superoxide dismutase and catalase activities were significantly lower than sham group. There was a high correlation between VEP wave amplitudes and oxidative stress markers.

Conclusion: Findings obtained in this study support optic nerve damage. These results point out an important risk that may decrease the quality of life. <https://pubmed.ncbi.nlm.nih.gov/33653184/>

Excerpts

In recent years, everyone, from child to old, has a smartphone, and everyday a long time is passed looking at the screen of this phone. Comfort and efficiency achieved thanks to the high data transfer rate provided by LTE- Advanced Pro technology increase this time day by day. Eyes are the most affected body parts from this condition. In addition to the effects such as strabismus and eye impairment arising from looking at a small screen, it is also important to examine the hidden risks that the RF magnetic field created by the phone will cause on the eye. In this study, the effects of RF emission created by a LTE-Advanced Pro technology phone on the optic nerve were examined in all aspects and the findings were given in the previous section. Briefly it can be said that, for the first time in scientific literature, the findings of the present study indicate that the LTE- Advanced Pro mobile phone radiation causes significant damage by triggering oxidative stress in the optic nerve. LTE-Advanced Pro technology uses a wider RF band between 800 MHz and 2600 MHz and the network system selects the most appropriate band itself according to the user's requirements. It is known that penetration depth of RF increases with decreasing frequency [39]. Since effects of RF radiation were observed on the optic nerve which is behind the eye, it can be said that low frequency bands such as 800 MHz were mostly active during the experiments. Maybe this inference cannot be generalized for all communication purposes, but usage probability of low frequency bands during LTE-Advanced Pro smart phone usage will always keep the damage risk on optic nerve alive.

Fig. 3A and 3B show the distribution of electric field and SAR, respectively. As seen, maximum E field was 5.0 V/m (Fig. 3A) and maximum SAR (10 g) was 0.01 W/kg (Fig. 3B). The SAR value in the area of eyes was about 0.0035 W/kg (Fig. 3B).

Before and after exposure, body surface temperatures were 28.08 ± 0.19 and 28.07 ± 0.26 °C, respectively in the sham group. These values were 28.37 ± 0.29 and 28.39 ± 0.22 °C, respectively for the RF groups. There was no significant difference within sham ($p = 0.275$) and RF ($p = 0.120$) groups before and after exposure. Also, there was no significant differences in surface body temperature between sham and RF exposed groups before ($p = 0.142$) and after ($p = 0.321$) exposure.

In the present study, for the first time, it was shown that exposition to 4.5 G mobile phone radiation for 2 hours/day for 6 weeks causes optic nerve damage. The optic nerve transmits all visual information to the visual cortex, and any damage in this nerve can cause permanent and serious vision loss. This study demonstrated that RF exposure may be an environmental risk factor for eye toxicity and potential eye disorders. Further studies are needed to reveal the potentiality of the risk in this area.

POTENTIAL INFLUENCE OF PRENATAL 2.45 GHZ RADIOFREQUENCY ELECTROMAGNETIC FIELD EXPOSURE ON WISTAR ALBINO RAT TESTIS

Viera Almášiová, Katarína Holovská, Sandra Andrašková, Viera Cigánková, Zuzana Ševčíková, Adam Raček, Zuzana Andrejčáková, Katarína Beňová, Štefan Tóth, Eva Tvrdá, Ján Molnár, Enikő Račeková. Potential influence of prenatal 2.45 GHz radiofrequency electromagnetic field exposure on Wistar albino rat testis. *Histol Histopathol.* 2021 Mar 25;18331. doi: 10.14670/HH-18-331.

Abstract

An ever-increasing use of wireless devices over the last decades has forced scientists to clarify their impact on living systems. Since prenatal development is highly sensitive to numerous noxious agents, including radiation, we focused on the assessment of potential adverse effects of microwave radiation (MR) on testicular development. Pregnant Wistar albino rats (3 months old, weighing 282 ± 8 g) were exposed to pulsed MR at a frequency of 2.45 GHz, mean power density of 2.8 mW/cm^2 , and a specific absorption rate of 1.82 W/kg for 2 hours/day throughout pregnancy. Male offspring were no longer exposed to MR following birth. Samples of biological material were collected after reaching adulthood (75 days). In utero MR exposure caused degenerative changes in the testicular parenchyma of adult rats. The shape of the seminiferous tubules was irregular, germ cells were degenerated and often desquamated. The diameters of the seminiferous tubules and the height of the germinal epithelium were significantly decreased (both at $**p < 0.01$), while the interstitial space was significantly increased ($**p < 0.01$) when compared to the controls. In the group of rats prenatally exposed to MR, the somatic and germ cells were rich in vacuoles and their organelles were often altered.

Necrotizing cells were more frequent and empty spaces between Sertoli cells and germ cells were observed. The Leydig cells contained more lipid droplets. An increased Fluoro Jade - C and superoxide dismutase 2 positivity was detected in the rats exposed to MR. Our results confirmed adverse effects of MR on testicular development.

Open access paper: <https://www.hh.um.es/Articles-Proofs/18-331-manuscript.pdf>

PERSONAL EXPOSURE ASSESSMENT TO WI-FI RADIOFREQUENCY ELECTROMAGNETIC FIELDS IN MEXICAN MICROENVIRONMENTS

Raquel Ramirez-Vazquez, Jesus Gonzalez-Rubio, Isabel Escobar, Carmen Del Pilar Suarez Rodriguez, Enrique Arribas. Personal Exposure Assessment to Wi-Fi Radiofrequency Electromagnetic Fields in Mexican Microenvironments. *Int J Environ Res Public Health.* 2021 Feb 14;18(4):1857. doi: 10.3390/ijerph18041857

Abstract

In recent years, personal exposure to Radiofrequency Electromagnetic Fields (RF-EMF) has substantially increased, and most studies about RF-EMF with volunteers have been developed in Europe. To the best of our knowledge, this is the first study carried out in Mexico with personal exposimeters. The main objective was to measure personal exposure to RF-EMF from Wireless Fidelity or wireless Internet connection (Wi-Fi) frequency bands in Tamazunchale, San Luis Potosi, Mexico, to compare results with maximum levels permitted by international recommendations and to find if there are differences in the

microenvironments subject to measurements. The study was conducted with 63 volunteers in different microenvironments: home, workplace, outside, schools, travel, and shopping. The mean minimum values registered were 146.5 $\mu\text{W}/\text{m}^2$ in travel from the Wi-Fi 2G band and 116.8 $\mu\text{W}/\text{m}^2$ at home from the Wi-Fi 5G band, and the maximum values registered were 499.7 $\mu\text{W}/\text{m}^2$ and 264.9 $\mu\text{W}/\text{m}^2$ at the workplace for the Wi-Fi 2G band and the Wi-Fi 5G band, respectively. In addition, by time period and type of day, minimum values were registered at nighttime, these values being 129.4 $\mu\text{W}/\text{m}^2$ and 93.9 $\mu\text{W}/\text{m}^2$, and maximum values were registered in the daytime, these values being 303.1 $\mu\text{W}/\text{m}^2$ and 168.3 $\mu\text{W}/\text{m}^2$ for the Wi-Fi 2G and Wi-Fi 5G bands, respectively. In no case, values exceeded limits established by the International Commission on Non-Ionizing Radiation Protection (ICNIRP). Of the study participants ($n = 63$), a subgroup ($n = 35$) answered a survey on risk perception. According to these results, the Tamazunchale (Mexico) population is worried about this situation in comparison with several European cities; however, the risk perception changes when they are informed about the results for the study.

Open access paper: <https://pubmed.ncbi.nlm.nih.gov/33673014/>

RADIO-FREQUENCY ELECTROMAGNETIC FIELD EXPOSURE AND CONTRIBUTION OF SOURCES IN THE GENERAL POPULATION: AN ORGAN-SPECIFIC INTEGRATIVE EXPOSURE ASSESSMENT

Luuk van Wel, Ilaria Liorni, Anke Huss, Arno Thielens, Joe Wiart, Wout Joseph, Martin Rössli, Milena Foerster, Amelie Massardier-Pilonchery, Myles Capstick, Elisabeth Cardis, Roel Vermeulen. Radio-frequency electromagnetic field exposure and contribution of sources in the general population: an organ-specific integrative exposure assessment. *J Expo Sci Environ Epidemiol*. 2021 Mar 2. doi: 10.1038/s41370-021-00287-8.

Abstract

In order to achieve an integrated radio-frequency electromagnetic fields (RF-EMF) dose assessment, detailed information about source-specific exposure duration and output power is needed. We developed an Integrated Exposure Model (IEM) to combine energy absorbed due to use of and exposure to RF-EMF sources and applied it to a sample of the general population to derive population RF-EMF estimates. The IEM used specific absorption rate transfer algorithms to provide RF-EMF daily dose estimates ($\text{mJ}/\text{kg}/\text{day}$) using source-specific attributes (e.g. output power, distance), personal characteristics and usage patterns. Information was obtained from an international survey performed in four European countries with 1755 participants. We obtained median whole-body and whole-brain doses of 183.7 and 204.4 $\text{mJ}/\text{kg}/\text{day}$. Main contributors to whole-brain dose were mobile phone near the head for calling (2G networks) and far-field sources, whereas the latter together with multiple other RF-EMF sources were main contributors for whole-body dose. For other anatomical sites, 2G phone calls, mobile data and far-field exposure were important contributors. The IEM provides insight into main contributors to total RF-EMF dose and, applied to an international survey, provides an estimate of population RF-dose. The IEM can be used in future epidemiological studies, risk assessments and exposure reduction strategies.

<https://pubmed.ncbi.nlm.nih.gov/33654268/>

Excerpts

Relative contributions per source

For the whole-brain the use of mobile phones near the head remains by far the main contributor to total dose. Phone calls performed on a 2G network, which generally uses a higher output power, provide a high contribution followed at a distance by nearby Wi-Fi-routers. Regarding whole-body dose, the contribution of other sources becomes more important. Far-field exposure, Wi-Fi-routers, laptops, tablets and even other functions than calling on a mobile phone provide a higher contribution to the whole-body dose. This indicates that while just looking at mobile phone calls may include most RF-EMF exposure for health outcomes focused on the brain, this is not the case for the whole-body. In addition, adaptive output power control depending on the mobile phones' function may further influence exposure levels, as explained below. When looking at potential health endpoints at anatomical sites other than the brain (e.g. heart, liver), multiple devices should be included.

RF-EMF dose reduction

With the relative contributions found in this study, various non-technical interventions may be considered to reduce overall RF-EMF dose. The avoidance of using a mobile phone near the head when using 2G networks may be an efficient way to reduce overall exposure by half for the whole-brain and up to 25% for the whole-body.

This can be achieved on modern smartphones by disabling the use of 2G networks altogether or by using a wired hands-free kit instead. In the latter case, the exposure will shift from the head to other parts of the body when the device is held in a hand or pocket. In general, we observed a higher RF-EMF dose with device functions that require higher amounts of data, such as video streaming. Placing the device on a nearby surface or stand with data intensive uses can be considered to reduce dose. For far-field exposure, it is generally difficult to achieve individual reduction as these are continuous exposures generally not controlled by the subject, such as FM radio broadcast and mobile phone antennas.

Conclusion

In conclusion, we developed the most comprehensive RF-EMF dose estimation tool to date. Realistic population exposure scenarios were obtained by using data on mobile phone use from an international survey in the model. Overall RF-EMF dose for the whole-body and whole-brain was found to be higher in younger age groups in comparison with older groups. Mobile phone calls on 2G networks were found to be the main contributor to whole-brain RF-EMF dose. For whole-body dose, far-field of telecommunications and multiple other RF-EMF sources played a prominent role as well. These findings can be used in the creation of non-technical interventions aimed at lowering RF-EMF exposure from current technologies, with the modular structure of the model allowing inclusion of new technologies such as 5th generation networks. Future epidemiological studies involving RF-EMF exposure should take multiple RF-EMF sources into account by adding detailed questions on exposure duration when investigating other anatomical sites than the brain.

EVIDENCE OF BYSTANDER EFFECT INDUCED BY RADIOFREQUENCY RADIATION IN A HUMAN NEUROBLASTOMA CELL LINE

Olga Zeni, Stefania Romeo, Anna Sannino, Rosanna Palumbo, Maria Rosaria Scarfì. Evidence of bystander effect induced by radiofrequency radiation in a human neuroblastoma cell line. *Environ Res.* 2021 Feb 26;110935. doi: 10.1016/j.envres.2021.110935.

Abstract

In previous studies we demonstrated that radiofrequency (RF) electromagnetic fields (EMF) is able to reduce DNA damage induced by a subsequent treatment with genotoxic agents, resembling the adaptive response, a phenomenon well known in radiobiology. In this study we report on the capability of the culture medium from SH-SY5Y neuroblastoma cells exposed to 1950 MHz to elicit, in recipient non-exposed cells, a reduction of menadione-induced DNA damage ($P < 0.05$; comet assay), indicating the capability of non-ionizing radiation to elicit a bystander effect. A comparable reduction was also detected in cultures directly exposed to the same EMF conditions ($P < 0.05$), confirming the adaptive response. In the same exposure conditions, we also evidenced an increase of heat shock protein 70 (hsp70) in culture medium of cells exposed to RF with respect to sham exposed ones (western blot analysis; $P < 0.05$), while no differences were detected in the intracellular content of hsp70. On the whole, our results evidence a protective effect of RF against menadione-induced DNA damage in directly and non-directly exposed cells, and suggest hsp70 pathway to be investigated as one of the potential candidate underpinning the interaction between RF exposure and biological systems. <https://pubmed.ncbi.nlm.nih.gov/33647301/>

THE EFFECT OF PULSED ELECTROMAGNETIC FIELDS ON ANGIOGENESIS

Lihong Peng, Chenying Fu, Lu Wang, Qing Zhang, Zejun Liang, Chengqi He, Quan Wei. The Effect of Pulsed Electromagnetic Fields on Angiogenesis. *Bioelectromagnetics.* 2021 Mar 6. doi: 10.1002/bem.22330.

Abstract

A pulsed electromagnetic field (PEMF) has been used to treat inflammation-based diseases such as osteoporosis, neurological injury, and osteoarthritis. Numerous animal experiments and in vitro studies have shown that PEMF may affect angiogenesis. For ischemic diseases, in theory, blood flow may be richer by increasing the number of blood vessels which supply blood to ischemic tissue. PEMF plays a role in enhancing angiogenesis, and their clinical application may go far beyond the current scope. In this review, we analyzed and summarized the effects and possible mechanisms of PEMF on angiogenesis. Most studies have shown that PEMF with specific parameters can promote angiogenesis, which is manifested by an increased vascular growth rate and increased capillary density. The potential mechanisms consist of promoting vascular endothelial cell proliferation, migration, and tube formation, and increasing the expression level of vascular endothelial growth factor (VEGF), fibroblast growth factor 2 (FGF2), angiopoietin-2 (Ang-2), and other angiogenic growth factors. Additionally, PEMF has an impact on the activation of voltage-gated calcium channels (VGCC). <https://pubmed.ncbi.nlm.nih.gov/33675261/>

EFFECTS OF RADIOFREQUENCY ELECTROMAGNETIC RADIATION EMITTED FROM A MOBILE PHONE BASE STATION ON THE REDOX HOMEOSTASIS IN DIFFERENT ORGANS OF SWISS ALBINO MICE

Mary Zosangzuali, Marina Lalremruati, C Lalmuansangi, F Nghakliana, Lalrinthara Pachuau, Priyanka Bandara, Zothan Siama. Effects of radiofrequency electromagnetic radiation emitted from a mobile phone base station on the redox homeostasis in different organs of Swiss albino mice. *Electromagn Biol Med*. 2021 Mar 9;1-15. doi: 10.1080/15368378.2021.1895207.

Abstract

This study was designed to investigate the possible effects of exposure to mobile phone base station (MPBS) emits 1800-MHz RF-EMR on some oxidative stress parameters in the brain, heart, kidney and liver of Swiss albino mice under exposures below thermal levels. Mice were randomly assigned to three experimental groups which were exposed to RF-EMR for 6 hr/day, 12 hr/day and 24 hr/day for 45 consecutive days, respectively, and a control group. The glutathione (GSH) levels and activities of glutathione-s-transferase (GST) and superoxide dismutase (SOD) were significantly reduced in mice brain after exposure to RF-EMR for 12 hr and 24 hr per day. Exposure of mice to RF-EMR for 12 hr and 24 hr per day also led to a significant increase in malondialdehyde (an index of lipid peroxidation) levels in mice brain. On the contrary, exposures used in this study did not induce any significant change in various oxidative stress-related parameters in the heart, kidney and liver of mice. Our findings showed no significant variations in the activities of aspartate amino-transferase (AST), alanine amino-transferase (ALT), and on the level of creatinine (CRE) in the exposed mice. This study also revealed a decrease in RBC count with an increase in WBC count in mice subjected to 12 hr/day and 24 hr/day exposures. Exposure to RF-EMR from MPBS may cause adverse effects in mice brain by inducing oxidative stress arising from the generation of reactive oxygen species (ROS) as indicated by enhanced lipid peroxidation, and reduced levels and activities of antioxidants. <https://pubmed.ncbi.nlm.nih.gov/33687298/>

EFFECTS OF 1.5-GHZ HIGH-POWER MICROWAVE EXPOSURE ON THE REPRODUCTIVE SYSTEMS OF MALE MICE

Guofu Dong, Hongmei Zhou, Yan Gao, Xuelong Zhao, Qi Liu, Zhihui Li, Xi Zhao, Jiye Yin, Changzhen Wang. Effects of 1.5-GHz high-power microwave exposure on the reproductive systems of male mice. *Electromagn Biol Med*. 2021 Mar 10;1-10. doi: 10.1080/15368378.2021.1891091.

Abstract

High-power microwaves (HPMs) have been reported to have hazardous effects on multiple human and animal organs. However, the biological effects of 1.5-GHz HPMs on the reproductive system are not clear. Here, we studied the effects of 1.5 -GHz HPM whole-body exposure on the pathological structure of the testicles and changes in spermatozoa mobility. C57BL/6 mice of groups L, M, and H were exposed to 1.5-GHz HPM fields for two 15-min intervals at the average specific absorption rates of 3, 6, and 12 W/Kg, respectively. The pathological structure of the testicles and spermatozoa, as well as serum testosterone and sperm motility parameters, were evaluated at 6 h, 1 d, 3 d, and 7 d after exposure. As a result, there were no significant pathological or ultrastructural changes in the testicles or spermatozoa and serum testosterone levels. The number of progressively motile spermatozoa, curvilinear velocity, linear velocity, and average path velocity of the exposure

group increased at 6 h, decreased at 1 d, and recovered at 3 d. The opposite results were considered a stress response to the thermal effect of the microwaves. Our results indicated that 1.5-GHz HPM whole-body exposure in mice at SARs of 3, 6, and 12 W/Kg for 30 min did not cause obvious damage to the reproductive system.
<https://pubmed.ncbi.nlm.nih.gov/33688776/>

DIRECTIONAL MIGRATION OF BREAST CANCER CELLS HINDERED BY INDUCED ELECTRIC FIELDS MAY BE DUE TO ACCOMPANYING ALTERATION OF METABOLIC ACTIVITY

Travis H. Jones, Kirti Kaul, Ayush A. Garg, Jonathan W. Song, Ramesh K. Ganju, Vish V. Subramaniam. Directional Migration of Breast Cancer Cells Hindered by Induced Electric Fields May Be Due to Accompanying Alteration of Metabolic Activity. *Bioelectricity*. 20 Jan 2021. <https://doi.org/10.1089/bioe.2020.0048>

Abstract

Background: Induced electric fields (iEFs) control directional breast cancer cell migration. While the connection between migration and metabolism is appreciated in the context of cancer and metastasis, effects of iEFs on metabolic pathways especially as they relate to migration, remain unexplored.

Materials and Methods: Quantitative cell migration data in the presence and absence of an epidermal growth factor (EGF) gradient in the microfluidic bidirectional microtrack assay was retrospectively analyzed for additional effects of iEFs on cell motility and directionality. Surrogate markers of oxidative phosphorylation (succinate dehydrogenase [SDH] activity) and glycolysis (lactate dehydrogenase activity) were assessed in MDA- MB-231 breast cancer cells and normal MCF10A mammary epithelial cells exposed to iEFs and EGF.

Results: Retrospective analysis of migration results suggests that iEFs increase forward cell migration speeds while extending the time cells spend migrating slowly in the reverse direction or remaining stationary. Furthermore, in the presence of EGF, iEFs differentially altered flux through oxidative phosphorylation in MDA- MB-231 cells and glycolysis in MCF10A cells.

Conclusions: iEFs interfere with MDA-MB-231 cell migration, potentially, by altering mitochondrial metabolism, observed as an inhibition of SDH activity in the presence of EGF. The energy intensive process of migration in these highly metastatic breast cancer cells may be hindered by iEFs, thus, through hampering of oxidative phosphorylation.

Conclusion

A retrospective analysis of the momentary speeds and velocities (forward and reverse directions) reported in Garg et al.,⁹ revealed that cancer cell migration is hindered by iEFs because migrating cells slow down or are immobile some of the time in the 700 μ m long microchannel over the duration of the experiment. Thus, iEFs not only reduce the number of cancer cells migrated but, on average, also slow them down. We have separately shown that SDH activity is hindered in MDA-MB-231 cells and LDH activity is enhanced in MCF10A cells by iEFs in the presence of EGF. This additional data on the effects of iEF on cellular metabolism may explain the previously reported changes in migration characteristics in microtrack microchannels. This work has paved the way for better understanding of how iEFs interact

with MDA-MB-231 cells and points to possible new therapeutic options for treating metastasis that involve targeting specific metabolic pathways. <https://www.liebertpub.com/doi/10.1089/bioe.2020.0048>

HEART RATE VARIABILITY AND MAGNETIC FIELD EXPOSURE AMONG TRAIN ENGINE DRIVERS-A PILOT STUDY

Kjell Hansson Mild, Roland Bergling, Rolf Hörnsten. Heart Rate Variability and Magnetic Field Exposure Among Train Engine Drivers-A Pilot Study. *Bioelectromagnetics*. 2021 Mar 1. doi: 10.1002/bem.22329.

No abstract. <https://pubmed.ncbi.nlm.nih.gov/33644873/>

EFFECTS INDUCED BY A WEAK STATIC MAGNETIC FIELD OF DIFFERENT INTENSITIES ON HT-1080 FIBROSARCOMA CELLS

Hakki Gurhan, Rodolfo Bruzon, Sahithi Kandala, Ben Greenebaum, Frank Barnes. Effects Induced by a Weak Static Magnetic Field of Different Intensities on HT-1080 Fibrosarcoma Cells. *Bioelectromagnetics*. 2021 Mar 18. doi: 10.1002/bem.22332.

Abstract

In this study, we investigated the effects of weak static magnetic fields (SMFs) on HT-1080 human fibrosarcoma cells. Exposures to SMFs for four consecutive days were varied from 0.5 to 600 μ T for treated units, while exposures to control units were held at 45 μ T. Growth rates were measured by comparing cell counts, whereas membrane potentials, mitochondrial calcium, mitochondrial superoxide (O_2^-), nitric oxide (NO), hydrogen peroxide (H_2O_2), intercellular pH, and oxidative stress were measured by using fluorescent dyes. The relative cell growth rates vary with the angle of the SMFs. Increases in the magnitude of the SMFs increased concentrations of mitochondrial calcium and membrane potential and decreased intracellular pH. H_2O_2 , an important reactive oxygen species (ROS), increases at 100 and 200 μ T, decreases at 300 and 400 μ T and increases again at 500 and 600 μ T. Overall, oxidative stress increases slightly with increasing SMFs, while superoxide and NO concentrations decrease. These results indicate that weak SMFs can accelerate and inhibit cell growth rates and induce alterations in ROS. Changes in ROS and oxidative stress are important for various cell functions. Calcium influx into mitochondria was one of the initial steps into the corresponding changes. <https://pubmed.ncbi.nlm.nih.gov/33735454/>

Excerpts

In the present study, we looked at mitochondrial ROS generation as an important source for ROS signaling and concentrated our attention on the initial parts of the mitochondrial signaling pathways. Mitochondrial superoxide, H_2O_2 , and NO are considered important intra-mitochondrial signaling molecules. In our experiments, ROS responses to the SMF were more complex. Changes in H_2O_2 are not linearly correlated with cell growth. This might be due to increases in antioxidant concentrations [Rosenspire et al., 2005]. Decreases in H_2O_2 correspond to where we observed the highest increase in cell growth rates. There are various studies showing that low levels of H_2O_2 can initiate cell proliferation [Antunes and Brito, 2017; Sies, 2017]. The mitochondrial superoxide and the NO tend to decrease with the increase in MF. The mitochondrial superoxide decreased with respect to the control for all SMF values, being highly significant for 200 and 400 μ T, and the NO decreased with respect

to the control for 0.5, 100, 400, and 600 μT . H_2O_2 and oxidative stress tend to increase with the increase in SMF. In both parameters, a highly significant increase was observed for 100 and 200 μT .

The importance of these results is that variations of background MFs can both increase and decrease cell growth rates and corresponding concentrations of ROS and other signaling molecules. These observed changes are significant as different concentration levels of ROS could have beneficial or adverse effects on biology. In conclusion, the basic effect observed here is that changes in SMFs create a change in the growth rate of the cells and changes in some important molecular concentrations on both sides of the cell and mitochondrial membrane. Though the growth rate is one of the parameters that we have studied, the changes in the signaling molecules and other cell parameters might cause long-term effects in biological systems. An example of this is long-term changes in oxidative stress related to cancer and aging-related diseases. Canceling out the earth's MF substantially, down to 0.5 μT by Mu metal cages, not only gave us the capability to eliminate background noise but also allowed us to observe effects at a level much lower than the earth's MF. Thus, in this study, we believe that measuring and controlling the SMFs and eliminating the effects of temperature and background noise in experiments are important for attempting to understand the effects of electromagnetic fields on biological systems. Additionally, we believe that the amplitude, angle of incidence, and length of the exposure can lead to variable results and need to be included in the description of experimental protocols [Barnes and Greenebaum, 2015].

Future work will include time-varying MFs, where we expect to see both electric and MF effects. Time-varying MFs induce electric fields; other data indicate that both electric and MFs can induce changes in cell growth rates and other biological parameters [Bingham, 1996]. More research work will be required to determine the mechanisms by which these two different fields modify the behavior of cells.

NO EVIDENCE FOR GENOTOXICITY IN MICE DUE TO EXPOSURE TO INTERMEDIATE-FREQUENCY MAGNETIC FIELDS USED FOR WIRELESS POWER-TRANSFER SYSTEMS

Shin Ohtani, Akira Ushiyama, Keiji Wada, Yukihisa Suzuki, Kazuyuki Ishii, Kenji Hattori. No evidence for genotoxicity in mice due to exposure to intermediate-frequency magnetic fields used for wireless power-transfer systems. *Mutat Res.* Mar-Apr 2021;863-864:503310. doi: 10.1016/j.mrgentox.2021.503310.

Abstract

Time varying magnetic fields (MFs) are used for the wireless power-transfer (WPT) technology. Especially, 85 kHz band MFs, which are included in the intermediate frequency (IF) band (300 Hz - 10 MHz), are commonly used WPT system for charging electric vehicles. Those applications of WPT technology have elicited public concern about health effects of IF-MF. However, existing data from health risk assessments are insufficient and additional data are needed. We assessed the genotoxic effects of IF-MF exposure on erythroid differentiation in mice. A high-intensity IF-MF mouse exposure system was constructed to induce an average whole-body electric field of 54.1 V/m. Blood samples were obtained from male mice before and after a 2-week IF-MF exposure (1 h/day, total: 10 h); X-irradiated mice were used as positive controls. We analyzed the blood samples with the micronucleus (MN) test and the

Pig-a mutation assay. No significant differences were seen between IF-MF- exposed and sham-exposed mice in the frequencies of either MN or Pig-a mutations in mature erythrocytes and reticulocytes. IF-MF exposure did not induce genotoxicity in vivo under the study conditions (2.36× the basic restriction for occupational exposure, 22.9 V/m, in the International Commission on Non-Ionizing Radiation Protection (ICNIRP) guidelines). The absence of significant biological effects due to IF-MF exposure supports the practical application of this technology.

Highlights

- We examined the genotoxicity of IF-MF exposure >2× the ICNIRP guideline.
- IF-MF had no genotoxic effects in an in vivo micronucleus test and Pig-a assay.
- These observations are useful for assessing the safety of a WPT system.
<https://pubmed.ncbi.nlm.nih.gov/33678242/>

OCCUPATIONAL EXPOSURE TO EXTREMELY LOW-FREQUENCY MAGNETIC FIELDS AND RISK OF AMYOTROPHIC LATERAL SCLEROSIS: RESULTS OF A FEASIBILITY STUDY FOR A POOLED ANALYSIS OF ORIGINAL DATA

Dan Baaken, Dagmar Dechent, Maria Blettner, Sarah Drießen, Hiltrud Merzenich. Occupational Exposure to Extremely Low-Frequency Magnetic Fields and Risk of Amyotrophic Lateral Sclerosis: Results of a Feasibility Study for a Pooled Analysis of Original Data. Bioelectromagnetics. 2021 Mar 25. doi: 10.1002/bem.22335.

Abstract

Previous meta-analyses have suggested an increased risk of amyotrophic lateral sclerosis (ALS) associated with occupational exposure to extremely low-frequency magnetic fields (ELF-MF). However, results should be interpreted with caution since studies were methodologically heterogeneous. Here, we assessed the feasibility of a pooling study to harmonize and re-analyze available original data. A systematic literature search was conducted. Published epidemiological studies were identified in PubMed and EMF-Portal from literature databases' inception dates until January 2019. The characteristics of all studies were described, including exposure metrics, exposure categories, and confounders. A survey among the principal investigators (PI) was carried out to assess their willingness to provide their original data. The statistical power of a pooling study was evaluated. We identified 15 articles published between 1997 and 2019. Studies differed in terms of outcome, study population, exposure assessment, and exposure metrics. Most studies assessed ELF-MF as average magnetic flux density per working day; however, exposure categories varied widely. The pattern of adjustment for confounders was heterogeneous between studies, with age, sex, and socioeconomic status being most frequent. Eight PI expressed their willingness to provide original data. A relative risk of ≥ 1.14 for ALS and occupational exposure to ELF-MF can be detected with a power of more than 80% in a pooled study. The pooling of original data is recommended and could contribute to a better understanding of ELF-MF in the etiology of ALS based on a large database and reduced heterogeneity due to a standardized analysis protocol with harmonized exposure metrics and exposure categories.

Open access paper: <https://onlinelibrary.wiley.com/doi/10.1002/bem.22335>

THE INFLUENCE OF THE EXTREMELY LOW FREQUENCY ELECTROMAGNETIC FIELD ON CLEAR CELL RENAL CARCINOMA

Cios A, Ciepielak M, Stankiewicz W, Szymański Ł. The Influence of the Extremely Low Frequency Electromagnetic Field on Clear Cell Renal Carcinoma. *International Journal of Molecular Sciences*. 2021; 22(3):1342. <https://doi.org/10.3390/ijms22031342>.

Abstract

The development of new technologies and industry is conducive to the increase in the number and variety of electromagnetic field (EMF) sources in our environment. The main sources of EMF are high-voltage lines, household appliances, audio/video devices, mobile phones, radio stations, and radar devices. In the growing use of electronic devices, scientists are increasingly interested in the effects of EMF on human health. Even though many studies on the effects of EMF have already been carried out, none of them has shown a significant effect on mammals, including humans. Moreover, it is not entirely clear how EMF influences cell behavior. The International Agency for Research on Cancer on 31 May 2011, classified PEM as a possible carcinogenic factor. This study aimed to investigate the effect of the electromagnetic field on morphological and functional changes in clear cell renal carcinoma. The research was carried out on in vitro cultures of four cell lines: HEK293, 786-O 769-P, and Caki1. The results of the research showed that the EMF of low frequency had a slight effect on the viability of cells. EMF, which induced cell arrest in the G1 phase, increased the number of early apoptotic cells and decreased the number of viable cells in the 786-O line. EMF did not affect the proliferation and viability of HEK293 cells. Extreme low-frequency EMF (ELF-EMF) also showed an inhibitory effect on the migration and metastatic properties of clear cell kidney cancer cells. Moreover, shortly after the end of ELF-EMF exposure, significant increases in ROS levels were observed in all tested cell lines. As part of the work, it was shown that low-frequency EMF shows an inhibitory effect on the proliferation of primary cancer cells, diminishing their migratory, invasive, and metastatic abilities. It also increases the apoptosis of cancer cells and the amount of reactive oxygen species. Based on the results of our research, we want to point up that the effect of ELF-EMF depends on a specific metabolic state or at a specific stage in the cell cycle of the cells under study

Open access paper: <https://www.mdpi.com/1422-0067/22/3/1342/htm>

INTERFERENCE OF CARDIOVASCULAR IMPLANTABLE ELECTRONIC DEVICES BY STATIC ELECTRIC AND MAGNETIC FIELDS

Kai Jagielski, Thomas Kraus, Dominik Stunder. Interference of Cardiovascular Implantable Electronic Devices by Static Electric and Magnetic Fields. *Expert Rev Med Devices*. 2021 Mar 12. doi: 10.1080/17434440.2021.1902802.

Abstract

Introduction: Electromagnetic interference (EMI) of cardiovascular implantable electronic devices (CIED) can lead to malfunctions and pose a danger for implant carriers. The increased use of DC technologies, e.g., in electric mobility, creates more static fields representing an increasing hazard for implant carriers.

Areas covered: A combination of approaches was used to determine thresholds for EMI by static fields. A literature search was conducted to identify relevant EMI mechanisms and to extract possible thresholds. The literature search revealed four interference mechanisms

caused by static magnetic fields and none for static electric fields. Due to the scarce information on motion-induced EMI, numerical simulations were performed to obtain a threshold. The simulation results were evaluated using medical product standards and benchmark tests on commercially available CIEDs. The results show that motion-induced interference should not occur below the activation of the magnetic safety switch (reed switch or Hall-effect sensor, MSS).

Expert opinion: The determined threshold for motion-induced EMI at 24.8 mT shows that the MSS activation is still the most relevant mechanism that can occur at 0.8 mT. Limit values for the general population do not protect implant carriers from EMI.
<https://pubmed.ncbi.nlm.nih.gov/33710950/>

Conclusion

The aim of this study was the assessment of risks for CIED patients by SEF and SMF. Regarding SEF, no indication was found that CIED can be interfered with when implant wearers are exposed to SEF. This does not include potential EMI from contact currents as they may occur from charged objects in the vicinity of SEF. Contact currents were not the focus of this study due to the significantly different scenario between field exposure and contact current, in which skin resistance and voltage level play a decisive role. Under the influence of SEF exposure, the shielding of the human body as a Faraday cage is sufficient to prevent electric induction into CIED that are located entirely under the human skin.

Regarding SMF, four interference mechanisms have been identified. As a reed switch or Hall effect sensor, the unintended activation of the MSS is the most investigated effect and can occur at SMF greater than 0.8 mT. Magnetic forces on CIED may be relevant for strong magnetic fields in the magnitude of teslas as they arise from MRI devices. A parameter reset or reprogramming of the CIED by SMF seems to be relatively uncommon, and only one publication reported a reproducible case with a PM by earphones. Rather few publications treat motion-induced EMI with the iEGM of the CIED, which has been investigated with numerical simulations and benchmark tests in this study.

In this study, activation thresholds for motion-induced EMI were determined and published for the first time. However, an estimated occurrence starting at 24.8 mT shows that motion-induced EMIs are less relevant than the activation of MSS that can already occur at 0.8 mT. However, motion-induced EMIs still pose a potential risk to implant wearers because the CIED may not provide the necessary therapy.

Other DC technologies and SMF sources will likely emerge in the future, both in professional and private environments. As a result, EMI from static fields will continue to be significant. The question arises whether the triggering of MSS around 1 mT is still practical, where even today, many technical applications already generate static magnetic fields of more than 1 mT, or whether there is a need for innovation on the part of the manufacturers.

SWIMMING DIRECTION OF THE GLASS CATFISH IS RESPONSIVE TO MAGNETIC STIMULATION

Ryan D Hunt, Ryan C Ashbaugh, Mark Reimers, Lalita Udpa, Gabriela Saldana De Jimenez, Michael Moore, Assaf A Gilad, Galit Pelled. Swimming direction of the glass catfish is responsive to magnetic stimulation. *PLoS One*. 2021 Mar 5;16(3):e0248141. doi: 10.1371/journal.pone.0248141.

Abstract

Several marine species have developed a magnetic perception that is essential for navigation and detection of prey and predators. One of these species is the transparent glass catfish that contains an ampullary organ dedicated to sense magnetic fields. Here we examine the behavior of the glass catfish in response to static magnetic fields which will provide valuable insight on function of this magnetic response. By utilizing state of the art animal tracking software and artificial intelligence approaches, we quantified the effects of magnetic fields on the swimming direction of glass catfish. The results demonstrate that glass catfish placed in a radial arm maze, consistently swim away from magnetic fields over 20 μ T and show adaptability to changing magnetic field direction and location.

Open access paper: <https://pubmed.ncbi.nlm.nih.gov/33667278/>

LOST OPPORTUNITIES FOR CANCER PREVENTION: HISTORICAL EVIDENCE ON EARLY WARNINGS WITH EMPHASIS ON RADIOFREQUENCY RADIATION

Lennart Hardell and Michael Carlberg. Lost opportunities for cancer prevention: historical evidence on early warnings with emphasis on radiofrequency radiation. *Reviews on Environmental Health*. 2021. doi: <https://doi.org/10.1515/reveh-2020-0168>.

Abstract

Some historical aspects on late lessons from early warnings on cancer risks with lost time for prevention are discussed. One current example is the cancer-causing effect from radiofrequency (RF) radiation. Studies since decades have shown increased human cancer risk. The fifth generation, 5G, for wireless communication is about to be implemented world-wide despite no comprehensive investigations of potential risks to human health and the environment. This has created debate on this technology among concerned people in many countries. In an appeal to EU in September 2017, currently endorsed by more than 400 scientists and medical doctors, a moratorium on the 5G deployment was required until proper scientific evaluation of negative consequences has been made (www.5Gappeal.eu). That request has not been taken seriously by EU. Lack of proper unbiased risk evaluation of the 5G technology makes adverse effects impossible to be foreseen. This disregard is exemplified by the recent report from the International Commission on non-ionizing radiation protection (ICNIRP) whereby only thermal (heating) effects from RF radiation are acknowledged despite a large number of reported non-thermal effects. Thus, no health effects are acknowledged by ICNIRP for non-thermal RF electromagnetic fields in the range of 100 kHz–300 GHz. Based on results in three case-control studies on use of wireless phones we present preventable fraction for brain tumors. Numbers of brain tumors of not defined type were found to increase in Sweden, especially in the age group 20–39 years in both genders, based on the Swedish Inpatient Register. This may be caused by the high prevalence of wireless phone use among children and in adolescence taking a reasonable latency period and the higher vulnerability to RF radiation among young persons.

Radiofrequency radiation (Excerpts)

In 2011 radiofrequency electromagnetic fields (RF-EMF) in the frequency range 30 kHz–300 GHz were evaluated by IARC at WHO to be possibly carcinogenic to humans, Group 2B [62], [, 63]. This was based on evidence of increased risk for glioma and acoustic neuroma in human epidemiology studies on use of mobile and/or cordless phone (DECT) [64], [65], [66], [67], [68], [69]. The increased cancer risk was supported by laboratory studies [70],[, 71].

Extremely low frequency (ELF)-EMF was in 2001 evaluated by IARC to be a possible human carcinogen, Group 2B [72]. This was the first time that non-ionizing radiation at low intensity levels can be a possible cause of cancer. It predated the IARC finding for RF-EMF by a decade.

Since then the evidence on RF-EMF carcinogenesis has strengthened based on further human studies on use of wireless phones, as reviewed [73], [, 74]. Also animal studies show increased cancer risk, both near field RF-EMF exposure [75], [76], [77] and far field exposure [78], [, 79]. Mechanistic studies show increase of reactive oxygen species (ROS) [80] as well as DNA damage [81]. These results give support to the increased cancer risk in humans and laboratory tested animals for RF radiation. In fact, RF-EMF may now be classified as a human carcinogen, Group 1 [82], [, 83]. However, such classification can only be made by IARC.

Of course, these well documented health hazards from RF-EMF are not well accepted by the telecom industry and its allied experts. Several methods are used to create doubt. Studies are discredited, only partly cited, or even not cited at all [84], [85], [86]. Thereby the uniformed reader gets the wrong information on actual risks. This includes also regulatory agencies and policy makers. Even agencies aimed at setting exposure guidelines may include pro-industry and biased scientists that obscure the true risks [87], [, 88].

ICNIRP

The International Commission on Non-Ionizing Radiation Protection (ICNIRP) is a private non-governmental (NGO) organization registered in Munich, Germany. ICNIRP appoints its own members and is closed to transparency. It was started in 1992 with the biophysicist Michael Repacholi as the first chairman, now emeritus member. ICNIRP has published three articles with guidelines on RF-EMF exposure [86], [89], [90]. Only thermal (heating) effects from RF radiation are recognized, thereby excluding all studies showing harmful effects at lower non-thermal intensities. In contrast to ICNIRP, some other expert panels such as European Academy of Environmental Medicine [91], the Bioinitiative group [92], and the Russian Commission for Protection from Non- Ionizing Radiation [93], take into account non-thermal RF effects and suggest much lower guidelines for RF exposure.

ICNIRP has managed to get collaborative status with WHO, as discussed previously [88]. The aim is to harmonize the RF-radiation guidelines all over the world. For that purpose ICNIRP has been successful. The guidelines are set to allow very high exposure levels so that the deployment of this technology is not hampered, in favor of industry but at disadvantage to human health and environment. In fact, the ICNIRP guidelines have never been challenged by industry in peer-reviewed articles, which must be taken as a green card for acceptance by industry.

Attributable fraction

The attributable fraction (AF), sometimes also called the etiologic fraction, is the number of cases in which exposure played an etiologic role. This is the preventable fraction if exposure would not be present. In Belpomme et al. [73] we published meta-analyses for longest cumulative use of mobile phones with odds ratio (OR) and 95 % confidence interval (CI), both for total and for ipsilateral wireless phone use. Note that only the Hardell group assessed also use of cordless phones (DECT). We present here AF based on statistically significant increased risks in the meta-analyses. AF is the proportion of cases that can be attributed to the particular exposure. This is calculated as the exposed case fraction multiplied by $[(OR-1)/OR]$.

As displayed in Table 1 the AF for glioma was calculated to 4.88%, 95% CI = 2.44–6.57%, corresponding to 211 preventable cases, 95% CI = 105–284 cases in the longest time for all cumulative use of wireless phones. Regarding ipsilateral use of the wireless phone AF was 6.03%, 95% CI = 4.51–7.12%, yielding 150 cases; 95% CI 112–177 to be preventable.

Rates of brain tumors in the Swedish National Inpatient Register ICD-code D43

Rates of brain tumors of unknown type, D43, were studied using the Swedish Inpatient Register (IPR) without any personal identification information [94]. It was established in 1964 and has complete national coverage since 1987 [95]. Register data on D43 are available from 1998. Currently more than 99% of hospital discharges are registered. For outpatients the data are less reliable due to missing information. The reporting of outpatients has increased during more recent years so these time trends may give spurious results, thus we omitted outpatients from the analysis.

Data were analyzed for the time period 1998–2019. Age-standardized rates are not available in the register. Instead numbers of patients per 100,000 inhabitants are reported. The Joinpoint Regression Analysis program version 4.1.1.1 was used to examine numbers of patients per 100,000 in inpatient care and incidence per 100,000 person-years in the Swedish Inpatient Register, by fitting a model of 0–3 joinpoints using permutation tests with Bonferroni correction for multiple testing to calculate the number of joinpoints that best fits the material [96]. When joinpoints were detected annual percentage changes (APC) and 95% CIs were calculated for each linear segment. Average annual percentage changes (AAPC) were also calculated for the whole time period using the average of the APCs weighted by the length of the segment. To be able to calculate APC and AAPC the data was log-transformed prior to analysis. Thus, it was not possible to perform joinpoint regression analysis when there were years with no cases during that time period. Since the data do not include any personal identification no ethical approval was needed.

In men AAPC increased during 1998–2019 with +1.77%, 95% confidence interval (CI) –0.02, +3.58%, Table 2; Figure 1. The increase was highest in the age group 20–39 years, +2.90%, 95% CI +1.66, +4.16 %, Figure 2. AAPC increased statistically significant in all age groups, except 0–19 years.

Similar results were found in women with AAPC +1.70%, 95% CI +0.38, +3.05% during 1998–2019, Table 3; Figure 3. Also in women the highest increase of AAPC was found in the age group 20–39 years, +2.89%, 95% CI + 1.54, +4.27%, Figure 4. AAPC increased statistically significant in all age groups except 0–19 years and 80+ years. Especially high increase of APC

was seen in women aged 60–79 years during 2005–2019, and women aged 80+ years during 2010–2019.

Discussion

RF radiation is a current controversy regarding cancer risks. The 2011 IARC evaluation on carcinogenesis [62], [, 63] has been downplayed and detracted by industry and captured agencies from the very beginning in spite of increasing evidence on harmful effects. However, IARC has decided that a new evaluation of cancer risks is top priority within a few years [100].

In this article we give some further data on the RF carcinogenesis. The attributable fraction gives the number of cases that could have been prevented if no risk exists for a specific exposure. Based on results in case-control studies from three study groups that have shown statistically significant increased risk for glioma and acoustic neuroma 211 glioma cases (all exposure) and 42 acoustic neuroma cases (ipsilateral exposure) would have been preventable in the longest cumulative exposure group. The preventable fraction was 4.88 and 4.63%, respectively. Highest preventable fraction was found for glioma with ipsilateral wireless phone use, 6.03% corresponding to 150 cases. Lower AF was calculated for meningioma, 1.75%, yielding 39 preventable cases (ipsilateral exposure). As displayed in Belpomme et al. [73] these results were based on Interphone [67], Coureau et al. [101], and Carlberg, Hardell [102], each without statistically significant increased risk. However, meta-analysis of these studies yielded, OR = 1.49, 95% CI = 1.08–2.06.

We have previously published results on increasing rates of tumors of unknown type in the brain or CNS both in the Swedish Inpatient Register and Causes of Death Register during 1998–2013 [103]. There was a clear

increasing trend in both genders during that time period, especially during more recent years with AAPC +1.78%, 95% CI + 0.76, 2.81% for both genders combined. A joinpoint was found in men in 2007; time period 2007– 2013 APC +4.95%, 95% CI +1.59, +8.42%. Similarly, in women a joinpoint was detected in 2008; time period 2008–2013 APC +4.08%, 95% CI +1.80, +6.41%.

We have now extended the time period up to 2019. Thus, we report increasing AAPC in both genders during 1998–2019 of similar magnitude as previously. In men the result was of borderline significance although the AAPC overlapped previous findings. Lower APC was found during more recent years in both men and women, see Figures 1 and 3. This may reflect a better diagnostic procedure and thus decreasing numbers of unknown brain tumor type. A delay in reporting to the register during recent years may also have an impact on the results.

It is noteworthy that we found highest AAPC in the age group 20–39 years in both men and women, Tables 2 and 3. We found in our case-control study on glioma a median latency period for use of mobile phone of 9.0 years (mean 10.1 years). The corresponding results for cordless phones (DECT) were 7.0 and 8.0 years, respectively [104]. In a population-based study during 2005–2006 on use of mobile and cordless phones among Swedish children aged 7–14 years 79.1% reported access to mobile phone and use of cordless phone was reported by 83.8% [105]. Thus, our current findings with increasing numbers of brain tumors in the age group 20–39 years may be consistent with use of wireless phones taking a reasonable latency

period. Moreover, our previous results showed highest risk for subjects that started the use of mobile or cordless phone before 20 years of age [104]. That age groups would also be more vulnerable to RF radiation [106]. In legends to Figures 1 and 3 we report the history for wireless phone use in Sweden. Figure 5 displays the number of out-going mobile phone minutes in millions during 2000–2019 in Sweden. The major increase is since early 21st century and may be associated with our findings of increasing numbers of brain tumors of unknown type considering a reasonable latency time.

As we have discussed elsewhere the Swedish Cancer Register is not reliable to study the incidence of brain tumors [103], [107]. The register is mainly based on reporting of cases with histopathological diagnosis. Now diagnosis may be based on CT and/or MRI without further investigations especially of patients with poor outcome. Biopsy or operation may be difficult to perform due to tumor location, age and co-morbidity. In the Swedish Cancer Register about 90% of the cases are diagnosed with cytology or histology, a number that has increased somewhat during recent years [107]. This fact indicates that brain tumors of unknown type are under-reported to the Cancer Register.

This review gives insight into missed opportunities for cancer prevention exemplified by asbestos, tobacco, certain pesticides and now RF radiation. No doubt economic considerations are favored instead of cancer prevention. The cancer victim is the loser in terms of suffering, life quality and shorter life expectancy. Also the life for the next-of-kin is affected. A strategy to sow doubt on cancer risks was established decades ago and is now adopted and implemented in a more sophisticated way by the telecom industry regarding RF-EMF risks to human beings and the environment. Industry has the economic power, access to politicians and media whereas concerned people are unheard.

Open access paper: <https://www.degruyter.com/document/doi/10.1515/reveh-2020-0168/html> or <https://www.degruyter.com/document/doi/10.1515/reveh-2020-0168/pdf>

REVIEW: GENETIC EFFECTS OF NON-IONIZING ELECTROMAGNETIC FIELDS

Henry Lai. Genetic effects of non-ionizing electromagnetic fields. *Electromagnetic Biology and Medicine*. Published online: 04 Feb 2021. doi: 10.1080/15368378.2021.1881866

Abstract

This is a review of the research on the genetic effects of non-ionizing electromagnetic field (EMF), mainly on radiofrequency radiation (RFR) and static and extremely low frequency EMF (ELF-EMF). The majority of the studies are on genotoxicity (e.g., DNA damage, chromatin conformation changes, etc.) and gene expression. Genetic effects of EMF depend on various factors, including field parameters and characteristics (frequency, intensity, wave-shape), cell type, and exposure duration. The types of gene expression affected (e.g., genes involved in cell cycle arrest, apoptosis and stress responses, heat-shock proteins) are consistent with the findings that EMF causes genetic damages. Many studies reported effects in cells and animals after exposure to EMF at intensities similar to those in the public and occupational environments. The mechanisms by which effects are induced by EMF are basically unknown. Involvement of free radicals is a likely possibility. EMF also interacts synergistically with different entities on genetic functions. Interactions, particularly with chemotherapeutic compounds, raise the possibility of using EMF as an adjuvant for cancer treatment to increase

the efficacy and decrease side effects of traditional chemotherapeutic drugs. Other data, such as adaptive effects and mitotic spindle aberrations after EMF exposure, further support the notion that EMF causes genetic effects in living organisms.

<https://www.tandfonline.com/doi/abs/10.1080/15368378.2021.1881866>

For excerpts from this paper see: <https://www.saferemr.com/2021/02/genetic-effects-of-non-ionizing.html>

THE ASSOCIATION BETWEEN SMARTPHONE USE AND BREAST CANCER RISK AMONG TAIWANESE WOMEN: A CASE-CONTROL STUDY

Ya-Wen Shih, Chin-Sheng Hung, Cheng-Chiao Huang, Kuei-Ru Chou, Shu-Fen Niu, Sally Chan, and Hsiu-Ting Tsai. The Association Between Smartphone Use and Breast Cancer Risk Among Taiwanese Women: A Case-Control Study. *Cancer Manag Res.* 2020; 12: 10799–10807. 2020 Oct 29. doi: 10.2147/CMAR.S267415

Abstract

Introduction Breast cancer is a common malignancy worldwide. Smartphones have gradually become indispensable to our modern lives and have already changed lifestyles of human beings. To our best knowledge, no study has investigated the relationship between smartphone use and breast cancer. This case-control study purposely investigated the relationship between smartphone use and breast cancer risk.

Materials and Methods This was a case-control study comprising 894 healthy controls and 211 patients with breast cancer. All participants were asked to respond to standard questionnaires to collect information on sleep quality, smartphone addiction, and smartphone use.

Results Participants with smartphone addiction had a significantly higher 1.43-fold risk of breast cancer. Individuals with the habitual behavior of smartphone use >4.5 minutes before bedtime had a significantly increased 5.27-fold risk of breast cancer compared to those who used a smartphone for ≤4.5 minutes before bedtime. Additionally, a closer distance between the smartphone and the breasts when using the smartphone exhibited a significantly increased 1.59-fold risk. Participants who carried their smartphone near their chest or waist-abdomen area had significantly increased 5.03-fold and 4.06-fold risks of breast cancer, respectively, compared to those who carried the smartphone below the waist. Moreover, there was a synergistic effect of smartphone addiction and smartphone use of >4.5 minutes before bedtime which increased the breast cancer risk.

Conclusion Excessive smartphone use significantly increased the risk of breast cancer, particularly for participants with smartphone addiction, a close distance between the breasts and smartphone, and the habit of smartphone use before bedtime.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7605549/>

CHEMICAL POLARIZATION EFFECTS OF ELECTROMAGNETIC FIELD RADIATION FROM THE NOVEL 5G NETWORK DEPLOYMENT AT ULTRA HIGH FREQUENCY

Ugochukwu O. Matthew, Jazuli S. Kazaure. Chemical polarization effects of electromagnetic field radiation from the novel 5G network deployment at ultra high frequency. Health and Technology. Published online Jan. 27, 2021.

Abstract

The wide-spectrum of non-ionizing, non-visible radiation emitted from the novel 5G network deployment was investigated and found liable to produce effects capable of heating up and altering human body nomenclature. The Ultra-high frequency magnetic fields, induced circulation of currents in the surrounding human body when potentially exposed. The quantum of these electromagnetic charges is influenced by the magnitude of the external magnetic field. The Magnetic fields warming is the major organic consequence of the electromagnetic fields radiofrequency radiation emitted from 5G network installation especially at a very high frequencies. From the current research, the levels of electromagnetic fields to which individuals are naturally unmasked under 4G network and 5G network technology in SCENARIO1, SCENARIO 2 and SCENARIO 3 are very negligible to alter human body dipolar chemistry. On the several findings of the research, deploying 5G network technology under the ultra-high frequency above 20 GHz will produce effect that will heat up the human body tissues due to electromagnetic field inducement since human body is dipolar in nature. The research established that while the current digital society will continue investment into 5G network technology, caution must be applied not to deploy 5G network under ultra-high frequency above 20 GHz due to its adverse health effects.

Conclusions

From the knowledge and principle of electromagnetism, human beings are constituted of substantial amount of oriented cells with diverse electromagnetic field attributes. The Biological attributes of the human tissue under diverse electromagnetic radiative emission are studied and that had provided the basis upon which the current research on the effects of electromagnetic fields on the human body. The heating consequences of the radio electromagnetic waves from 5G network technology deployment had formed the fundamental basis for current research. On the several findings of the research, deploying 5G network technology under the ultra-high baseband above 20 GHz will produce effects such as heating up of the body tissues due to electromagnetic field inducement on the account that human body is dipolar in nature. The effects will extend to produce dielectric polarization, ionic polarization, interfacial polarization and orientational polarization. This is generally on the account that variations on dielectric properties of biological tissues with the frequency of the electromagnetic field inducement are very dissimilar. While it is very imperative to determine the frequency distribution in deploying the novel 5G network to avoid adverse dielectric dispersion that may flow into the human body.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7839940/>

AGE-DEPENDENCE OF ELECTROMAGNETIC POWER AND HEAT DEPOSITION IN NEAR-SURFACE TISSUES IN EMERGING 5G BANDS

Giulia Sacco, Stefano Pisa, Maxim Zhadobov. Age-dependence of electromagnetic power and heat deposition in near-surface tissues in emerging 5G bands. *Sci Rep.* 2021 Feb 17;11(1):3983. doi: 10.1038/s41598-021-82458-z.

Abstract

With the development of 5th generation (5G) mobile networks people of different ages will be exposed in the upper part of the microwave spectrum. From the perspective of non-ionizing radiation dosimetry, an accurate analysis of age-dependent electromagnetic power deposition and resulting heating is required. In this study, we evaluate the effect of age on exposure at 26 GHz and 60 GHz. A near-surface tissue model illuminated by a plane wave is used to assess the exposure considering both frequency-independent and frequency-dependent limits.

The age-related variation of the skin thickness and tissue electromagnetic properties has been considered. Moreover, the blood flow decrease rate has been taken into account to assess the age-dependent heating. Our results demonstrate that the overall variations of the power density, specific absorption rate (SAR) and heating in the near-surface tissues are limited to about 10-15%. These variations are mainly due to the tissue permittivity and blood flow change with age. In contrast to the transmitted power density that increases with age, the peak SAR decreases at both frequencies. The peak steady-state heating increases from 5 to 70 years old by roughly 11% at 26 GHz and 13% at 60 GHz.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7889919/>

REDUCTION OF 5G CELLULAR NETWORK RADIATION IN WIRELESS MOBILE PHONE USING AN ASYMMETRIC SQUARE SHAPED PASSIVE METAMATERIAL DESIGN

Tayaallen Ramachandran, Mohammad Rashed Iqbal Faruque, Air Mohammad Siddiky, Mohammad Tariqul Islam. Reduction of 5G cellular network radiation in wireless mobile phone using an asymmetric square shaped passive metamaterial design. *Sci Rep.* 2021 Jan 29;11(1):2619. doi: 10.1038/s41598-021-82105-7.

Abstract

This study aims to demonstrate the feasibility of metamaterial application in absorption reduction of 5G electromagnetic (EM) energy in the human head tissue. In a general sense, the radio frequency (RF) energy that received by wireless mobile phone from the base station, will emit to surrounding when the devices are in active mode. Since the latest fifth generation technology standard for cellular networks is upon us, the emission of radiation from any wireless devices needs to be taken into consideration. This motivation helps to prepare this paper that focuses on construction of novel and compact square-shaped metamaterial (SM) design to reduce electromagnetic exposure to humans. The commercially available substrate material known as FR-4 with thickness of 1.6 mm was selected to place the metamaterial design on it. The electromagnetic properties and Specific Absorption Rate (SAR) analyses were carried out numerically by utilising high-performance 3D EM analysis, Computer Simulation Technology Studio (CST) software. Meanwhile, for the validation purpose, the metamaterial designs for both unit and array cells were fabricated to measure the electromagnetic

properties of the material. From the numerical simulation, the introduced SM design manifested quadruple resonance frequencies in multi bands precisely at 1.246 (at L-band), 3.052, 3.794 (at S-band), and 4.858 (C-band) GHz.

However, the comparison of numerically simulated and measured data reveals a slight difference between them where only the second resonance frequency was decreased by 0.009 GHz while other frequencies were increased by 0.002, 0.045, and 0.117 GHz in sequential order. Moreover, the SAR analysis recorded high values at 3.794 GHz with 61.16% and 70.33% for 1 g and 10 g of tissue volumes, respectively. Overall, our results demonstrate strong SAR reduction effects, and the proposed SM design may be considered a promising aspect in the telecommunication field.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7846749/>
or <https://www.nature.com/articles/s41598-021-82105-7>

RADIOFREQUENCY RADIATION: A POSSIBLE THREAT TO MALE FERTILITY

Himanshi, Umesh Rai, Rajeev Singh. Radiofrequency Radiation: A possible threat to male fertility. *Reprod Toxicol.* 2021 Jan 23;S0890-6238(21)00021-6. doi: 10.1016/j.reprotox.2021.01.007.

Abstract

Radiofrequency exposure from man-made sources has increased drastically with the era of advanced technology. People could not escape from such RF radiations as they have become the essential part of our routine life such as Wi-Fi, microwave ovens, TV, mobile phones, etc. Although non-ionizing radiations are less damaging than ionizing radiations but its long term exposure effect cannot be avoided. For fertility to be affected, either there is an alteration in germ cell, or its nourishing environment, and RF affects both the parameters subsequently, leading to infertility. This review with the help of in vitro and in vivo studies shows that RF could change the morphology and physiology of germ cells with affected spermatogenesis, motility and reduced concentration of male gametes. RF also results in genetic and hormonal changes. In addition, the contribution of oxidative stress and protein kinase complex after RFR exposure is also summarized which could also be the possible mechanism for reduction in sperm parameters. Further, some preventative measures are described which could help in reverting the radiofrequency effects on germ cells.

<https://pubmed.ncbi.nlm.nih.gov/33497741/>

MALE CELLULAR TELEPHONE EXPOSURE, FECUNDABILITY, AND SEMEN QUALITY: RESULTS FROM TWO PRECONCEPTION COHORT STUDIES

E E Hatch, S K Willis, A K Wesselink, E M Mikkelsen, M L Eisenberg, G J Sommer, H T Sorensen, K J Rothman, L A Wise. Human Reproduction, deab001, 10 February 2021. <https://doi.org/10.1093/humrep/deab001>

Abstract

STUDY QUESTION To what extent is exposure to cellular telephones associated with male fertility?

SUMMARY ANSWER Overall, we found little association between carrying a cell phone in the front pants pocket and male fertility, although among leaner men (BMI <25 kg/m²), carrying a cell phone in the front pants pocket was associated with lower fecundability.

WHAT IS KNOWN ALREADY Some studies have indicated that cell phone use is associated with poor semen quality, but the results are conflicting

STUDY DESIGN, SIZE, DURATION Two prospective preconception cohort studies were conducted with men in Denmark (n = 751) and in North America (n = 2349), enrolled and followed via the internet from 2012 to 2020.

PARTICIPANTS/MATERIALS, SETTING, METHODS On the baseline questionnaire, males reported their hours/day of carrying a cell phone in different body locations. We ascertained time to pregnancy via bi-monthly follow-up questionnaires completed by the female partner for up to 12 months or until reported conception. We used proportional probabilities regression models to estimate fecundability ratios (FRs) and 95% confidence intervals (CIs) for the association between male cell phone habits and fecundability, focusing on front pants pocket exposure, within each cohort separately and pooling across the cohorts using a fixed-effect meta-analysis. In a subset of participants, we examined selected semen parameters (semen volume, sperm concentration and sperm motility) using a home-based semen testing kit.

MAIN RESULTS AND THE ROLE OF CHANCE There was little overall association between carrying a cell phone in a front pants pocket and fecundability: the FR for any front pants pocket exposure versus none was 0.94 (95% CI: 0.0.83–1.05). We observed an inverse association between any front pants pocket exposure and fecundability among men whose BMI was <25 kg/m² (FR = 0.72, 95% CI: 0.59–0.88) but little association among men whose BMI was ≥25 kg/m² (FR = 1.05, 95% CI: 0.90–1.22). There were few consistent associations between cell phone exposure and semen volume, sperm concentration, or sperm motility.

LIMITATIONS, REASONS FOR CAUTION Exposure to radiofrequency radiation from cell phones is subject to considerable non-differential misclassification, which would tend to attenuate the estimates for dichotomous comparisons and extreme exposure categories (e.g. exposure 8 vs. 0 h/day). Residual confounding by occupation or other unknown or poorly measured factors may also have affected the results.

WIDER IMPLICATIONS OF THE FINDINGS Overall, there was little association between carrying one's phone in the front pants pocket and fecundability. There was a moderate inverse association between front pants pocket cell phone exposure and fecundability among men with BMI <25 kg/m², but not among men with BMI ≥25 kg/m². Although several previous studies have indicated associations between cell phone exposure and lower sperm motility, we found few consistent associations with any semen quality parameters.

STUDY FUNDING/COMPETING INTEREST(S) The study was funded by the National Institutes of Health, grant number R03HD090315. In the last 3 years, PRESTO has received in-kind donations from Sandstone Diagnostics (for semen kits), Swiss Precision Diagnostics (home pregnancy tests), Kindara.com (fertility app), and FertilityFriend.com (fertility app). Dr. L.A.W. is a fibroid consultant for AbbVie, Inc. Dr. H.T.S. reports that the Department of Clinical Epidemiology is involved in studies with funding from various companies as research grants to and administered by Aarhus University. None of these studies are related to the current study. Dr. M.L.E. is an advisor to Sandstone Diagnostics, Ro, Dadi, Hannah, and Underdog. Dr. G.J.S. holds ownership in Sandstone Diagnostics Inc., developers of the Trak Male Fertility Testing System. In addition, Dr. G.J.S. has a patent pending related to Trak Male Fertility Testing System issued.

<https://academic.oup.com/humrep/advance-article-abstract/doi/10.1093/humrep/deab001/6132077>

WHAT IS THE RADIATION BEFORE 5G? A CORRELATION STUDY BETWEEN MEASUREMENTS IN SITU AND IN REAL TIME AND EPIDEMIOLOGICAL INDICATORS IN VALLECAS, MADRID

Isabel López, Nazario Félix, Marco Rivera, Adrián Alonso, Ceferino Maestú. Environmental Research. Volume 194, March 2021, 110734. <https://doi.org/10.1016/j.envres.2021.110734>.

Highlights

- People who are exposed to higher radiation values present more severe headaches, dizziness and nightmares.
- The methodology for obtaining electromagnetic radiation measurements should be reviewed.
- The population continues to receive radiation peaks in distances greater than 200 m, no one is free from exposure.

Abstract

Background Exposure of the general population to electromagnetic radiation emitted by mobile phone base stations is one of the greater concerns of residents affected by the proximity of these structures due to the possible relationship between radiated levels and health indicators.

Objectives This study aimed to find a possible relationship between some health indicators and electromagnetic radiation measurements.

Methods A total of 268 surveys, own design, were completed by residents of a Madrid neighborhood surrounded by nine telephone antennas, and 105 measurements of

electromagnetic radiation were taken with a spectrum analyzer and an isotropic antenna, in situ and in real – time, both outside and inside the houses.

Results It was shown statistically significant p - values in headaches presence ($p = 0.010$), nightmares ($p = 0.001$), headache intensity ($p < 0.001$), dizziness frequency ($p = 0.011$), instability episodes frequency ($p = 0.026$), number of hours that one person sleeps per day ($p < 0.001$) and three of nine parameters studied from tiredness. Concerning cancer, there are 5.6% of cancer cases in the study population, a percentage 10 times higher than that of the total Spanish population.

Discussion: People who are exposed to higher radiation values present more severe headaches, dizziness and nightmares. Moreover, they sleep fewer hours.

<https://www.sciencedirect.com/science/article/abs/pii/S0013935121000281>

ASSOCIATION BETWEEN THE PATTERN OF MOBILE PHONE USE AND SLEEP QUALITY IN NORTHEAST CHINA COLLEGE STUDENTS

Jia Meng, Fang Wang, Rentong Chen, Hui Hua, Qian Yang, Dan Yang, Nan Wang, Xin Li, Fangfang Ma, Liting Huang, Zhenzhen Zou, Menglin Li, Tieting Wang, Yannan Luo, Yunda Li, Yang Liu. Sleep Breath. 2021 Feb 2. doi: 10.1007/s11325-021-02295-2.

Abstract

Objectives: Currently, mobile penetration is high amongst college students. The aims of this study were to investigate the characteristics of mobile phone use and to explore the influence of mobile phone use characteristics on sleep quality amongst college students.

Methods: From December 2016 to January 2017, we collected mobile phone use characteristics and sleep quality data using the Pittsburgh Sleep Quality Index (PSQI) and standardised questionnaires that were answered by 4500 medical university students in Liaoning Province (actual response rate of 94%, $n = 4234$ college students). This study used the SPSS 21.0 software to establish the database and perform the statistical analysis.

Results: One hundred percent of the college students had mobile phones and used mobile phones for entertainment (91%), work (51%), obtaining information (61%), and other purposes (23%). Additionally, there was a statistically significant difference in the PSQI score between students who held the phone at a distance of more than 10 cm from their eyes and those who held it a distance of less than 10 cm ($P = 0.002$). Multiple logistic regression analysis showed that the risk of poor sleep quality was 1.21-1.53 times higher for those who spent more than 5 h a day using their phones and 1.41-1.59 times higher for those who used their phones for more than half an hour before going to bed when the lights were off.

Conclusions: Daily cumulative mobile phone use and use with the lights off before sleep are associated with poorer sleep quality.

<https://pubmed.ncbi.nlm.nih.gov/33532989/>

THE EFFECT OF CONTINUOUS LOW-INTENSITY EXPOSURE TO ELECTROMAGNETIC FIELDS FROM RADIO BASE STATIONS TO CANCER MORTALITY IN BRAZIL

Nádia Cristina Pinheiro Rodrigues, Adilza Condessa Dode, Mônica Kramer de Noronha Andrade, Gisele O'Dwyer, Denise Leite Maia Monteiro, Inês Nascimento Carvalho Reis, Roberto Pinheiro Rodrigues, Vera Cecília Frossard, Valéria Teresa Saraiva Lino. *Int J Environ Res Public Health*. 2021 Jan 29;18(3):1229. doi: 10.3390/ijerph18031229.

Abstract

Background: this study aims to estimate the rate of death by cancer as a result of Radio Base Station (RBS) radiofrequency exposure, especially for breast, cervix, lung, and esophagus cancers.

Methods: we collected information on the number of deaths by cancer, gender, age group, gross domestic product per capita, death year, and the amount of exposure over a lifetime. We investigated all cancer types and some specific types (breast, cervix, lung, and esophagus cancers).

Results: in capitals where RBS radiofrequency exposure was higher than 2000/antennas-year, the average mortality rate was 112/100,000 for all cancers. The adjusted analysis showed that, the higher the exposure to RBS radiofrequency, the higher cancer mortality was. The highest adjusted risk was observed for cervix cancer (rate ratio = 2.18). The spatial analysis showed that the highest RBS radiofrequency exposure was observed in a city in southern Brazil that also showed the highest mortality rate for all types of cancer and specifically for lung and breast cancer.

Conclusion: the balance of our results indicates that exposure to radiofrequency electromagnetic fields from RBS increases the rate of death for all types of cancer.

Open access paper: <https://www.mdpi.com/1660-4601/18/3/1229>

SONIC HEALTH ATTACKS BY PULSED MICROWAVES IN HAVANA REVISITED [HEALTH MATTERS]

James C. Lin. *Sonic Health Attacks by Pulsed Microwaves in Havana Revisited [Health Matters]*. IEEE Microwave Magazine. 22(3):71-73, March 2021. doi: 10.1109/MMM.2020.3044125.

Abstract

Presents the results of the U.S. National Academies of Sciences, Engineering, and Medicine (NASEM) report, "An Assessment of Illness in U.S. Government Employees and Their Families at Overseas Embassies." it is almost exactly three years since the publication of my article "Strange Reports of Weaponized Sound in Cuba" [2].

There, it was first hypothesized that "[a]ssuming that the reported events are reliable, there is actually a scientific explanation for the source of sonic energy. It could well be from a targeted beam of high-power microwave pulse radiation" [2]. In examining plausible causes of the described illnesses, the NASEM report makes that point that, among the mechanisms the study committee considered, the most plausible mechanism to explain these cases,

especially in individuals with distinct early symptoms, appears to be directed, pulsed RF (microwave) energy.

<https://ieeexplore.ieee.org/document/9343783>

NOTICE OF RETRACTION AND REPLACEMENT. LI ET AL. ASSOCIATION BETWEEN MATERNAL EXPOSURE TO MAGNETIC FIELD NONIONIZING RADIATION DURING PREGNANCY AND RISK OF ATTENTION-DEFICIT/HYPERACTIVITY DISORDER IN OFFSPRING IN A LONGITUDINAL BIRTH COHORT

De-Kun Li. Notice of Retraction and Replacement. Li et al. Association Between Maternal Exposure to Magnetic Field Nonionizing Radiation During Pregnancy and Risk of Attention-Deficit/Hyperactivity Disorder in Offspring in a Longitudinal Birth Cohort. Letter: February 18, 2021. JAMA Netw Open. 2020;3(3):e201417

To the Editor On behalf of my coauthors, I write to report a concern raised about the methods and analyses for the Original Investigation, “Association Between Maternal Exposure to Magnetic Field Nonionizing Radiation During Pregnancy and Risk of Attention-Deficit/Hyperactivity Disorder in Offspring in a Longitudinal Birth Cohort,”¹ that was published in JAMA Network Open on March 24, 2020. The concern was brought to the attention of the journal editors by a reader and the editors forwarded this concern to us. After discussion and assessment, we have determined that there were errors in the statistical analyses, which required use of different analyses and which result in changes to some of the original findings and limitations to the interpretations of this study. As a result, the editors have asked us to retract and replace this article.

This study was conducted to determine whether maternal exposure to high levels of magnetic field (MF) nonionizing radiation was associated with an increased risk of attention-deficit/hyperactivity disorder (ADHD) in offspring by using measurements of MF nonionizing radiation levels and physician-diagnosed ADHD. We included a longitudinal birth cohort of 1482 mother-child pairs whose mothers were participants of an existing birth cohort and whose level of exposure to MF nonionizing radiation was captured by wearing a monitor for 24 hours during pregnancy in 2 previous studies conducted from October 1, 1996, to October 31, 1998, and from May 1, 2006, to February 29, 2012.

To examine the association of high levels of MF nonionizing radiation with risk of ADHD, we used the 90th percentile of the 24-hour measurements as the MF index, which reflects the MF nonionizing radiation level at or above which a participant was exposed for 10% of the time during the day. We had originally classified participating mother-child dyads into low or high MF nonionizing radiation exposure groups based on the experience of previous studies.²⁻⁴ However, we erroneously reported using a 1.3-mG cut point based on the 25th percentile of the MF index’s distribution. And we then conducted analyses with participants dichotomized to a low exposure level (those whose MF nonionizing radiation level was <1.3 mG) or a high exposure level (those whose MF nonionizing radiation level was ≥1.3 mG). These analyses led to the conclusions that “in utero exposure to high levels of MF nonionizing radiation was associated with an increased risk of ADHD, especially ADHD with immune-related comorbidity.”¹

At the request of the editors, we have conducted new analyses based on continuous MF exposure levels as well as analyses based on categorical MF exposure levels, without using a cut point level. We have also conducted a dose-response analysis based on the categorical exposure levels with increased MF exposure levels being compared with the same reference group because a dose-response association is commonly examined in epidemiological studies.

With these new analyses, we continue to find in utero exposures to some, but not all, high levels of MF nonionizing radiation were associated with a higher risk of ADHD. However, the associations observed were inconsistent and nonlinear, limiting interpretations. Thus, changes are needed to the original article, including the following:

We removed the cut point–based results in the original Table 1 and replaced the data using mean (SD) based on continuous MF exposure levels. Tables 2 through 4, which previously reported findings based on dichotomous low and high MF exposure levels, have been replaced with findings from the analysis using continuous measure and categorical levels of MF exposure. We also replaced the eTable in Supplement 1 with similar analyses.

We have added an eFigure in Supplement 2 showing the findings of the dose-response analysis between MF exposure in quintiles of mother-child dyads and the risk of ADHD. In addition, we have replaced the original dichotomized Kaplan-Meier survival curve with multicategorical Kaplan-Meier survival curves for multiple maternal exposure levels.

Finally, we have added to the Discussion section the following limitations: “no corrections were made for multiple comparisons and for many of the significant associations observed among children with ADHD and concurrent immune-related comorbidities, the CIs were wide, indicating that these findings should be interpreted with caution.”

On behalf of my coauthors, I apologize for any confusion this has caused readers and have requested that the original article be retracted and replaced with a corrected version. The corrections affect the Abstract, text, Tables, Figure, and Supplement 1. The replacement article includes new supplements with a copy of the original article with the errors highlighted and another copy with the corrections highlighted.

Open access paper:

<https://jamanetwork.com/journals/jamanetworkopen/fullarticle/2776553>

ASSOCIATIONS BETWEEN ARTIFICIAL LIGHT AT NIGHT AND RISK FOR THYROID CANCER: A LARGE US COHORT STUDY

Dong Zhang, Rena R Jones, Peter James, Cari M Kitahara, Qian Xiao. *Cancer*. 2021 Feb 8. doi: 10.1002/cncr.33392.

Abstract

Background: Light at night (LAN) inhibits nighttime secretion of melatonin and may cause circadian disruption, which may be a risk factor for cancer. Recent studies have linked high LAN exposure with elevated breast cancer risk. Given that breast cancer may share a common hormone-dependent etiology with thyroid cancer and that circadian rhythms play a role in

regulating thyroid function, the authors hypothesized that exposure to LAN is positively associated with thyroid cancer incidence.

Methods: This study examined the association between LAN and thyroid cancer incidence in the National Institutes of Health-American Association of Retired Persons Diet and Health Study. LAN exposure was estimated from satellite data and was linked to residential addresses at the baseline. Incident thyroid cancer cases were ascertained via linkage to state cancer registries. Cox regression was used to determine the relationship between LAN and thyroid cancer risk, with adjustments made for sociodemographic, lifestyle, and other environmental factors.

Results: Among 464,371 participants, a positive association was found between LAN and thyroid cancer risk. Specifically, in comparison with the lowest quintile of LAN, the highest quintile was associated with a 55% increase in risk (hazard ratio [HR], 1.55; 95% confidence interval [CI], 1.18-2.02). The association was primarily driven by papillary thyroid cancer and was stronger in women (HR, 1.81; 95% CI, 1.26-2.60) than men (HR, 1.29; 95% CI, 0.86-1.94). In women, the association was stronger for localized cancer, whereas in men, the association was stronger for a more advanced stage. Results were consistent across different tumor sizes.

Conclusions: LAN was positively associated with thyroid cancer risk. Future studies are needed to confirm this association and identify underlying biological mechanisms.

<https://pubmed.ncbi.nlm.nih.gov/33554351/>

EFFECTS OF SELECTIVE OUTCOME REPORTING ON RISK PERCEPTION

F Freudenstein, R J Croft, S P Loughran, B M Zeleke, P M Wiedemann. Effects of selective outcome reporting on risk perception. *Environ Res.* 2021 Feb 3;110821. doi: 10.1016/j.envres.2021.110821.

Highlights

- An experimental study investigated how selective reporting of study results - indicating increased health effects - influenced its receiver's risk perception.
- Selective reporting of risk information increased risk perception when compared to receiving the full information.
- Selectively informed subjects revealed a stronger tendency towards overgeneralization of health risks.
- The results indicate that reporting of full study results is an important factor in effective risk communication.

Abstract

The current study aimed to investigate how selective reporting of study results indicating increased health effects will influence its receiver's risk perception. Using the example of the Interphone Study from 2010 on mobile phone usage and cancer, an online experiment was conducted separating respondents into two groups. One group of subjects was informed selectively about a relationship between heavy mobile phone use and an elevated risk of glioma (brain cancer) only. The other group of subjects was informed about the full results of the analyses of glioma risk by cumulative call time, which suggests that other than for the heavy users, there were no statistically significant elevated risks related to mobile phone use.

The results showed that selective reporting of risk information increased risk perception when compared to receiving the full information. Additionally, the selectively informed subjects revealed a stronger tendency towards overgeneralization of the 'elevated brain cancer risk' to all mobile phone users, although this did not extend to an overgeneralization to other electromagnetic field sources or differences in the perception of a usage time dependency for possible health risks. These results indicate that reporting of full results is an important factor in effective risk communication.

<https://pubmed.ncbi.nlm.nih.gov/33548295/>
<https://pubmed.ncbi.nlm.nih.gov/33548295/>

ELECTROMAGNETIC RADIATION AS AN EMERGING DRIVER FACTOR FOR THE DECLINE OF INSECTS

Alfonso Balmori. Electromagnetic radiation as an emerging driver factor for the decline of insects. *Sci Total Environ.* Available online 28 January 2021, 144913. <https://doi.org/10.1016/j.scitotenv.2020>.

Highlights

- Biodiversity of insects is threatened worldwide
- This reductions is mainly attributed to agricultural practice and pesticide use
- There is sufficient evidence on the damage caused by electromagnetic radiation
- Electromagnetic radiation may be a complementary driver in this decline
- The precautionary principle should be applied before any new deployment (e.g. 5G)

Abstract

The biodiversity of insects is threatened worldwide. Numerous studies have reported the serious decline in insects that has occurred in recent decades. The same is happening with the important group of pollinators, with an essential utility for pollination of crops. Loss of insect diversity and abundance is expected to provoke cascading effects on food webs and ecosystem services. Many authors point out that reductions in insect abundance must be attributed mainly to agricultural practices and pesticide use. On the other hand, evidence for the effects of non-thermal microwave radiation on insects has been known for at least 50 years. The review carried out in this study shows that electromagnetic radiation should be considered seriously as a complementary driver for the dramatic decline in insects, acting in synergy with agricultural intensification, pesticides, invasive species and climate change. The extent that anthropogenic electromagnetic radiation represents a significant threat to insect pollinators is unresolved and plausible. For these reasons, and taking into account the benefits they provide to nature and humankind, the precautionary principle should be applied before any new deployment (such 5G) is considered.

Excerpt

The precautionary principle and the importance of seriously considering EMR as a factor of insect decline.

Despite the strong scientific evidence of the negative impacts of electromagnetic radiation on insects, a recent study funded by the European Union's Horizon 2020 Research and Innovation Programme (EKLIPSE) stated that our current knowledge concerning the impact of anthropogenic RF-EMR on pollinators (and other invertebrates) is inconclusive (Vanbergen et al., 2019). Thus, the extent to which anthropogenic EMR represents a significant threat to

insect pollinators is unresolved. For these reasons, and taking into account the benefits they provide to nature and humankind, the precautionary principle of the European Union (Communication from the Commission on the Precautionary Principle, 2000) should be applied.

The potential effects of RF-EMFs on most taxonomic groups, including migratory birds, bats and insects, are largely unknown, and the potential effects on wildlife could become more relevant with the expected adoption of new mobile network technology (5G), raising the possibility of unintended biological consequences (Sutherland et al., 2018). Thus, before any new deployment (such 5G) is considered, its effects should be clearly assessed, at least while conclusions are drawn and these existing uncertainties are overcome, according to the official document 'Late Lessons of Early Warnings' (European Environment Agency, 2013).

A letter by the United States Department of the Interior sent to the National Telecommunications and Information Administration in the Department of Commerce warns about the scarcity of studies carried out on the impacts from non-ionising EMR emitted by communication towers (United States Department of the Interior, 2014). The precise potential effects of increases in EMR on wildlife, which are not yet well recognised by the global conservation community, have been identified as an important emerging issue for global conservation and biological diversity (Sutherland et al., 2018). Thus, as we have explained in this review, EMR should be seriously considered as a complementary driver for the dramatic decline in insects in recent studies, acting in synergy with agricultural intensification, pesticides, invasive species and climate change.

<https://www.sciencedirect.com/science/article/abs/pii/S0048969720384461>

BIOLOGICAL EFFECTS OF EXPOSURE TO A RADIOFREQUENCY ELECTROMAGNETIC FIELD ON THE PLACENTAL BARRIER IN PREGNANT RATS

Hye Sun Kim, Hyung-Do Choi, Jeong-Ki Pack, Nam Kim, Young Hwan Ahn. Biological Effects of Exposure to a Radiofrequency Electromagnetic Field on the Placental Barrier in Pregnant Rats. Bioelectromagnetics. 2021 Feb 1. doi: 10.1002/bem.22322.

Abstract

The placenta protects the fetus against excessive stress-associated maternal cortisol during pregnancy. We studied whether exposure to radiofrequency electromagnetic field (RF-EMF) radiation during pregnancy can cause changes in dams and their placentas. Pregnant Sprague-Dawley rats were divided into cage-control, sham-exposed, and RF-exposed groups. They were exposed to RF-EMF signals at a whole-body specific absorption rate of 4 W/kg for 8 h/day from gestational Day 1 to 19. Levels of cortisol in the blood, adrenal gland, and placenta were measured by enzyme-linked immunosorbent assay. Levels of adrenocorticotrophic hormone and corticotropin-releasing hormone were monitored in maternal blood. Expression levels of placental 11 β -hydroxysteroid dehydrogenase type 2 (11 β -HSD2) messenger RNA (mRNA) were measured by reverse transcription polymerase chain reaction. Morphological changes in the placenta were analyzed using hematoxylin and eosin staining. Fetal parts of the placenta were measured using Zen 2.3 blue edition software. Maternal cortisol in circulating blood (RF: 230 $\pm$ 24.6 ng/ml and Sham: 156 $\pm$ 8.3 ng/ml) and the adrenal gland (RF:

58.3 ± 4.5 ng/ml and Sham: 30 ± 3.8 ng/ml) was significantly increased in the RF-exposed group ($P < 0.05$).

Placental cortisol was stably maintained, and the level of placental 11 β -HSD2 mRNA expression was not changed in the RF-exposed group. RF-EMF exposure during pregnancy caused a significant elevation of cortisol levels in circulating blood; however, no changes in the placental barrier were observed in pregnant rats.

<https://pubmed.ncbi.nlm.nih.gov/33527465/>

ANALYZING THE IMPACT OF 900 MHZ EMF SHORT-TERM EXPOSURE TO THE EXPRESSION OF 667 MIRNAS IN HUMAN PERIPHERAL BLOOD CELLS

Andreas Lamkowski, Matthias Kreitlow, Jörg Radunz, Martin Willenbockel, Marcus Stiemer, Lars Ole Fichte, Carl Friedrich Rädcl, Matthäus Majewski, Patrick Ostheim, Matthias Port, Michael Abend. Analyzing the impact of 900 MHz EMF short-term exposure to the expression of 667 miRNAs in human peripheral blood cells. *Sci Rep.* 2021 Feb 24;11(1):4444. doi: 10.1038/s41598-021-82278-1.

Abstract

More than ever before, people around the world are frequently exposed to different sections of the electromagnetic spectrum, mainly emitted from wireless modern communication technologies. Especially, the level of knowledge on non-thermal biological EMF effects remains controversial. New technologies allow for a more detailed detection of non-coding RNAs which affect the post-transcriptional control. Such method shall be applied in this work to investigate the response of human blood cells to electromagnetic irradiation. In this ex vivo in vitro study, we exposed peripheral blood cells from 5 male donors to a continuous wave of 900 MHz EMF for 0, 30, 60 and 90 min. Significant micro RNA (miRNA) expression changes ($p \leq 0.05$) above or below the SHAM exposed samples were evaluated using a quantitative real time PCR platform for simultaneous detection of 667 miRNAs called low density array. Only significant miRNA expression changes which were detectable in at least 60% of the samples per exposure group were analyzed. The results were compared with data from room temperature + 2 °C (RT + 2 °C) samples (here referred to as hyperthermia) to exclude miRNA expression altered by hyperthermia. The validation study by using the same donors and study design was performed after an interval of 2 years. When analyzing a total of 667 miRNAs during the screening study, 2 promising candidate miRNAs were identified, which were down regulated almost twice and showed a complete separation from the unexposed control group (miR-194 at 30 min and miR-939 at 60 min). The p-values even survived the Bonferroni correction for multiple comparisons ($p = 0.0007$ and $p = 0.004$, respectively). None of these miRNAs were expressed at a second time point after EMF exposure. Following an alternative analysis approach, we examined for miRNAs revealing an expected significant association of differential miRNA expression with the dose-time EMF exposure product, separately for each donor. Donors 2 and 3 revealed 11 and 10 miRNA species being significantly associated with EMF exposure which differed significantly from the other donors showing a minor number of differentially expressed miRNAs and could identify donors 2 and 3 as particularly EMF-responsive.

The measurements were repeated after 2 years. The number of expressed/non-expressed miRNAs was almost similar (97.4%), but neither the number nor the previously differentially expressed miRNAs could be reproduced. Our data neither support evidence of early changes at miRNA expression level in human whole blood cells after 900 MHz EMF exposure nor the identification of EMF-responsive individuals.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7904780/>

CHRONIC EXPOSURE TO 2.45 GHZ MICROWAVE RADIATION IMPROVES COGNITION AND SYNAPTIC PLASTICITY IMPAIRMENT IN VASCULAR DEMENTIA MODEL

Mahnaz Bayat, Narges Karimi, Mohammad Karami, Afshin Borhani Haghighi, Kamjoo Bayat, Somayeh Akbari, Masoud Haghani. Chronic exposure to 2.45 GHz microwave radiation improves cognition and synaptic plasticity impairment in vascular dementia model. *Int J Neurosci.* 2021 Feb 26;1-13. doi: 10.1080/00207454.2021.1896502.

Abstract

In this study, we evaluated the effects of 2.45 GHz microwave radiation on cognitive dysfunction induced by vascular dementia (VaD). The VaD was induced by bilateral-common carotid occlusion (2-VO). The rats were divided into 4 groups including: control (n = 6), sham (n = 6), 2-VO (n = 8), and 2-VO + Wi-Fi (n = 10) groups. Wi-Fi modem centrally located at the distance of 25 cm from the animal's cages and the animals were continuously exposed to Wi-Fi signal while they freely moved in the cage (2 h/day for forty-five days). Therefore, the power density (PD) and specific absorption rate value (SAR) decreased at a distance of 25 to 60 cm (PD = 0.018 to 0.0032 mW/cm², SAR = 0.0346 to 0.0060 W/Kg). The learning, memory, and hippocampal synaptic-plasticity were evaluated by radial arm maze (RAM), passive avoidance (PA), and field-potential recording respectively.

The number of hippocampal CA1 cells was also assessed by giemsa-staining. Our results showed that VaD model led to impairment in the spatial learning and memory performance in RAM and PA that were associated with long-term potentiation (LTP) impairment, decrease of basal-synaptic transmission (BST), increase of GABA transmission, and decline of neurotransmitter release-probability as well as hippocampal cell loss. Notably, chronic Wi-Fi exposure significantly recovered the learning-memory performance, LTP induction, and cell loss without any effect on BST. The LTP recovery by Wi-Fi in the 2-VO rats was probably related to significant increases in the hippocampal CA1 neuronal density, partial recovery of neurotransmitter release probability, and reduction of GABA transmission as evident by rescue of paired-pulse ratio 10ms.

<https://pubmed.ncbi.nlm.nih.gov/33635159/>

HISTOPATHOLOGICAL AND EPIGENETIC ALTERATIONS IN THE SPINAL CORD DUE TO PRENATAL ELECTROMAGNETIC FIELD EXPOSURE: AN H3K27ME3-RELATED MECHANISM

Ayşe İkinci Keleş, Burcu Biterge Süt. Histopathological and epigenetic alterations in the spinal cord due to prenatal electromagnetic field exposure: An H3K27me3-related mechanism. *Toxicol Ind Health*. 2021 Feb 23. doi: 10.1177/0748233721996947.

Abstract

Neural system development is one of the most important stages of embryogenesis. Perturbations in this crucial process due to genetic and environmental risk factors cause neural tube defects and other central nervous system diseases. We investigated the effects of prenatal exposure to 900-MHz electromagnetic field (EMF) on the spinal cord. Pregnant rats were exposed to 900-MHz EMF for 1 h/day from E13.5 until birth. Six pups from the control and EMF groups were sacrificed at postnatal day 32, and the upper thoracic region of the spine was removed and processed for histological procedures. For histopathological analyses, hematoxylin&eosin staining and, for stereological analyses and the quantitation of motor neurons, cresyl violet staining was performed.

H3K27me3 levels were determined via immunofluorescence staining. Histopathological analysis identified structural alterations of ependymal cells, enlarged central canals, as well as degenerated and shrunken motor neurons in the EMF group, while the control group tissues had normal appearances. We also observed enrichment of H3K27me3 in the ependymal cells and the motor neurons in the spinal cord of the control group rats, while the EMF group had low levels of H3K27me3 staining. Our results suggest that the loss of H3K27me3 signals might correlate with reduced neuronal stem cell potential in the EMF group and result in anatomical and structural differences in the spinal cord. This study provided a comprehensive histopathological analysis of the spinal cord after prenatal EMF exposure and offered an H3K27me3-dependent molecular explanation for the detrimental effects of EMF exposure on the spine.

<https://pubmed.ncbi.nlm.nih.gov/33620299/>

AN EVALUATION OF THE GENOTOXIC EFFECTS OF ELECTROMAGNETIC RADIATION AT 900 MHZ, 1800 MHZ, AND 2100 MHZ FREQUENCIES WITH A SMART ASSAY IN DROSOPHILA MELANOGASTER

Merve Gunes, Kayhan Ates, Burcin Yalcin, Sibel Akkurt, Sukru Ozen, Bulent Kaya. An Evaluation of the Genotoxic Effects of Electromagnetic Radiation at 900 MHz, 1800 MHz, and 2100 MHz Frequencies with a SMART Assay in *Drosophila melanogaster*. *Electromagn Biol Med*. 2021 Feb 24;1-10. doi: 10.1080/15368378.2021.1878210.

Abstract

With the development of today's technology, the electromagnetic radiation spread by mobile phones and base stations is also rapidly increasing, and this causes serious concerns about the environment and human health. The *Drosophila* model organism is widely used in genetic toxicology studies because its genome is highly similar to the genes identified in human diseases. In this study, the genotoxic effects of radiofrequency electromagnetic radiation were evaluated by the wing Somatic Mutation and Recombination Test (SMART) in *Drosophila melanogaster* at 900 MHz, 1800 MHz, and 2100 MHz. The SMART method is based on the

observation of genetic changes occurring in the trichomes of the *Drosophila* wings appearing as mutant clones under the microscope. Throughout the study, total clone parameters were evaluated by exposing the *Drosophila* larvae to electromagnetic fields for two, four, and six hours per day for two days. As a result of the study, it was observed that the number of mutant clones was statistically increased according to the negative control group in all applications except for the six-hour application at 1800 MHz. <https://pubmed.ncbi.nlm.nih.gov/33622140/>

Conclusion

... To summarise, this article includes data on *Drosophila* larvae exposure to RF-EMR at 900 MHz, 1800 MHz, and 2100 MHz for evaluating genotoxicity by virtue of the SMART test. Even if the six-hour experiment's results at 1800 MHz showed no statistical significance, the rest of the obtained results showed that the number of mutant clones was increased as compared with the negative control group. Also, experimental results showed that the effects of RF-EMR depend on the exposed electrical field, the frequency, and the span between *D. melanogaster* and the antenna.

Studies have shown that EMR can induce various types of cancer and can cause DNA damage (Brainard et al. 1999; Hardell and Sage 2008; Phillips et al. 2009). In addition, there are experimental studies in the production of ROS which have shown no significant results in apoptosis (Hou et al. 2015). Although studies have shown that EMR has a genotoxic effect, the opposite results have also been observed (Liu et al. There are contradictions in the studies on the biological effects of EMR (Lonn et al. 2005; Cao et al. 2009). This suggests that various cell lines or different systems may have different results. To our knowledge, this is the first study in which the genotoxic effects of the 900 MHz, 1800 MHz, and 2100 MHz radiofrequencies have been studied in *D. melanogaster*. Today, extensive studies are needed to demonstrate the toxicity of RF-EMF (Vila et al. 2018).

GENOTOXICITY OF RADIOFREQUENCY ELECTROMAGNETIC FIELDS: PROTOCOL FOR A SYSTEMATIC REVIEW OF IN VITRO STUDIES

Stefania Romeo, Olga Zeni, Anna Sannino, Susanna Lagorio, Mauro Biffoni, Maria Rosaria Scarfi. Genotoxicity of radiofrequency electromagnetic fields: Protocol for a systematic review of in vitro studies. *Environ Int.* 2021 Jan 21;148:106386. doi: 10.1016/j.envint.2021.106386.

Abstract

Background: Exposure to radiofrequency electromagnetic fields (RF-EMF, 100 kHz - 300 GHz) emitted by wireless communication technologies is pervasive and ubiquitous. Concern has been raised about possible adverse effects to human health. In 2011 the International Agency for Research on Cancer has classified RF-EMF as possibly carcinogenic to humans, highlighting that the evidence is weak and far from conclusive. Updated systematic reviews of the scientific literature on this topic are lacking, especially for mechanistic studies.

Objectives: To develop a protocol for a systematic review of experimental studies investigating genotoxic effects induced by RF-EMF in in vitro cellular models. Genotoxicity is one of the key-biological indicators of carcinogenicity, and the most common characteristics

of established carcinogens. The predefined procedures for conducting the systematic review are outlined below.

Methods: We will follow the guidelines developed by the National Toxicology Program-Office of Health Assessment and Translation (NTP-OHAT), adapted to the evaluation of in vitro studies.

Eligibility criteria: We will include experimental in vitro studies addressing the relationship between controlled exposures to RF-EMF and genotoxicity in mammalian cells only. Eligibility for inclusion will be further restricted to peer reviewed articles reporting findings from primary studies.

Information sources: We will search the scientific literature databases NCBI PubMed, Web of Science, and EMF- Portal. No filter on publication date will be applied. Only studies published in English will be considered. The reference lists of the included papers and available reviews will be screened for unidentified relevant papers. References will be managed through Endnote X9 software.

Data extraction and synthesis of results: Data from included papers will be extracted according to predefined forms. Heterogeneity within the available evidence will determine the type of evidence synthesis that is appropriate. Findings will be summarized in tables, graphical displays and in a narrative synthesis of the available evidences. A meta-analysis will be carried out if subgroups of studies homogeneous in terms of exposure characteristics, endpoint, and cell types will be identified.

Risk of bias: The internal validity of included studies will be assessed using the NTP-OHAT Risk of Bias Rating Tool for animal studies, adapted to in vitro studies. This stage of the process will be managed through the Health Assessment Workspace Collaborative (HAWC).

Evidence appraisal: To rate confidence in the body of evidence, we will use the OHAT GRADE-based approach for animal studies.

Framework and funding: This protocol concerns one of the evidence streams considered in a larger systematic review of the scientific literature on the potential carcinogenicity of RF-EMF, performed by scientists from several Italian public research agencies. The project is supported by the Italian Workers' Compensation Authority (INAIL) in the framework of the CRA with the Istituto Superiore di Sanità "BRiC 2018/06 - Scientific evidence on the carcinogenicity of radiofrequency electromagnetic fields".

<https://pubmed.ncbi.nlm.nih.gov/33486297/>

EFFECTS OF NON-IONIZING ELECTROMAGNETIC POLLUTION ON INVERTEBRATES, INCLUDING POLLINATORS SUCH AS HONEY BEES: WHAT WE KNOW, WHAT WE DON'T KNOW, AND WHAT WE NEED TO KNOW

Friesen M, Havas M. 2020. Pages 127-138 In Working Landscapes. Proceedings of the 12th Prairie Conservation and Endangered Species Conference, February 2019, Winnipeg, Manitoba. Edited by D. Danyluk. Critical Wildlife Habitat Program, Winnipeg, Manitoba. <http://pcesc.ca/media/45404/final-2019-pcesc-proceedings.pdf>.

Abstract

Invertebrates, including pollinators such as honey bees, can be adversely affected by non-ionizing electromagnetic radiation (EMR). Sources contributing to common environmental EMR exposures include antennae (cell phone, broadcast, and radar), communications satellites, and power lines. Adverse biochemical changes and disorientation have been reported for honey bees and other invertebrates. Field studies have reported changes in abundance and composition of “key pollinator groups” (wild bees, hoverflies, bee flies, beetles, and wasps) that have been attributed to emissions from telecommunications towers. We take a close look at the biological effects on invertebrates of EMR reported in the scientific literature and a general look at evidence from studies on plants, birds, humans, and other animals (domestic, laboratory, wild). We discuss possible implications of excessive electromagnetic pollution on ecosystems and identify knowledge gaps and what we need to know before more electromagnetic pollution is added to the environment, especially in the form of 5G.

Introduction

Invertebrates (animals without backbones) are major components of most ecosystems. Insects are key to the integrity of many ecosystems in many roles including as pollinators. Honey bees play a role in pollination of domestic as well as wild plants and are often used as bio-indicator species and as a “model” to examine environmental problems. The global decline of pollinators is of grave concern and efforts are being made to identify the reasons (Potts et al. 2010; Sánchez-Bayo and Wyckhuys 2019). One factor not widely considered is the possible role of anthropogenic electromagnetic radiation (EMR).

Electromagnetic fields (EMFs) are invisible electric and magnetic fields of force. All living organisms have evolved in Earth’s natural EMFs and depend on them to live. Natural sources include Earth’s static magnetic field, and static electricity, including differences in charges among clouds and the earth that can lead to lightning. Electromagnetic radiation (EMR) originates when fields change.

Anthropogenic (human-made, artificial) EMR sources are sometimes referred to as electromagnetic pollution or electrosmog. The main frequency ranges of interest in this article are: 1) extremely low frequencies (ELF) of 50/60 to 90 Hz that emanate from sources such as power lines and building wiring; and 2) radiofrequency radiation (RFR) of 700 MHz to 6 GHz, commonly used for devices such as cell phones, radio and television, and their supporting infrastructure, e.g., cell towers, antennae on buildings, and orbiting communications satellites. Also discussed are frequencies currently being developed and deployed above 6 GHz for 5G (5th Generation) for faster and more pervasive connectivity, including the “Internet of Things”. <http://pcesc.ca/media/45404/final-2019-pcesc-proceedings.pdf>

EFFECTS OF ELECTROMAGNETIC FIELD (EMF) RADIATION ON ANDROGEN SYNTHESIS AND RELEASE FROM THE PIG ENDOMETRIUM DURING THE FETAL PERI-IMPLANTATION PERIOD

Wiktoria Kozłowska, Ewa Monika Drzewiecka, Agata Zmijewska, Anna Koziorowska, Anita Franczak. Effects of electromagnetic field (EMF) radiation on androgen synthesis and release from the pig endometrium during the fetal peri-implantation period. *Anim Reprod Sci.* 2021 Jan 21;226:106694. doi: 10.1016/j.anireprosci.2021.106694.

Abstract

An electromagnetic field (EMF) may have effects on female reproduction. This study was conducted to determine whether EMF [50 and 120 Hz, 2 and 4 h of incubation in the presence or absence of progesterone (P4, 10⁻⁵ M)] affects androgen synthesis and release from the pig endometrium. Endometrial slices were collected from pigs (n = 5) during the fetal peri-implantation period (i.e., days 15-16 of gestation) and treated in vitro with EMF. The selected endometrial slices were treated with P4 to determine whether this hormone has effects on protection of the tissue from EMF radiation. The CYP17A1 and HSD3B1 mRNA transcript abundance, steroid 17 α hydroxylase/17, 20-lyase (cytochrome P450c17) and hydroxy Δ 5steroid dehydrogenase/3 β and steroid Δ isomerase (3 β HSD) protein abundance were examined using Real-Time PCR and Western Blot procedures, respectively. In media collected after incubation, the concentrations of androstenedione (A4) and testosterone (T) were quantified using a RIA. When P4 was added to the culture medium, EMF radiation had suppressive effects on endometrial T release after 2 and 4 h of incubation when the EMF treatment was occurring and increased A4 release after 4 h of incubation with EMF at 120 Hz. When there was no inclusion of P4, release of A4 was decreased after 2 h of EMF treatment at 120 Hz and after 4 h of EMF treatment at 50 and 120 Hz. Progesterone did not have functions that protected the pig endometrium against EMF radiation during the fetal peri-implantation period.

<https://pubmed.ncbi.nlm.nih.gov/33486154/>

CHANGES IN HONEYBEE BEHAVIOR PARAMETERS UNDER THE INFLUENCE OF THE E-FIELD AT 50 HZ AND VARIABLE INTENSITY

Paweł Migdał, Agnieszka Murawska, Paweł Bieńkowski, Ewelina Berbeć, Adam Roman. *Animals (Basel).* 2021 Jan 20;11(2):E247. doi: 10.3390/ani11020247.

Abstract

EM-fields come from both natural and anthropogenic sources. This study aimed to investigate changes in honeybee behavior parameters under the influence of an electric field at 50 Hz and variable intensity. Bees were exposed for 1 h, 3 h, or 6 h to the following artificial E-field intensities: 5.0 kV/m, 11.5 kV/m, 23.0 kV/m, or 34.5 kV/m. Bees in the control group were under the influence of an E-field <2.0 kV/m. Six basic behaviors were selected for bee observation (walking, grooming, flight, stillness, contact between individuals, and wing movement). Our research shows the impact of bee exposure time on behavioral change within groups. Exposure for 3 h caused a decrease in the time that bees spent on behaviors and in the number of occurrences. After 6 h, the parameters increased within the groups, as was the case with 1 h exposure. This may indicate that there is a behavioral barrier that allows the pattern to normalize for some time. Open access paper: <https://www.mdpi.com/2076-2615/11/2/247>

THE EFFECT OF AN ANTHROPOGENIC MAGNETIC FIELD ON THE EARLY DEVELOPMENTAL STAGES OF FISHES-A REVIEW

Krzysztof Formicki, Agata Korzelecka-Orkisz, Adam Tański. The Effect of an Anthropogenic Magnetic Field on the Early Developmental Stages of Fishes-A Review. *Int J Mol Sci.* 2021 Jan 26;22(3):1210. doi: 10.3390/ijms22031210.

Abstract

The number of sources of anthropogenic magnetic and electromagnetic fields generated by various underwater facilities, industrial equipment, and transferring devices in aquatic environment is increasing. These have an effect on an array of fish life processes, but especially the early developmental stages. The magnitude of these effects depends on field strength and time of exposure and is species-specific. We review studies on the effect of magnetic fields on the course of embryogenesis, with special reference to survival, the size of the embryos, embryonic motor function, changes in pigment cells, respiration hatching, and directional reactions. We also describe the effect of magnetic fields on sperm motility and egg activation. Magnetic fields can exert positive effects, as in the case of the considerable extension of sperm capability of activation, or have a negative influence in the form of a disturbance in heart rate or developmental instability in inner ear organs.

<https://pubmed.ncbi.nlm.nih.gov/33530555/>

EXPOSURE TO 50 HZ EXTREMELY-LOW-FREQUENCY MAGNETIC FIELDS INDUCES NO DNA DAMAGE IN CELLS BY GAMMA H2AX TECHNOLOGY

Ye Lv, Shuchang Chen, Bing Zhu, Hong Xu, Shanshan Xu, Weiyan Liu, Yunyun Shen, Qunli Zeng. Exposure to 50 Hz Extremely-Low-Frequency Magnetic Fields Induces No DNA Damage in Cells by Gamma H2AX Technology. *Biomed Res Int.* 2021 Feb 15;2021:8510315. doi: 10.1155/2021/8510315.

Abstract

The current results for extremely-low-frequency magnetic fields (ELF-MF) on DNA damage are still debated. A sensitive indicator and systematic research are needed to assess the effects of ELF-MF. In this study, we used γ H2AX as an early and sensitive molecular marker to evaluate the DNA damage effects of ELF-MF in vitro.

Human amnion epithelial cells (FLs), human skin fibroblast cells (HSFs), and human umbilical vein endothelial cells (HUVECs) were exposed to 50 Hz ELF-MF at 0.4, 1, and 2 mT for 15 min, 1 h, and 24 h, respectively. After exposure, cells were subjected to γ H2AX immunofluorescence and western blot. The results showed no significant difference in the average number of foci per cell, the percentage of γ H2AX foci-positive cells, or the expression of γ H2AX between the sham and 50 Hz ELF-MF exposure groups ($P > 0.05$). In conclusion, 50 Hz ELF-MF did not induce DNA damage in FLs, HSFs, or HUVECs, which was independent of the intensity or duration of the exposure.

Open access paper: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7899753/>

STATIC MAGNETIC FIELD INDUCES ABNORMALITY OF GLUCOSE METABOLISM IN RATS' BRAIN AND RESULTS IN ANXIETY-LIKE BEHAVIOR

Tang Shuo, Ye Yumeng, Yang Leilei, Hao Yanhui, Yu Chao, Yan Hua, Xing Yuan, Jia Zhaoqian, Hu Cuicui, Zuo Hongyan, Li Yang. Static Magnetic Field Induces Abnormality of Glucose Metabolism in rats' brain and results in anxiety-like behavior. *J Chem Neuroanat.* 2021 Feb 4;101923. doi: 10.1016/j.jchemneu.2021.101923.

Abstract

In this study, fifty-four male Wistar rats were randomly divided into four groups according to the static magnetic field (SMF) intensity, namely, control, low-intensity, moderate-intensity, and high-intensity groups. The rats' whole body was exposed to a superconducting magnet exposure source. The exposure SMF intensity for the low-intensity, moderate-intensity, and high-intensity groups was 50 m T, 100 m T, and 200 m T, respectively, and the exposure time was 1 h/day for consecutive 15 days. After different exposure times, glucose metabolism in rats' brain was evaluated by micro-positron emission tomography (micro-PET), and the expression of hexokinase 1(HK1) and 6-phosphate fructokinase-1(PFK1) was detected by western blot. The exploration and locomotion abilities of the rats were evaluated by conducting open field test (OFT). Furthermore, pathological changes of rats' brain were observed under a microscope by using hematoxylin-eosin staining. PET results showed that moderate-intensity SMFs could cause fluctuant changes in glucose metabolism in rats' brain and the abnormalities were SMF intensity dependent. The expression of the two rate-limiting enzymes HK1 and PFK1 in glucose metabolism in brain significantly decreased after SMF exposure. The OFT showed that the total distance, surrounding distance, activity time, and climbing and standing times significantly decreased after SMF exposure. The main pathological changes in the brain were pyknosis, edema of neurons, and slight widening of the perivascular space, which occurred after 15 times of exposure. This study indicated that SMF exposure could lead to abnormal glucose metabolism in the brain and might result in anxiety-like behaviors.

<https://pubmed.ncbi.nlm.nih.gov/33549700/>

TURNING PREFERENCE IN DOGS: NORTH ATTRACTS WHILE SOUTH REPELS

Jana Adámková, Kateřina Benediktová, Jan Svoboda, Luděk Bartoš, Lucie Vynikalová, Petra Nováková, Vlastimil Hart, Michael S Painter, Hynek Burda. Turning preference in dogs: North attracts while south repels. *PLoS One.* 2021 Jan 28;16(1):e0245940. doi: 10.1371/journal.pone.0245940.

Abstract

It was shown earlier that dogs, when selecting between two dishes with snacks placed in front of them, left and right, prefer to turn either clockwise or counterclockwise or randomly in either direction. This preference (or non-preference) is individually consistent in all trials but it is biased in favor of north if they choose between dishes positioned north and east or north and west, a phenomenon denoted as "pull of the north". Here, we replicated these experiments indoors, in magnetic coils, under natural magnetic field and under magnetic field shifted 90° clockwise. We demonstrate that "pull of the north" was present also in an environment without any outdoor cues and that the magnetic (and not topographic) north

exerted the effect. The detailed analysis shows that the phenomenon involves also "repulsion of the south". The clockwise turning preference in the right- preferring dogs is more pronounced in the S-W combination, while the counterclockwise turning preference in the left-preferring dogs is pronounced in the S-E combination. In this way, south-placed dishes are less frequently chosen than would be expected, while the north-placed dishes are apparently more preferred.

Turning preference did not correlate with the motoric paw laterality (Kong test). Given that the choice of a dish is visually guided, we postulate that the turning preference was determined by the dominant eye, so that a dominant right eye resulted in clockwise, and a dominant left eye in counterclockwise turning. Assuming further that magnetoreception in canines is based on the radical-pair mechanism, a "conflict of interests" may be expected, if the dominant eye guides turning away from north, yet the contralateral eye "sees the north", which generally acts attractive, provoking body alignment along the north-south axis.

Open access paper:

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0245940>

ELECTROMAGNETIC FIELDS, 5G AND HEALTH: WHAT ABOUT THE PRECAUTIONARY PRINCIPLE?

John William Frank. Electromagnetic fields, 5G and health: what about the precautionary principle? *J Epidemiol Community Health*. Published Online First: 19 January 2021. doi: 10.1136/jech-2019-213595.

Abstract

New fifth generation (5G) telecommunications systems, now being rolled out globally, have become the subject of a fierce controversy. Some health protection agencies and their scientific advisory committees have concluded that there is no conclusive scientific evidence of harm. Several recent reviews by independent scientists, however, suggest that there is significant uncertainty on this question, with rapidly emerging evidence of potentially harmful biological effects from radio frequency electromagnetic field (RF-EMF) exposures, at the levels 5G roll-out will entail. This essay identifies four relevant sources of scientific uncertainty and concern: (1) lack of clarity about precisely what technology is included in 5G; (2) a rapidly accumulating body of laboratory studies documenting disruptive in vitro and in vivo effects of RF-EMFs—but one with many gaps in it; (3) an almost total lack (as yet) of high-quality epidemiological studies of adverse human health effects from 5G EMF exposure specifically, but rapidly emerging epidemiological evidence of such effects from past generations of RF-EMF exposure; (4) persistent allegations that some national telecommunications regulatory authorities do not base their RF-EMF safety policies on the latest science, related to unmanaged conflicts of interest. The author, an experienced epidemiologist, concludes that one cannot dismiss the growing health concerns about RF-EMFs, especially in an era when higher population levels of exposure are occurring widely, due to the spatially dense transmitters which 5G systems require. Based on the precautionary principle, the author echoes the calls of others for a moratorium on the further roll-out of 5G systems globally, pending more conclusive research on their safety.

Conclusions and Recommendation

In assessing causal evidence in environmental epidemiology, Bradford Hill himself pointed out that ‘the whole picture matters;’ he argued against prioritising any subset of his famous nine criteria for causation. One’s overall assessment of the likelihood that an exposure causes a health condition should take into account a wide variety of evidence, including ‘biological plausibility’. 34 35 After reviewing the evidence cited above, the writer, an experienced physician-epidemiologist, is convinced that RF-EMFs may well have serious human health effects. While there is also increasing scientific evidence for RF-EMF effects of ecological concern in other species, 6–8 16–18 23 both plant and animal, these have not been reviewed here, for reasons of space and the author’s disciplinary limitations. In addition, there is convincing evidence, cited above, that several nations’ regulatory apparatus, for telecommunications innovations such as the 5G roll-out, is not fit for purpose. Indeed, significant elements in that apparatus appear to have been captured by vested interests. Every society’s public health—and especially the health of those most likely to be susceptible to the hazard in question (in the case of EMFs, children and pregnant women)—needs to be protected by evidence-based regulations, free from significant bias.

Finally, this commentary would be remiss if it did not mention a widely circulating conspiracy theory, suggesting that 5G and related EMF exposures somehow contributed to the creation or spread of the current COVID-19 pandemic. There are knowledgeable commentators’ reports on the web debunking this theory, and no respectable scientist or publication has backed it. 40 41 Indeed, combatting it is widely viewed by the scientific community as critical to dealing with the pandemic, as conspiracy theorists holding this view have already carried out violent attacks on mobile phone transmission facilities and other symbolic targets, distracting the public and authorities at a time when pandemic control actions are paramount. 42 This writer completely supports that view of the broader scientific community: the theory that 5G and related EMFs have contributed to the pandemic is baseless.

It follows that, for the current 5G roll-out, there is a sound basis for invoking ‘the precautionary principle’. 43 This is the environmental and occupational health principle by which significant doubt about the safety of a new and potentially widespread human exposure should be a reason to call a moratorium on that exposure, pending adequate scientific investigation of its suspected adverse health effects. In short, one should ‘err on the side of caution’. In the case of 5G transmission systems, there is no compelling public health or safety rationale for their rapid deployment. The main gains being promised are either economic (for some parties only, not necessarily with widely distributed financial benefits across the population) or related to increased consumer convenience. Until we know more about what we are getting into, from a health and ecological point of view, those putative gains need to wait.

Open access paper: <https://jech.bmj.com/content/early/2021/01/04/jech-2019-213595> or <https://jech.bmj.com/content/jech/early/2021/01/04/jech-2019-213595.full.pdf>

EFFECTS OF ELECTROMAGNETIC WAVES WITH LTE AND 5G BANDWIDTH ON THE SKIN PIGMENTATION IN VITRO

Kyuri Kim, Young Seung Lee, Nam Kim, Hyung-Do Choi, Dong-Jun Kang, Hak Rim Kim, Kyung-Min Lim. Effects of Electromagnetic Waves with LTE and 5G Bandwidth on the Skin Pigmentation In Vitro. *Int J Mol Sci.* 2020 Dec 26;22(1):E170. doi: 10.3390/ijms22010170.

Abstract

With the rapid growth of wireless communication devices, the influences of electromagnetic fields (EMF) on human health are gathering increasing attention. Since the skin is the largest organ of the body and is located at the outermost layer, it is considered a major target for the health effects of EMF. Skin pigmentation represents one of the most frequent symptoms caused by various non-ionizing radiations, including ultraviolet radiation, blue light, infrared, and extremely low frequency (ELF). Here, we investigated the effects of EMFs with long-term evolution (LTE, 1.762 GHz) and 5G (28 GHz) bandwidth on skin pigmentation in vitro. Murine and Human melanoma cells (B16F10 and MNT-1) were exposed to either LTE or 5G for 4 h per day, which is considered the upper bound of average smartphone use time. It was shown that neither LTE nor 5G exposure induced significant effects on cell viability or pigmentation. The dendrites of MNT-1 were neither lengthened nor regressed after EMF exposure. Skin pigmentation effects of EMFs were further examined in the human keratinocyte cell line (MNT-1-HaCaT) co-culture system, which confirmed the absence of significant hyperpigmentation effects of LTE and 5G EMFs. Lastly, MelanoDerm™, a 3D pigmented human epidermis model, was irradiated with LTE (1.762 GHz) or 5G (28 GHz), and image analysis and special staining were performed. No changes in the brightness of MelanoDerm™ tissues were observed in LTE- or 5G-exposed tissues, except for only minimal changes in the size of melanocytes. Collectively, these results imply that exposure to LTE and 5G EMFs may not affect melanin synthesis or skin pigmentation under normal smartphone use condition.

Open access paper: <https://www.mdpi.com/1422-0067/22/1/170>

ASSOCIATION BETWEEN REPRODUCTIVE HEALTH AND NONIONIZING RADIATION EXPOSURE

Pooja Negi, Rajeev Singh. Association between reproductive health and nonionizing radiation exposure. *Electromagnetic Biology and Medicine.* Published online: 20 Jan 2021. DOI: 10.1080/15368378.2021.1874973.

Abstract

Recently, a decreasing rate of fertility has to be credited to an array of factors such as environmental, health and lifestyle. Male infertility is likely to be affected by the strong exposure to heat and radiations. The most common sources of nonionizing radiations are cell phones, laptops, Wi-Fi and microwave ovens, which may participate to the cause of male infertility. One of the major sources of daily exposure to non-ionizing radiation is mobile phones. A mobile phone is now basically dominating our daily life through better services such as connectivity, smartphone devices. However, the health consequences are linked with their usage are frequently ignored.

Constant exposure to non-ionizing radiations produced from a cell phone is one of the possible reasons for growing male infertility. Recently, several studies have shown that cell

phone users have altered sperm parameters causing declining reproductive health. Cell phone radiation harms male fertility by affecting the different parameters like sperm motility, sperm count, sperm morphology, semen concentration, morphometric abnormalities, increased oxidative stress along with some hormonal changes. This review is focusing on the prevailing literature from in vitro and in vivo studies suggesting that non-ionizing exposure negatively affects human male infertility. <https://pubmed.ncbi.nlm.nih.gov/33471575/>

Conclusion

Generally, the outcome of the studies has indicated that mobile phone usage changes different sperm parameters in both ways in-vitro (human) and in-vivo (animals). Several studies disclose that the exposure to cell phones produces harmful effects on the testes, which may affect sperm motility, sperm number, sperm concentration, and morphology and an increased DNA damage, causing micronuclei formation and reactive oxygen species within the cell. So many evidences showed that exposure from cell phones results in elevated oxidative stress with disintegrated DNA and it is directly and indirectly dependent on the time of cell phone use.

Further researches are required to provide strong evidence that the use of mobile phones may disturb sperm and testicular activity. Several evidences suggest that the irregularities reported due to RF-EMF-exposure depend on physical parameters such as utilized RF wavelength, penetration range into the object, and transmission length of the radiation. Unfortunately, existing studies are not able to suggest a true mechanism between the harmful effects of RF-EMF radiation and the male reproductive system. To conclude all of the above, government bodies and agencies should form strong guidelines against cell phone exposure and take preventive actions such as in the usage of mobile phones, preventing chatting, reducing the overall contact time, and holding the gadget away from the groin may be of significant help to people pursuing fertility. Moreover, very limited studies are available on protective actions so far so a large-scale analysis is also required to determine the reproductive parameters.

IMPLICATIONS OF MMWAVE RADIATION ON HUMAN HEALTH: STATE OF THE ART THRESHOLD LEVELS

Ravilla Dilli. Implications of mmWave Radiation on Human Health: State of the Art Threshold Levels. IEEE Access. 18 January 2021. DOI: 10.1109/ACCESS.2021.3052387.

Abstract

millimeter (mmWave) frequencies are covering from 30GHz to 300GHz in the electromagnetic spectrum and their uses in various applications like next-generation wireless communication systems (massive 5G telecommunications network), medical devices, airport security and automatic collision avoidance systems are growing vastly in the near future. Therefore, it is important to study the effects of mmWave radiation (non- ionization radiation) on biological systems and biophysical mechanisms. This paper focus on thorough review of nascent literature about current understandings of biological effects and epidemiological studies due to mmWave radiation in human beings. It presents latest guidelines with quantitative electromagnetic field thresholds by considering the realistic exposure scenarios of “general public” and “occupational” who undergo through wireless communication sources in their daily life. It also gives necessary safety measures to be taken while using the emerging

mmWave technologies for future generation wireless communication networks.
<https://ieeexplore.ieee.org/document/9328127>

Conclusions and Futurescope

This paper has investigated mainly on the latest and current research work on various adverse health effects in human beings due to continuous and discontinuous, short-term and long-term RF EMFs exposures at mmWave frequency bands. It has mentioned the state-of-art threshold levels for exposures to RF EMFs at mmWave frequency bands. However, there are few exposure scenarios like RF EMFs interfering with electrical equipment (also called EMC), potential harms to volunteer research participants, EMFs due to metallic implants which are part of medical treatment are out of scope to the threshold levels mentioned in this paper. The future scope of this work is to establish guidelines and safety measures at THz frequency bands, also consider the EMC influences in defining threshold values. Currently, the EHS health disorders are greatly increasing day by day in people who use smart phones for a long time in a day. Future continuations of this research work include deriving scientific basis which gives relation between the usage of mmWave communication devices and EHS health disorders in humans.

REDEFINING ELECTROSENSITIVITY: A NEW LITERATURE-SUPPORTED MODEL

Mary Redmayne, Siobhan Reddel. Redefining electrosensitivity: Jan 2021. DOI: 10.1080/15368378.2021.1874971.

Abstract

In critically examining literature on electrohypersensitivity and the reported somatic responses to anthropogenic modulated radiofrequency radiation (RFR) exposure, it becomes apparent that electrohypersensitivity is one part of a range of consequences. Current evidence on the necessity of considering patients' overall health status leads us to propose a new model in which electrohypersensitivity is but part of the electrosensitive status inherent in being human. We propose the likelihood and type of response to environmental RFR include i) a linear somatic awareness continuum, ii) a non-linear somatic response continuum, and iii) the extent of each individual's capacity to repair damage (linked to homeostatic response). We anticipate this last, dynamic, aspect is inextricably linked to the others through the autonomic nervous system. The whole is dependent upon the status of the interconnected immune and inflammatory systems. This holistic approach leads us to propose various outcomes. For most, their body maintains homeostasis by routine repair. However, some develop electrohypersensitivity either due to RFR exposure or as an ANS-mediated, unconscious response (aka nocebo effect), or both. We suggest RFR exposure may be one factor in the others developing an auto-immune disease or allergy. A few develop delayed catastrophic disease such as glioma. This model gives the blanket term ElectroMagnetic Illness (EMI) to all RFR-related conditions. Thus, EHS appears to be one part of a range of responses to a novel and rapidly changing evolutionary situation.

Conclusion and consequences of the model

Overall, there is compelling evidence in the broader literature indicating that EHS is but one part of a more complex range of responses related to humans being intrinsically electrosensitive.

In fact, “Many of the general malaise symptoms associated with IEI (Idiopathic Environmental Intolerance) 7 are common in the general population” p.42 (World Health Organisation 2004). This is not a sign of a healthy general population even by the WHO’s definition of health, 8 and causes must be identified. Life is dependent upon, and arguably defined by the electrical activity, of the brain, the broader nervous system and indeed the body’s ongoing efforts to maintain homeostasis. As we said earlier, internal electrical impulses are inherent to life; when they stop, we die. There are also natural, static and low-frequency, non-modulated, electromagnetic fields with which we have evolved and most of which we use and even rely upon. Our bodies and cells have their own resonant frequencies, and research has shown cell sensitivity to RFR exposure at ultra-low intensities (Kositsky, 2001). It is almost to be expected that exposures never encountered during the course of evolution would affect us. Indeed, “Since the low-level environmental exposures associated with IEI are commonly found in everyone’s daily living environment, the exposed population is indistinguishable from the general population.

Everyone is exposed” p.44 (World Health Organisation 2004).

Our current model of electrosensitivity includes the “general population” and expresses that differing responses are a normal part of being alive. For some of those who do not appear to respond (no symptoms), there will mostly likely be cellular impacts. For most, these will be promptly repaired or otherwise dealt with (e.g., apoptosis); for some, damage will gradually or eventually accumulate and may lead, in time, to a catastrophic response in the form of a disease that began as oxidative stress that the body has fought but finally to which it has catastrophically succumbed. For others, effects will be felt and cause early symptoms. Among these people, some will respond by avoiding the exposure either subconsciously or consciously, while others will ignore them and eventually pay the price. Others will develop EHS. At this stage, it will often go undiagnosed due to lack of training in recognising this possible diagnosis.

Importantly, this model implies that there is a need to acknowledge that EHS (with all its variety) is one part of a range of symptomless pre-cursor and EMI responses to a novel and rapidly changing evolutionary situation.

There should be no stigma to EHS, and those affected deserve the same level of concern and extent of assistance as those with allergies or auto-immune diseases, the principal actions being minimisation of the harmful stimulant until personal repair capacity has improved, then slowly re-introducing exposure, to the extent manageable on an individual basis. Assistance with this would be necessary at several stages so the body is not pushed into the hypo-responsive state, but rather assisted to reach the best possible health with the best practicable repair capacity.

We propose that to work towards avoiding, ameliorating, or attempting to overcome EHS it is necessary not only to observe the positive actions recommended, but also to build capacity. And capacity provides the energy necessary to build resilience (Gerritsen and Band 2018).

<https://www.tandfonline.com/doi/full/10.1080/15368378.2021.1874971>

THE IMPACT OF MOBILE PHONE USE ON TINNITUS: A SYSTEMATIC REVIEW AND META-ANALYSIS

Artur Kacprzyk, Tomasz Stefura, Marta Krzysztofik, Tomasz Rok, Eugeniusz Rokita, Grzegorz Tatoń. The Impact of Mobile Phone Use on Tinnitus: A Systematic Review and Meta-Analysis. Bioelectromagnetics. 2021 Jan 13. doi: 10.1002/bem.22316.

Abstract

Tinnitus is a perception of sound in the absence of an external source. The aim of our study was to investigate with a meta-analytical approach, whether mobile phone (MP) use increases the risk of tinnitus. Eight studies reporting the risk of tinnitus in relation to MP use were identified, and six high-quality studies (two cohort studies, one case-control study, and three cross-sectional ones) were included in the meta-analysis. The quality assessment was performed using the Newcastle-Ottawa scale. The risk of tinnitus was analyzed depending on the exposure to MPs in subgroups according to the study design and method of exposure assessment. Two cohort studies, which assessed the exposure to MPs using network operator data, indicated no significantly increased risk of tinnitus among highly exposed MP users in comparison to lightly exposed individuals (odds ratio [OR]: 1.03 [95% confidence interval [CI]: 0.93-1.15]). Likewise, the self-reported exposure data from two cohorts/case-control and four cross-sectional studies did not find an association between exposure to MPs and tinnitus (OR: 1.20 [95% CI: 0.40-3.61] and OR: 1.73 [95% CI: 0.67-4.49], respectively). Current scientific knowledge, including high-quality studies with a reliable exposure assessment based on network operator data, does not support the hypothesis that MP use is associated with tinnitus.

<https://pubmed.ncbi.nlm.nih.gov/33440459/>

IS THERE EVIDENCE FOR OXIDATIVE STRESS CAUSED BY ELECTROMAGNETIC FIELDS? A SUMMARY OF RELEVANT OBSERVATIONS IN EXPERIMENTAL ANIMAL AND CELL EXPERIMENTS RELATED TO HEALTH EFFECTS IN THE LAST TEN YEARS

Prof Meike Mevissen, David Schürmann. Is there evidence for oxidative stress caused by electromagnetic fields? A summary of relevant observations in experimental animal and cell experiments related to health effects in the last ten years. BERENIS – The Swiss expert group on electromagnetic fields and non-ionising radiation Newsletter – Special Issue January 2021.

Introduction

This special issue of the BERENIS newsletter contains an up-to-date assessment of a possible correlation between oxidative stress and exposure to magnetic and electromagnetic fields and their putative effects on health. For this purpose, relevant animal and cell studies published between 2010 and 2020 were identified and summarized. An extended report presenting these recent studies in more detail will be published soon by the FOEN¹. This special issue contains a short version of the report.

Conclusions

In summary, the majority of the animal and more than half of the cell studies provided evidence of increased oxidative stress caused by RF-EMF or ELF-MF. This notion is based on observations in a large number of cell types, applying different exposure times and dosages (SAR or field strengths), also in the range of the regulatory limits. Certainly, some studies are burdened with methodological uncertainties and weaknesses or are not very comprehensive

in terms of exposure time, dose, number and quantitative analysis of the biomarkers used, to name a few. Taking these methodological weaknesses into account, nonetheless, a tendency becomes apparent, namely that EMF exposure, even in the low dose range, can lead to changes in oxidative balance. Organisms and cells are generally able to react to oxidative stress, and many studies showed adaptation to EMF exposure after a recovery phase. Pre-existing conditions, such as immune deficiencies or diseases (diabetes, neurodegenerative diseases), compromise the body's defence mechanisms, including antioxidative protection, and it is therefore possible that individuals with these conditions experience more severe health effects. In addition, the studies show that very young and elderly individuals can react less efficiently to oxidative stress induced by EMF, which of course also applies to other stressors that cause oxidative stress. More extensive studies under standardised conditions are necessary, to better understand and confirm these phenomena and observations.

Open access report: <https://t.co/ubvIFOnyXd?amp=1>

BERENIS - SWISS EXPERT GROUP ON ELECTROMAGNETIC FIELDS AND NON-IONISING RADIATION

The FOEN has nominated a consultative group of Swiss experts from various disciplines with scientific competence regarding electromagnetic fields and NIR, which has commenced its work in July 2014. The group is called "BERENIS", based on the acronym of the respective German term. The BERENIS experts regularly screen the scientific literature, and assess the publications which they consider relevant for the protection of humans from potentially adverse effects.

The Swiss expert group on electromagnetic fields and non-ionising radiation (BERENIS)

In Switzerland, the Federal Office for the Environment (FOEN) is the responsible government body for monitoring and assessing research on health effects of non-ionising radiation (NIR) from stationary sources in the environment. This includes informing and updating the public about the current state of research, which is the basis for the ambient regulatory limits stated in the Swiss "ordinance relating to protection from non- ionising radiation". In the case of reliable new scientific knowledge and experiences, the FOEN would advise the Federal Council of Switzerland to adapt these ambient regulatory limits.

Assessing the results and conclusions of scientific studies enables early detection of potential health risks of NIR. The FOEN places particular emphasis to not overlook any evidence of harmfulness for public health demanding for corrective regulatory interventions. Furthermore, critical assessment of available scientific data is required to make firm statements about the validity of the presented evidence for biological effects, their relevance for public health, and if so, to estimate the number of potentially affected people.

NIR includes a broad spectrum of frequencies with varying intensities and radiation characteristics, which is becoming more complex and multifaceted as the development and application of technologies emitting them is very dynamic and rapidly increasing. In the context of the work of BERENIS, NIR at frequencies below 10 GHz is addressed. Potential biological effects and the underlying mechanisms of NIR are manifold, and research activities

range from the molecular to the population level. As a consequence, specific expertise in various disciplines is needed to assess the related scientific studies rooted in many different biological, medical and technical scientific realms.

The FOEN has therefore nominated a consultative group of Swiss experts from various disciplines with scientific competence regarding electromagnetic fields and NIR, which has commenced its work in July 2014. The group is called "BERENIS", based on the acronym of the respective German term. The BERENIS experts regularly screen the scientific literature, and assess the publications which they consider relevant for the protection of humans from potentially adverse effects. The results of this evaluation are published in quarterly newsletters, which can be downloaded from this webpage.

Members of BERENIS

- Prof. Dr. Martin Rössli, Swiss Tropical and Public Health Institute, Basel (in charge)
- Prof. Dr. Peter Achermann, The KEY Institute for Brain-Mind Research, Zurich
- Dr. Jürg Fröhlich, Fields at Work GmbH, Zurich
- Prof. Dr. med. Jürg Kesselring, previous head of the Department of Neurorehabilitation, Rehabilitation Centre, Clinics of Valens
- Prof. Dr. Meike Mevissen, Vet-Pharmacology & Toxicology, University of Bern
- Dr. David Schürmann, Molecular Genetics Group, Department of Biomedicine, University of Basel
- Dr. med. Edith Steiner, Ärztinnen und Ärzte für Umweltschutz, Basel Secretariat:
- Dr. Stefan Dongus, Swiss Tropical and Public Health Institute, Basel Federal Office for the Environment:
- Alexander Reichenbach
- Dr. Maurane Riesen Observers:
- Dr. Evelyn Stempfel (Federal Office of Public Health)
- Dr. Roland Krischek (Swiss National Accident Insurance Fund)
- Dr. Christian Monn (State Secretariat for Economic Affairs)
- Dr. Samuel Iff (State Secretariat for Economic Affairs)

EMPIRICAL STUDY ON SPECIFIC ABSORPTION RATE OF HEAD TISSUES DUE TO INDUCED HEATING OF 4G CELL PHONE RADIATION

Christopher B, Mary S, Khandaker MU, Jojo PJ. 178(Special Issue): 108910. Jan 2021. DOI:10.1016/j.radphyschem.2020.108910.

Abstract

Exposures to electromagnetic radiation mainly from the extended use of mobile phones may initiate biological damages in the human body at the macromolecular level. Several studies on human and animal models have shown significant changes in the functions of neural cells. Present empirical study analyses the thermal changes and the specific absorption rates (SAR) of brain, eye and skin tissues due to prolonged exposure to mobile phone radiation. A phantom, simulating human head with skin, skull and brain was used for the study. The Phantom was exposed to radiation for longer durations (600 s and more) and the temperature variations at different specific points were studied with sensitive thermocouple probes. SAR (1 g of contiguous tissue) values were determined using the variations of temperature and other parameters. The average rise in brain temperature was found to be 0.10 +/- 0.05

degrees C at 30 mm deep in the brain and the estimated SAR was 0.66 +/- 0.35 Wkg(-1). The increase in temperature for the eye socket was 0.03 +/- 0.02 degrees C with SAR 0.15 +/- 0.08 Wkg(-1). The average rise in temperature for skin was 0.14 +/- 0.05 degrees C and the SAR was 0.66 +/- 0.42 Wkg(-1). Although the measured SAR lie within the safe limit of 2 Wkg(-1) recommended by the international regulatory body, considering the tremendous growth in the number of mobile phone users and prolonged use of mobile phone in communication purposes, the cumulative effects could be a real concern for human health.

ASSOCIATION BETWEEN ESTIMATED WHOLE-BRAIN RADIOFREQUENCY ELECTROMAGNETIC FIELDS DOSE AND COGNITIVE FUNCTION IN PREADOLESCENTS AND ADOLESCENTS

Cabre-Riera A, van Wel, L, Liorni, I, Thielens A, Birks LE, Pierotti L, Joseph W, Gonzalez-Safont L, Ibarluzea J, Ferrero A, et al. *Int J Hygiene Envir Health*. 231:113659. Jan 2021. DOI: 10.1016/j.ijheh.2020.113659.

Abstract

Objective: To investigate the association between estimated whole-brain radiofrequency electromagnetic fields (RF-EMF) dose, using an improved integrated RF-EMF exposure model, and cognitive function in preadolescents and adolescents.

Methods: Cross-sectional analysis in preadolescents aged 9-11 years and adolescents aged 17-18 years from the Dutch Amsterdam Born Children and their Development Study (n = 1664 preadolescents) and the Spanish INfancia y Medio Ambiente Project (n = 1288 preadolescents and n = 261 adolescents), two population-based birth cohort studies. Overall whole-brain RF-EMF doses (mJ/kg/day) were estimated for several RF-EMF sources together including mobile and Digital Enhanced Cordless Telecommunications phone calls (named phone calls), other mobile phone uses than calling, tablet use, laptop use (named screen activities), and far-field sources. We also estimated whole-brain RF-EMF doses in these three groups separately (i.e. phone calls, screen activities, and far-field) that lead to different patterns of RF-EMF exposure. We assessed non-verbal intelligence in the Dutch and Spanish preadolescents, information processing speed, attentional function, and cognitive flexibility in the Spanish preadolescents, and working memory and semantic fluency in the Spanish preadolescents and adolescents using validated neurocognitive tests.

Results: Estimated overall whole-brain RF-EMF dose was 90.1 mJ/kg/day (interquartile range (IQR) 42.7; 164.0) in the Dutch and Spanish preadolescents and 105.1 mJ/kg/day (IQR 51.0; 295.7) in the Spanish adolescents. Higher overall estimated whole-brain RF-EMF doses from all RF-EMF sources together and from phone calls were associated with lower non-verbal intelligence score in the Dutch and Spanish preadolescents (-0.10 points, 95% CI-0.19; -0.02 per 100 mJ/kg/day increase in each exposure). However, none of the whole-brain RF-EMF doses was related to any other cognitive function outcome in the Spanish preadolescents or adolescents.

Conclusions: Our results suggest that higher brain exposure to RF-EMF is related to lower non-verbal intelligence but not to other cognitive function outcomes. Given the cross-sectional nature of the study, the small effect sizes, and the unknown biological mechanisms, we cannot discard that our results are due to chance finding or reverse causality. Longitudinal studies on RF-EMF brain exposure and cognitive function are needed.

THE EFFECT OF MOBILE PHONE WAVES ON SALIVARY SECRETION IN DENTAL STUDENTS OF RAFSANJAN UNIVERSITY OF MEDICAL SCIENCES

N. Kamyab, Z. Mordouei, M. Hosseini, M. Sheikh Fathollahi. The effect of mobile phone waves on salivary secretion in dental students of Rafsanjan University of Medical Sciences.

Int J Radiat Res 2021, 19(1): 81-87.

Abstract

Background: Excessive use of mobile phones is addictive and causes many complications. The severity of these complications depends on the duration of the call and the frequency of using mobile phones. Therefore, this study aimed to determine the effect of mobile phone waves on salivary secretion.

Materials and Methods: This descriptive study was conducted on 120 dental students in Rafsanjan, Iran, in 2015. Using the call history of the mobile phones, the number and duration of each person's daily calls were obtained. The spitting method was used to collect unstimulated saliva, and stimulated saliva was collected with the help of natural chewing gums without sugar and additive flavors. After sampling, the falcon tubes containing saliva were centrifuged at 2500 rpm for 10 minutes. Then the saliva volume was read from the tube and recorded. Independent two-sample t test, one-way ANOVA, and multiple linear regression model were used for data analysis.

Results: Students who talked for more than 40 minutes or 20-40 minutes a day on their mobile phones had a mean unstimulated salivary secretion of 0.885 and 0.331 ml/min, respectively and mean stimulated salivary secretion of 0.702 and 0.708 ml/min, respectively more than students who talked less than 20 minutes a day on their cell phones ($P < 0.05$).

Conclusion: The results showed that increased duration of cell phone calls was associated with increasing the amount of stimulated and unstimulated salivary secretion, which could decrease the risk of developing oral infections and tooth decay.

<http://ijrr.com/article-1-3459-en.html>

Conclusion

The results of the present study indicated that increased duration of cell phone calls was correlated to increasing the amount of salivary secretion. It is suggested future studies be conducted on all age groups and in addition to quantity, the quality of saliva is examined. Measuring the amount of saliva secreted on the dominant side of cell phone use and compare it with the recessive side is also recommended.

MEASURING THE IMPACT OF ICNIRP VS. STRICTER-THAN-ICNIRP EXPOSURE LIMITS ON QoS AND EMF FROM CELLULAR NETWORKS

Jaime Galán-Jiménez, Luca Chiaraviglio. Measuring the impact of ICNIRP vs. stricter-than-ICNIRP exposure limits on QoS and EMF from cellular networks. *Computer Networks*. 6 January 2021. 107824. <https://doi.org/10.1016/j.comnet.2021.107824>.

Abstract

The installation of new equipment (Base Stations, BSs) during the planning phase of a cellular network (including 5G BSs) is governed by exposure limits in terms of allowable ElectroMagnetic Field (EMF) levels. The exposure limits can be either defined by i) international bodies (e.g., ICNIRP) or ii) national regulations imposing limits stricter than i). In this work, we compare the impact of ICNIRP vs. stricter-than-ICNIRP exposure regulations on the Quality of Service (QoS) and EMF. To this aim, we perform a large-scale measurement campaign in one scenario in Spain subject to ICNIRP regulations and another one in Italy subject to EMF limits stricter than ICNIRP ones. Both the scenarios are characterized by similar exposure conditions, comparable user density, and common 4G performance targets by the operators. Results, obtained by measuring QoS and EMF at selected locations, reveal that the QoS in the scenario subject to strict EMF limits is heavily worsened compared to the one in which ICNIRP-based limits are enforced. Clearly, the scenario with strict EMF limits presents a lower level exposure over the territory compared to the one imposing ICNIRP limits.

<https://www.sciencedirect.com/science/article/abs/pii/S1389128621000128>

EFFECTS OF 2600 MHZ RADIOFREQUENCY RADIATION IN BRAIN TISSUE OF MALE WISTAR RATS AND NEUROPROTECTIVE EFFECTS OF MELATONIN

Kevser Delen, Bahriye Sırav, Sinem Oruç, Cemile M Seymen, Dilek Kuzay, Korkut Yeğin, Gülnur Take Kaplanoğlu. Effects of 2600 MHz Radiofrequency Radiation in Brain Tissue of Male Wistar Rats and Neuroprotective Effects of Melatonin. *Bioelectromagnetics*. 2021 Jan 13. doi: 10.1002/bem.22318.

Abstract

The debate on the biological effects of radiofrequency radiation (RFR) still continues due to differences in the design of studies (frequency, power density, specific absorption rate [SAR], exposure duration, cell, tissue, or animal type). The current study aimed to investigate the effects of 2,600 MHz RFR and melatonin on brain tissue biochemistry and histology of male rats. Thirty-six rats were divided into six groups randomly: cage-control, sham, RFR, melatonin, sham melatonin, and RFR melatonin. In RFR groups, animals were exposed to 2,600 MHz RFR for 30 days (30 min/day, 5 days/week) and the melatonin group animals were subcutaneously injected with melatonin (7 days/week, 10 mg/kg/day) for 30 days. SAR in brain gray matter was calculated as 0.44 and 0.295 W/kg for 1 and 10 g averaging, respectively. RFR exposure decreased the GSH, GSH-Px, and SOD levels and increased the MPO, MDA, and NOx levels ($P < 0.005$) significantly. RFR exposure also led to an increase in structural deformation and apoptosis in the brain tissue. This study revealed that exogenous high-dose melatonin could reduce these adverse effects of RFR. Limiting RFR exposure as much as possible is recommended, and taking daily melatonin supplements may be beneficial.

<https://pubmed.ncbi.nlm.nih.gov/33440456/>

THE EFFECT OF 900-MHZ RADIOFREQUENCY ELECTROMAGNETIC FIELDS DURING THE ADOLESCENCE ON THE HISTOLOGICAL STRUCTURE OF RAT TESTIS AND ITS ANDROGEN AND ESTROGEN RECEPTORS LOCALIZATION

Gur FM, Keles AI, Erol HS, Guven C, Taskin E, Kaya H, Gur HE, Odaci E, Halici MB, Timurkaan S. Int J Radiation Research. 19(1):135-144. Jan 2021. DOI: 10.18869/acadpub.ijrr.19.1.135

Abstract

Background: Mobile phones as an electronic device which are emitting radiofrequency-electromagnetic field (RF-EMF). In this study was intend to determine the contingent effects of cell phone induced RF-EMF on testicular tissue in adolescence.

Materials and Methods: Rats in the RF-EMF group were exposed to 900 MHz RF-EMF, while sham and control rats were not. After the completion of the test steps, the testicular tissues which were rapidly removed from the body of sacrificed rats were examined by using histopathological and biochemical methods. Testicular tissues cut to 5 mu m thickness undergo routine histological procedures. Thus, histopathological evaluation will be completed. Malondialdehyde (MDA), glutathione (GSH), catalase (CAT) and superoxide dismutase (SOD) levels in testicular tissues were measured by biochemical methods to determine whether oxidative stress occurred or not.

Results: Histopathologic findings were not observed in any of the studied groups. There was also no difference between the groups in terms of localization of androgen and estrogen receptors. The incidence of apoptotic index and TUNEL-positive cells was similar in all three groups. In the RF-EMF group, although the levels of MDA and CAT were significantly higher, GSH levels were lower than the other groups. There was no essential difference between the groups in terms of SOD level.

Conclusions: The obtained results of this study showed that exposed to 900 MHz RF-EMF in adolescents caused oxidative stress in the testes, but testicular damage which is caused by oxidative stress remained too low to be detected by histopathological methods in this study.

SENSITIVITY OF PLANTS TO HIGH FREQUENCY ELECTROMAGNETIC RADIATION: CELLULAR MECHANISMS AND MORPHOLOGICAL CHANGES

Shalinder Kaur, Alain Vian, Shikha Chandel, Harminder PalSingh, Daizy Rani Batish, Ravinder Kumar Kohli. Rev Environ Sci Biotechnol (2021). <https://doi.org/10.1007/s11157-020-09563-9>.

Abstract

The technological advancement and increased usage of wireless and other communication devices have greatly enhanced the level of radiofrequency electromagnetic field radiation (EMF-r) in the environment. It has resulted in unprecedented increased exposure of living organisms to these radiations. Most of the studies in past have, however, focused on animal systems and comparatively less attention has been paid to plants with studies reporting various, sometimes contradictory effects. This review is an attempt to provide a critical appraisal of the available reports regarding the impacts of these radiations on plant development and the underlying physiological, biochemical, and molecular mechanisms

involved. Here, we propose that the main entry point for the biological effects of EMF-r corresponds to an increase in ROS metabolism and cytosolic calcium that leads to various cellular responses including changes in gene expression and/or enzymatic activities, which could ultimately result in immediate cellular alterations or delayed plant growth. This may constitute a new perspective in the interpretation of plant responses to EMF-r exposure. Understanding the impacts of EMF-r and the inherent abilities of plants to cope up with such changes should lead to EMF-r being considered as full-fledged environmental signals that are perceived by the plants and integrated into their development patterns.

Conclusion and perspectives

Over the course of past decades, the extensive applications of EMF-r producing devices, and their potential to induce biological effects, have encouraged scientists to investigate the possible mechanisms of their action. Few studies have documented the progressive impacts of EMF-r on biota; however, studies with appropriate methodology suggested biological effects that require additional experimentation to understand their integration into plant development (rather than describing them in terms of positive or negative effects, as often found in the literature). The contradictory outcomes of studies suggest that the effects of EMF-r may be highly dependent upon exposure conditions (power density, frequency, and duration) and are species specific. A standardization of the procedure in use to expose plant to EMF-r, at least for model plants (arabidopsis, tomato, wheat, maize...) and for common frequencies (900, 1800 and MHz) would be highly valuable to allow a better comparison of the measured biological effects. However, the initial interaction and mechanism of EMF-r with plant tissue (the “primo-interaction”) is not yet understood, even if several putative mechanisms have been proposed. These include dipole transition of polar structures (Amat et al. 2006), forced vibration of free ions (Panagopoulos et al. 2000, 2002) or modification of ligand binding capacity (Chiabrera et al. 2000). These uncertainties make difficult the elaboration of efficient strategies to characterize the complexity of the plant response. The literature emphasizes that EMF-r interfere with the growth and development of plants at the molecular or whole plant level, clearly involving some factors (calcium, ROS, secondary metabolites, ATP) of plant responses to environmental cues. There are convincing evidences to consider EMF-r as real environmental signals’ that plants possibly integrate into their development. Nevertheless, in the real environment, EMF-r induced stress is certainly of secondary importance in comparison with other more serious stresses for plants (drought, pathogen attack, wind, etc.). However, an unintended consequence is that a constant level of exposure to electromagnetic fields may condition plants to respond secondarily more efficiently to a severe stress, installing a kind of memory in the plant (Thellier et al. 2000; Hilker et al. 2016). This hypothesis would be worth testing experimentally and may have valuable application in agriculture. In this perspective, global approaches to plant responses to EMF-r exposure (RNA sequencing, proteomics, metabolomics, DNA methylation, etc.) are still too few in the present literature for a more exhaustive knowledge of the metabolic pathways affected by exposure to EMF-r and should be investigated/deciphered in experimental designs. <https://link.springer.com/article/10.1007/s11157-020-09563-9>

ENVIRONMENTAL RISK FACTORS AND HEALTH: AN UMBRELLA REVIEW OF META-ANALYSES

Rojas-Rueda D, Morales-Zamora E, Alsufyani WA, Herbst CH, AlBalawi SM, Alsukait R, Alomran M. International Journal of Environmental Research and Public Health. 2021; 18(2):704.

Abstract

Background: Environmental health is a growing area of knowledge, continually increasing and updating the body of evidence linking the environment to human health.

Aim: This study summarizes the epidemiological evidence on environmental risk factors from meta-analyses through an umbrella review.

Methods: An umbrella review was conducted on meta-analyses of cohort, case-control, case-crossover, and time-series studies that evaluated the associations between environmental risk factors and health outcomes defined as incidence, prevalence, and mortality. The specific search strategy was designed in PubMed using free text and Medical Subject Headings (MeSH) terms related to risk factors, environment, health outcomes, observational studies, and meta-analysis. The search was limited to English, Spanish, and French published articles and studies on humans. The search was conducted on September 20, 2020. Risk factors were defined as any attribute, characteristic, or exposure of an individual that increases the likelihood of developing a disease or death. The environment was defined as the external elements and conditions that surround, influence, and affect a human organism or population's life and development. The environment definition included the physical environment such as nature, built environment, or pollution, but not the social environment. We excluded occupational exposures, microorganisms, water, sanitation and hygiene (WASH), behavioral risk factors, and no- natural disasters.

Results: This umbrella review found 197 associations among 69 environmental exposures and 83 diseases and death causes reported in 103 publications. The environmental factors found in this review were air pollution, environmental tobacco smoke, heavy metals, chemicals, ambient temperature, noise, radiation, and urban residential surroundings. Among these, we identified 65 environmental exposures defined as risk factors and 4 environmental protective factors. In terms of study design, 57 included cohort and/or case-control studies, and 46 included time-series and/or case-crossover studies. In terms of the study population, 21 included children, and the rest included adult population and both sexes. In this review, the largest body of evidence was found in air pollution (91 associations among 14 air pollution definitions and 34 diseases and mortality diagnoses), followed by environmental tobacco smoke with 24 associations. Chemicals (including pesticides) were the third larger group of environmental exposures found among the meta-analyses included, with 19 associations.

Conclusion: Environmental exposures are an important health determinant. This review provides an overview of an evolving research area and should be used as a complementary tool to understand the connections between the environment and human health. The evidence presented by this review should help to design public health interventions and the implementation of health in all policies approach aiming to improve population health.

Excerpts

Table 13: <https://www.mdpi.com/1660-4601/18/2/704/htm>

Radon, a radioactive natural, was found in a recent meta-analysis as a risk factor for lung cancer [100] at indoor radon exposure levels above 100 Bq/m³ (Table 13). In another meta-analysis, indoor radon exposure was also associated as a risk factor for childhood leukemia [101]. Finally, long-term exposures to extremely low-frequency electromagnetic fields were also found associated as a risk factor for amyotrophic lateral sclerosis [67] and childhood leukemia [99] (Table 13). Extremely low-frequency (ELF) magnetic fields are alternating fields generated by the distribution and supply of electricity.

Although this umbrella review found several publications and associations among environmental exposures and health outcomes, we also identified several evidence gaps. Most of the studies focus on identifying environmental risk factors, and only a few studies have been focusing on identifying environmental protective factors. Furthermore, few studies have focused on vulnerable and disadvantaged populations (children, elders, social disadvantaged, ethnic minorities, etc.). Furthermore, most studies do not provide a clear definition of the health outcomes using the international classification of diseases (ICD), nor a comparable exposure definition when the same pollutant is used. In terms of the meta-analysis, we exclude several studies from this review because, in the analyses, cross-sectional studies were mixed with other observational studies (i.e., cohorts).

Additionally, several studies did not report heterogeneity values (i.e., I²) or do not provide dose-response functions essential for population risk assessment, health impact assessments and policy translation. We have summarized a list of recommendations for future research in environmental health studies based on these gaps, and we have listed those recommendations in Table 16. Open access paper: <https://www.mdpi.com/1660-4601/18/2/704/htm>

REPEATED ELECTROMAGNETIC FIELD STIMULATION LOWERS AMYLOID-B PEPTIDE LEVELS IN PRIMARY HUMAN MIXED BRAIN TISSUE CULTURES

Felipe P Perez, Bryan Maloney, Nipun Chopra, Jorge J Morisaki, Debomoy K Lahiri. *ci Rep.* 2021 Jan 12;11(1):621. doi: 10.1038/s41598-020-77808-2.

Abstract

Late Onset Alzheimer's Disease is the most common cause of dementia, characterized by extracellular deposition of plaques primarily of amyloid- β (A β) peptide and tangles primarily of hyperphosphorylated tau protein. We present data to suggest a noninvasive strategy to decrease potentially toxic A β levels, using repeated electromagnetic field stimulation (REMFS) in primary human brain (PHB) cultures. We examined effects of REMFS on A β levels (A β 40 and A β 42, that are 40 or 42 amino acid residues in length, respectively) in PHB cultures at different frequencies, powers, and specific absorption rates (SAR). PHB cultures at day in vitro 7 (DIV7) treated with 64 MHz, and 1 hour daily for 14 days (DIV 21) had significantly reduced levels of secreted A β 40 ($p = 0.001$) and A β 42 ($p = 0.029$) peptides, compared to untreated cultures. PHB cultures (DIV7) treated at 64 MHz, for 1 or 2 hour during 14 days also produced significantly lower A β levels. PHB cultures (DIV28) treated with 64 MHz 1 hour/day during 4 or 8 days produced a similar significant reduction in A β 40 levels. 0.4 W/kg was the minimum SAR required to produce a biological effect. Exposure did not result in cellular toxicity nor significant changes in secreted A β precursor protein- α (sAPP α) levels, suggesting the decrease in A β did not likely result from redirection toward the α -secretase pathway. EMF frequency and power used in our work is utilized in human magnetic resonance imaging (MRI), thus suggesting REMFS can be further developed in clinical settings to modulate A β deposition.

<https://pubmed.ncbi.nlm.nih.gov/33436686/>