

Proposed Bioelectric Dissonance Pathway to Metabolic Dysfunction

Non-native EMF exposure → S4/CYB5B/ER-mitochondria timing disruption → insulin pulse distortion, impaired glucose uptake, excess hepatic glucose output, and metabolic-syndrome drift

Proposed mechanistic integration


A biologically grounded synthesis, not settled proof.

1. Non-native EMF / ELF Exposure




Waveform / phase structure may bias voltage-sensitive gating

2. Metabolic decision hardware




3. Metabolic desynchronization

Timing noise turns coordinated fuel handling into metabolic mis-coding




4. Tissue-specific consequences



Secondary inset (human):
Acute local brain glucose-metabolism changes have been reported in human exposure studies.

Findings vary by region, task, device, and exposure parameters; not all studies show effects.

5. Metabolic dysfunction phenotype



LEGEND: Arrow code

- Solid = established biology
- - - → Dashed = proposed bridge
- → Dotted = downstream interpretation

PATHWAY SUMMARY

Non-native EMF → S4/CYB5B/MAM timing disruption → Ca^{2+} / ATP / redox mis-coding → insulin-pulse distortion + impaired glucose uptake + excess hepatic glucose output + adipose spillover → systemic metabolic dysfunction

SELECTED EVIDENCE ANCHORS (Complementary lines of support)



Human PET:
acute local brain glucose-metabolism shift reported



Rat islets:
hyperglycemia, oxidative stress, impaired insulin secretion reported



Established physiology:
 β -cell Ca^{2+} pulsatility, liver Ca^{2+} control, muscle Ca^{2+} -glucose coupling, MAMs in insulin resistance



Results depend on exposure context, biology, intensity, and timing.