Pathophysiology 16 (2009) 113–122

Epidemiological evidence for an association between use of wireless phones and tumor diseases

ELSEVIER

PATHOPHYSIOLOGY

www.elsevier.com/locate/pathophys

Lennart Hardell [a,*], Michael Carlberg [a], Kjell Hansson Mild [b]

[a] *Department of Oncology, University Hospital, SE-701 85 Örebro, Sweden*
[b] *Department of Radiation Physics, Umeå University, SE-901 87 Umeå, Sweden*

Received 7 November 2008; accepted 30 January 2009

Abstract

During recent years there has been increasing public concern on potential cancer risks from microwave emissions from wireless phones. We evaluated the scientific evidence for long-term mobile phone use and the association with certain tumors in case–control studies, mostly from the Hardell group in Sweden and the Interphone study group. Regarding brain tumors the meta-analysis yielded for glioma odds ratio (OR) = 1.0, 95% confidence interval (CI) = 0.9–1.1. OR increased to 1.3, 95% CI = 1.1–1.6 with 10 year latency period, with highest risk for ipsilateral exposure (same side as the tumor localisation), OR = 1.9, 95% CI = 1.4–2.4, lower for contralateral exposure (opposite side) OR = 1.2, 95% CI = 0.9–1.7. Regarding acoustic neuroma OR = 1.0, 95% CI = 0.8–1.1 was calculated increasing to OR = 1.3, 95% CI = 0.97–1.9 with 10 year latency period. For ipsilateral exposure OR = 1.6, 95% CI = 1.1–2.4, and for contralateral exposure OR = 1.2, 95% CI = 0.8–1.9 were found. Regarding meningioma no consistent pattern of an increased risk was found. Concerning age, highest risk was found in the age group <20 years at time of first use of wireless phones in the studies from the Hardell group. For salivary gland tumors, non-Hodgkin lymphoma and testicular cancer no consistent pattern of an association with use of wireless phones was found. One study on uveal melanoma yielded for probable/certain mobile phone use OR = 4.2, 95% CI = 1.2–14.5. One study on intratemporal facial nerve tumor was not possible to evaluate due to methodological shortcomings. In summary our review yielded a consistent pattern of an increased risk for glioma and acoustic neuroma after >10 year mobile phone use. We conclude that current standard for exposure to microwaves during mobile phone use is not safe for long-term exposure and needs to be revised.

Keywords: Brain tumors; Glioma; Acoustic neuroma; Meningioma; Cellular phones; Cordless phones