

COMMENT

Open Access



Scientific evidence invalidates health assumptions underlying the FCC and ICNIRP exposure limit determinations for radiofrequency radiation: implications for 5G

International Commission on the Biological Effects of Electromagnetic Fields (ICBE-EMF)*

Abstract

In the late-1990s, the FCC and ICNIRP adopted radiofrequency radiation (RFR) exposure limits to protect the public and workers from adverse effects of RFR. These limits were based on results from behavioral studies conducted in the 1980s involving 40–60-minute exposures in 5 monkeys and 8 rats, and then applying arbitrary safety factors to an apparent threshold specific absorption rate (SAR) of 4W/kg. The limits were also based on two major assumptions: any biological effects were due to excessive tissue heating and no effects would occur below the putative threshold SAR, as well as twelve assumptions that were not specified by either the FCC or ICNIRP. In this paper, we show how the past 25 years of extensive research on RFR demonstrates that the assumptions underlying the FCC's and ICNIRP's exposure limits are invalid and continue to present a public health harm. Adverse effects observed at exposures below the assumed threshold SAR include non-thermal induction of reactive oxygen species, DNA damage, cardiomyopathy, carcinogenicity, sperm damage, and neurological effects, including electromagnetic hypersensitivity. Also, multiple human studies have found statistically significant associations between RFR exposure and increased brain and thyroid cancer risk. Yet, in 2020, and in light of the body of evidence reviewed in this article, the FCC and ICNIRP reaffirmed the same limits that were established in the 1990s. Consequently, these exposure limits, which are based on false suppositions, do not adequately protect workers, children, hypersensitive individuals, and the general population from short-term or long-term RFR exposures. Thus, urgently needed are health protective exposure limits for humans and the environment. These limits must be based on scientific evidence rather than on erroneous assumptions, especially given the increasing worldwide exposures of people and the environment to RFR, including novel forms of radiation from 5G telecommunications for which there are no adequate health effects studies.

Keywords: Federal Communications Commission (FCC), International commission on non-ionizing radiation protection (ICNIRP), Radiofrequency radiation (RFR), Exposure limits, Exposure assessment, Radiation health effects, Reactive oxygen species (ROS), DNA damage, 5G, Scientific integrity, Cell phone*, Mobile phone*

Introduction

In establishing exposure limits for toxic or carcinogenic agents, regulatory agencies generally set standards that take into account uncertainties of health risks for the general population [1] and for susceptible subgroups such as children [2]. That approach has not been applied in the same way to the setting of exposure limits for

*The terms cell phone and mobile phone are used interchangeably in this commentary; cell phone is the term used in the United States, while mobile phone is the term used in most of Europe.

*Correspondence: ron.melnick@gmail.comTucson, USA

¹Tucson, USA



radiofrequency radiation (RFR) (frequency range: 3 kHz to 300 GHz). Moreover, assumptions underlying the current RFR exposure limits are flawed; hence, the limits that are currently applied do not adequately protect human and environmental health. This issue is discussed in greater detail under Assumption #9.

The Federal Communications Commission's (FCC) limits for maximum permissible exposure to RF electromagnetic fields (EMF) [3] were established in 1996 [4], and currently include many recommendations from the International Commission on Non-Ionizing Radiation Protection [5]. These exposure limits were expected to protect against adverse health effects in humans that might occur from short-term (i.e., acute) exposures to RFR and have been maintained by the FCC for the past 26 years. The exposure limits that were established by the FCC in 1996 relied on criteria recommended by the National Council on Radiation Protection & Measurements (NCRP) [6] and the Institute of Electrical and Electronics Engineers (ANSI/IEEE) [7, 8]. The limits were "based on a determination that potentially harmful biological effects can occur at a SAR (specific absorption rate) level of 4.0 W/kg as averaged over the whole-body." The SAR is a measure of the rate of RF energy absorbed per unit mass.

The threshold for a behavioral response and for acute thermal damage in sensitive tissues was considered to be an exposure that produced a whole-body SAR greater than 4 W/kg. In parallel with the development of the FCC's RFR exposure limits, ICNIRP's guidelines for limiting exposure to RF-EMF were also based on behavioral studies conducted in rats and monkeys in the 1980s [9].

The harmful effects that served as the basis for the exposure criteria were changes in behavior observed in small numbers of rats and monkeys when exposed to RFR for up to 60 minutes to power densities at which the whole-body SAR was approximately 4 W/kg or higher [10, 11]. Those studies were conducted in the early 1980s (1980 and 1984, respectively) by investigators of the US Navy Department. Consequently, 4 W/kg was identified as the threshold SAR for adverse health effects induced by RFR. In food-deprived monkeys that were exposed to three different frequencies (225 MHz, 1.3 GHz, and 5.8 GHz) during 60-min sessions, lever-pressing response rates for the delivery of food pellets were reduced compared to sham exposure sessions. The threshold SAR for this decreased response was reported to range from 3.2 to 8.4 W/kg [11]. Similarly, in food-deprived rats exposed to 40-min sessions at 1.28 or 5.62 GHz radiation, the threshold SAR for a decrease in response rate was reported to range from approximately 3.8 to 4.9 W/kg [10]. In experimental studies in which monkeys were exposed in an anechoic chamber for 4 hours to 1.29 GHz

radiation at various power densities, an increase in mean body temperature of 0.7°C was associated with a whole-body SAR of 4 W/kg [12]. Behavior disruption associated with an increase in body temperature of approximately 1.0°C was assumed to be the most sensitive measure of harmful effects from RF-EMF exposure.

After establishing 4 W/kg as the threshold dose for acute harmful effects, both the FCC [3, 4] and ICNIRP [5, 9] set exposure limits for controlled occupational exposures to 0.4 W/kg SAR averaged over the whole body (based on applying a 10-fold safety/uncertainty factor). For the general population, the FCC's and ICNIRP's exposure limits were set at 0.08 W/kg SAR averaged over the whole body (by applying an additional 5-fold safety/uncertainty factor) for frequencies between 3 MHz and 3 GHz. The exposure limits established by the FCC and ICNIRP do not account for any impact of differing signal characteristics, such as carrier wave modulations or pulsing of the signal. Whole-body exposures for the general population are based simply on power levels averaged over 30-minute periods [3, 5].

Based on SAR distributions from whole-body exposures in which local (i.e., partial body) SARs were estimated to be 10 to 20 times the average value, local exposure limits were set 20 times higher than the average whole-body exposure limit [4–7]. For occupational exposures, local peak exposure limits were permitted up to 8 W/kg averaged over any 1-g cube of tissue [4] or 10 W/kg averaged over any 10 g of contiguous tissue [9] by the FCC and ICNIRP, respectively. For the general population, local peak SARs for partial-body exposures were not to exceed 1.6 W/kg averaged over any 1 g of cube-shaped tissue [3], or not to exceed 2.0 W/kg averaged over any 10 g of cube-shaped tissue [5]. Higher limit values are permissible for extremities. Extremities include the hands, wrists, feet, ankles, and pinnae (the external part of the ear), despite the close proximity of the ear to the brain. These adjustments were made long before the widespread use of wireless communication devices in which the emitting antenna is typically held close to local body organs such as the brain. The NCRP document [6] acknowledges that exposures could be greater than the recommended safety limit values when people are in close proximity to emitters of RFR.

The setting of exposure limits for the prevention of excessive tissue heating was based on the following assumptions: 1) electromagnetic waves at frequencies used in wireless communications do not have sufficient energy to break chemical bonds or ionize molecules [13]; 2) RFR could not damage DNA; and 3) tissue heating was the only possible biological effect of nonionizing radiation [5, 9, 14–16]. For potential environmental and human health issues that are not addressed in the

setting of exposure limits (for example effects of chronic exposures, or effects of co-exposure of skin to RFR and other environmental agents, such as would occur with 5G exposure in combination with sunlight), the implicit assumption is that such effects do not matter, or that the arbitrarily selected safety/uncertainty factor is sufficient to deal with those concerns. In any case, it is expected that underlying assumptions applied to health risk assessments would be clearly described [1].

Exposure limits for RF radiation are based on numerous assumptions; however, research studies published over the past 25 years show that most of those assumptions are not supported by scientific evidence. In the NCRP report [6], the authors noted that when further understanding of biological effects of RF radiation becomes available, exposure guidelines will need to be evaluated and possibly revised. The ANSI/IEEE document [7] also notes that effects of chronic exposure or evidence of non-thermal interactions could result in revising exposure standards. Unfortunately, these recommendations were never implemented. Assumptions of

safety from exposures that could adversely affect human or environmental health should be tested and validated *before* widespread exposures occur, not afterwards, by agencies responsible for protecting public health.

In this paper, we highlight studies that demonstrate the fallacy of inherent assumptions in the FCC/ICNIRP guidelines for RF radiation exposure limits, and we find that the limits fail to protect human and environmental health. Fourteen assumptions that underlie the RFR exposure limits established in the 1990s and reaffirmed in 2020 by the FCC [4, 5] and ICNIRP [5, 9] are addressed in this paper and are shown in Fig. 1.

Assumptions underlying exposure limits for RF radiation and the scientific evidence demonstrating that these assumptions are not valid

A. Effects of RF radiation at exposures below the putative threshold SAR of 4W/kg

Assumption 1) *There is a threshold exposure for any adverse health effect caused by RF radiation; in the*

frequency range of 100 kHz to 6 GHz it is a whole-body exposure that exceeds an SAR of 4 W/kg. Any biological effect of RF radiation above the threshold exposure is due to tissue heating.

Cardiomyopathy and carcinogenicity

In response to a request from the Food and Drug Administration's (FDA) Center for Devices and Radiological Health [17], the National Toxicology Program (NTP) conducted toxicity and carcinogenicity studies of cell phone (CDMA- or GSM-modulated) radiation in rats and mice exposed to RFR at frequencies of 900 MHz and 1800 MHz, respectively [18, 19]. Exposures to RFR for up to 2 years occurred in reverberation chambers over 18 hours/day on a continuous cycle of 10 minutes on and 10 minutes off. In rats, the whole-body SAR levels during the 10-minute on cycles were 0, 1.5, 3, or 6 W/kg.

The major histopathological findings from the NTP study in male rats [18] included dose-related increases in cardiomyopathy, increased incidence of cancers and preneoplastic lesions in the heart (schwannoma and Schwann cell hyperplasia) and brain (glioma and glial cell hyperplasia), increases in prostate gland tumors and hyperplasias, significant increases in adrenal gland tumors, and significant increases in the overall incidence of benign or malignant neoplasms in all organs in the 3 W/kg groups. The incidence of cardiomyopathy was also increased in GSM-exposed female rats, and significant increases in DNA damage were found in rats and mice [18, 19]. Similarly, an earlier study by Chou et al. [20] found a significant (3.6-fold) increase in the incidence of primary malignant neoplasms in male rats exposed to 2450 MHz pulsed RFR for 25 months (21.5 hr./day) at an SAR that ranged from 0.15 to 0.4 W/kg.

A 3-day external peer-review of the NTP studies confirmed there was "clear evidence of carcinogenic activity" in male rats for heart schwannomas, and "some evidence of carcinogenic activity" for brain gliomas and adrenal gland tumors with exposure to either GSM- or CDMA-modulated RF radiation [21]. In addition, a lifetime study by the Ramazzini Institute reported a significant increase in heart schwannomas in male rats exposed 19 hour/day to 1800 MHz GSM-modulated RFR at a field strength of 50 V/m, equivalent to a whole-body SAR of 0.1 W/kg [22]. The incidence of heart Schwann cell hyperplasia was also increased in that exposure group. These findings are consistent with results from the NTP study and demonstrate that the proliferative effect of modulated RFR in heart Schwann cells is a reproducible finding that can occur at doses far below the assumed whole-body threshold SAR of 4 W/kg.

ICNIRP [23] dismissed the evidence of carcinogenicity for RFR that was provided in the studies by the NTP [18] and the Ramazzini Institute [22] based on their earlier critique of those studies [24]. However, that critique demonstrated an unfortunate lack of understanding together with a misrepresentation of the design, conduct, and interpretation of experimental carcinogenicity studies in animal models [25], as well as a lack of appreciation for the remarkable concordance between the tumor responses observed in experimental animals with those identified in cancer epidemiology studies of mobile phone users described under Assumption #6.

Neither heating effects nor thermal stress was likely causal of the adverse health effects observed in the NTP [18] study, since there was no tissue damage observed in a 28-day study at the same SARs, there was no significant effect on body weight during the 2-year study, and there were no exposure-related clinical observations that would indicate thermal or metabolic stress. Furthermore, a preliminary thermal pilot study demonstrated that body temperatures did not increase by more than 1°C at the exposure levels used in the chronic studies [26], and there is no evidence that a small change in body temperature associated with the RFR exposures in the NTP study can cause the types of carcinogenic effects that were observed. The similar findings of GSM-modulated RFR on Schwann cells by the Ramazzini Institute [22] at much lower whole-body SARs confirm these effects to be independent of tissue heating.

Neurological effects

Though the FCC and ICNIRP exposure limits are based on a putative threshold dose of 4 W/kg due to behavioral disruption observed at higher doses in rats and monkeys [10, 11] numerous studies have shown consistent and reproducible deficits in spatial learning and memory in laboratory animals exposed to RF radiation at SARs below 4 W/kg. Examples of study exposures that demonstrated these neurological effects included 900 MHz GSM at 0.41–0.98 W/kg, 2 hr./day for 4 days in mice [27]; 900 MHz GSM at 0.52–1.08 W/kg, 2 hr./day for 1 month in rats [28]; 900 MHz GSM at 1.15 W/kg, 1 hr./day for 28 days in rats [29]; 900 MHz pulsed RFR at 0.3–0.9 W/kg for 6 hr./day in rats from conception to birth and tested at 30 days of age [30]; 900 MHz GSM and 1966 MHz UMTS at 0.4 W/kg for 6 months in rats [31]; and 900 MHz continuous wave EMF at 0.016 W/kg 3 hr./day for 28 days in rats [32]. The studies cited above are not the only studies showing these effects, but they clearly demonstrate that exposure to RFR at an SAR of 4 W/kg is not a threshold dose for neurological effects in rodents. The effects of RF radiation on spatial learning and memory indicate

the hippocampus as a target site of these exposures. For a more complete listing of neurological effects of RFR reported between 2007 and 2017 see Lai [33].

In addition, many studies have reported changes in brain electrical activities in human subjects, measured by electroencephalography (EEG), including sleep disturbance from single exposures to cell phone RF radiation. This is not surprising since the nervous system transmits messages based on electrical signals generated by nerve cells. Decreased β -trace protein, which is a key enzyme in the synthesis of a sleep-promoting neurohormone, has been seen in young adults with high-cumulative amounts of hours of mobile phone use [34]. Another frequently reported effect of RF radiation is increased blood-brain barrier permeability in rats at SARs much lower than 4 W/kg, e.g. [32, 35–41]. Oxidative stress induced in the brain of animals exposed to RF-EMF has been associated with observed neurological effects [42]. Although many studies did not observe significant changes in neurological effects in humans and several studies did not observe increased permeability in the blood-brain barrier in animal models [33], differences in EMF frequency, modulation, duration of exposure, and direction of incident waves to the exposed subject, as well as difference in dielectric properties and the size and shape of the exposed subject likely account for differences in observed effects [43, 44].

Sperm damage

The effect of non-ionizing microwave radiation on the testis (testicular degeneration in mice) was first reported 60 years ago [45]. Since then, and with the rapid increase in use of RF-EMF emitting devices, numerous studies have investigated testicular effects of RFR and potential associations with male infertility [46–50]. Human and animal studies have shown that the testis is one of the most sensitive organs to RF-EMF exposures, and that keeping a mobile phone in trouser pockets in talk mode can affect fertility parameters e.g., sperm motility, sperm count, sperm morphology, and apoptosis [48, 51]. Meta-analyses of published epidemiologic studies on the impact of mobile phone radiation on sperm quality in adult men have found significant decreases in sperm motility, sperm viability and/or sperm concentrations that were associated with mobile phone usage [52–55]. Several physical factors associated with exposure conditions can affect the outcome of human studies, including depth of energy penetration, duration of call, type of transmission technology, distance of the device to the body or testis, and power density with defined SAR. For example, Zilberlicht et al. [56] observed higher rates of

abnormal sperm concentrations among men who held their phones less than 50 cm from their groin.

The effects of RFR on reproductive parameters in humans are consistent with results from experimental studies in animals and in vitro studies. For example, exposure of human semen to 850 MHz radiation from mobile phones for 1 hour at an SAR of 1.46 W/kg caused a significant decrease in sperm viability that was associated with an increase in reactive oxygen species (ROS) [50] or an increase in sperm DNA fragmentation [57]. Exposure of isolated human spermatozoa to 1.8 GHz RF-EMF significantly reduced sperm motility and induced ROS generation at an SAR of 1.0 W/kg, and significantly increased oxidative DNA damage and DNA fragmentation at an SAR of 2.8 W/kg [58].

Some examples of effects of RFR on male fertility factors in studies with experimental animals at SARs below 4 W/kg include: a decrease in sperm count and an increase in ROS in rats exposed to mobile phone frequencies 2 hr./day, for 35 days (SAR=0.9 W/kg) [59]; increases in oxidative stress, 8-hydroxy-deoxyguanosine (8-OHdG), and DNA strand breaks in the testes of rats exposed to 900 MHz (SAR=0.166 W/kg), 1800 MHz (0.166 W/kg), or 2100 MHz (0.174 W/kg) 2 hr./day for 6 months [60]; an increase in ROS, a decrease in sperm count, and altered sperm morphology in rats exposed to 900 MHz 3G mobile phone radiation (SAR=0.26 W/kg) 2 hr./day for 45 days [61]; decreased sperm quality in rats in which local exposure of the scrotum to 2575–2635 MHz 4G smartphone time division LTE radiation occurred for 1 min over 10 min intervals 6 hr./day for 150 days [62]; impaired testicular development at 35 days of age in male offspring of pregnant rats that were exposed to 2.45 GHz RFR (SAR=1.75 W/kg) 2 hr./day throughout pregnancy [63]; decreased sperm motility in mice exposed to 905 MHz RFR (SAR=2.2 W/kg) 12 hr./day for 5 weeks, and increased ROS formation and DNA fragmentation after 1 week of exposure [64]. Although negative studies have also been reported, it is important to remember that the outcome of experimental studies can be affected by differences in exposure conditions, including the frequency, modulation, polarization, stray electromagnetic fields, local SAR, duration of exposure, and analytical methods [43, 44].

Although the mechanism of testicular effects from exposure to non-thermal levels of RFR is not fully known, numerous studies in rats and mice, and in human sperm have found associations between negative effects on fertility parameters and increases in ROS and/or DNA damage [48, 51, 57, 58, 60, 61, 64–68]. Thus, the adverse effects of RFR on sperm quality are likely due in large part to induced generation of ROS.

Assumption 2) *RF radiation is incapable of causing DNA damage other than by heating; there is no mechanism for non-thermal DNA damage.*

In 2009, ICNIRP [16] claimed that “low energy photons of RF radiation are too weak to affect ionization or cause significant damage to biological molecules such as DNA, under ordinary circumstances.” However, DNA damage and other genotoxic effects have been observed in numerous studies of low intensity RFR in animal models and in humans. For example, the NTP study found statistically significant increases in DNA damage in brain cells of exposed rats and mice compared to sham controls [18, 19, 69], and Akdag et al. [70] found statistically significant increases in DNA damage in hair cells in the ear canal among 30 to 60 year-old men who used mobile phones for 10 years for 0–30 min/day, 30–60 min/day, or greater than 60 min/day compared to people who did not use mobile phones. In the latter study, the extent of DNA damage increased with increasing daily exposure duration. In a review of published studies on genetic effects of ELF- and RF-EMF, Lai [71] listed more than 150 studies in which non-thermal exposures to RFR produced increases in DNA damage, chromosome aberrations, or micronuclei formation.

In addition, it is well established that DNA damage can also be caused by indirect processes, such as by the generation of reactive oxygen species (ROS), and numerous studies have demonstrated DNA damage at exposures below the putative threshold SAR of 4 W/kg. More than 120 published studies have demonstrated oxidative effects associated with exposure to low intensity RFR (Additional file 1: Appendix 1). An analysis of experimental studies on molecular effects of low intensity RF radiation (RFR) in biological systems found that the majority (93 of 100 studies) demonstrated the induction of oxidative effects [72]. More recent studies (from 2017) revealed that all 30 relevant publications (100%) detected significant oxidative effects under low intensity RFR exposures, and most of these studies used modulated RFR from wireless communication devices.

Increased production of ROS in living cells may be caused by weak magnetic fields altering recombination rates of short-lived radical pairs generated by normal metabolic processes leading to changes in free radical concentrations [73], or by low intensity extremely low frequency (ELF) EMFs resulting in alterations in voltage-gated ion channels in cell membranes causing changes in cation flow across membranes [74]. These mechanisms apply to both ELF-EMFs and to RFR modulated by pulsed fields at extremely low frequencies. Other biophysical mechanisms by which non-thermal RF-EMF can

cause biological effects through interactions with normal cellular processes have been described [75].

Increasing NADH oxidase activity is another mechanism by which RFR can increase ROS production. NADH oxidases, which are membrane-associated enzymes that catalyze one-electron reduction of oxygen to superoxide radical using NADH as the electron donor, have been identified as primary mediators of RFR interactions in cellular systems [76]. A significant (3-fold) increase in the activity of NADH oxidase was measured in purified plasma membranes from HeLa cells exposed to 875 MHz for 5 or 10 min at a power density of 200 $\mu\text{W}/\text{cm}^2$. This exposure intensity is significantly lower than the ICNIRP [5] safety limit.

The major source of ROS in living cells is the mitochondrial electron transport chain, where leakage of electrons generates superoxide radicals due to the partial reduction of oxygen [77]. A dose-dependent effect of 1.8 GHz modulated RFR exposure (SAR=0.15 and 1.5 W/kg) on mitochondrial ROS production was detected in mouse spermatogonial germ cells [65]. Exposure of quail embryos to extremely low intensity modulated RFR (GSM 900 or 1800 MHz, 0.25 or 0.32 $\mu\text{W}/\text{cm}^2$) during the initial days of embryogenesis resulted in a robust overproduction of superoxide radical and nitrogen oxide in mitochondria of embryonic cells [78, 79]. Thus, multiple mechanisms for the increased production of ROS by low intensity RF radiation have been demonstrated.

Numerous studies have been published on mutagenic effects of low intensity RF-EMFs, especially studies that identified increases in levels of a specific marker of oxidative DNA damage and a risk factor for cancer, 8-hydroxy-2'-deoxyguanosine (8-OHdG) [58, 60, 78–84]. For example, the level of 8-OHdG in human spermatozoa was increased significantly after in vitro exposure for 16 hr. to 1.8 GHz at a power level of 2.8 W/kg and correlated with levels of ROS generation [58]. Likewise, exposure of quail embryos *in ovo* to GSM-modulated 900 MHz of 0.25 $\mu\text{W}/\text{cm}^2$ for 1.5, 5, or 10 days was sufficient to produce a significant, two-threefold, increase in 8-OHdG levels in embryonic cells [79]. Umbilical cord blood and placenta tissue samples obtained after delivery from women who used mobile phones during pregnancy had significantly higher levels of oxidative stress parameters, including 8-OHdG and malondialdehyde, compared to cord blood and placental tissue from women who did not use mobile phones during pregnancy [85]. In addition, DNA damage, analyzed by the comet assay, was increased significantly in cord blood lymphocytes obtained from women who used mobile phones during pregnancy compared to cord blood lymphocytes obtained from women who did not use mobile phones.

As low intensity RF radiation does not have sufficient energy to ionize DNA molecules, and as increased production of ROS in living cells due to RF-EMF exposures has been reliably documented, an indirect effect of this type of radiation is the formation of oxidative damage to DNA. The most aggressive form of ROS that can cause oxidative DNA damage is the hydroxyl radical; this reactive oxygen species can be generated from superoxide radical and hydrogen peroxide [86], which may be produced in living cells exposed to low intensity RF radiation. Ultraviolet radiation (UVR, encompassing UVA, UVB, and UVC), which is classified by IARC as “carcinogenic to humans”, can also cause indirect DNA damage by generating ROS [87]. Thus, both RFR and UVR, which can similarly induce oxidative DNA damage, can increase cancer risk by a similar mechanism.

Increased production of ROS and depletion of antioxidant capacity in living cells exposed to low intensity RF radiation can result in oxidative DNA damage. Induction of oxidative stress, which is a key characteristic of many human carcinogens [88], including UVR and asbestos, can also lead to genotoxicity and carcinogenicity of non-ionizing RF radiation without causing direct DNA damage.

Assumption 3) *Two to seven exposures to RF radiation for up to 1 hour duration are sufficient to exclude adverse effects for any duration of exposure including chronic exposures.*

The behavioral studies in 8 male rats and 5 male monkeys that served as the basis for the exposure limits to RF radiation adopted by the FCC and ICNIRP involved 2 to 7 exposure sessions of 40-minute duration for rats [10] and 3 exposure sessions of 60-minute duration for monkeys at each power density [11]. Additional support for the threshold SAR of 4 W/kg in the frequency range of 100 kHz to 6 GHz came from behavioral studies conducted in rats and monkeys by D’Andrea et al. [89, 90]. However, D’Andrea et al. [91, 92] also reported that exposure of rats to continuous wave 2450 MHz RFR for 14 or 16 weeks caused significant differences in behavioral activity between sham-exposed rats and RFR-exposed rats at mean SARs of 0.7 W/kg and at 1.23 W/kg, indicating that 4 W/kg is not a threshold SAR with extended exposure durations. Since that time many studies have shown that responses to non-thermal RFR depend on both exposure intensity and exposure duration [93]. Importantly, the same response was observed with lower exposure intensity but prolonged exposure duration as at higher exposure intensity and shorter duration [94].

Recognizing that the exposure limits do not address potential health effects after long-term exposures to

RF radiation emitted from wireless devices that people are experiencing, the FDA [17] nominated RF radiation to the NTP for chronic toxicology and carcinogenicity studies out of concern that “existing exposure guidelines are based on protection from acute injury from thermal effects of RFR exposure, and may not be protective against any non-thermal effects of chronic exposures.” Adverse health effects noted in Assumption #1, including cardiomyopathy, carcinogenicity, sperm damage, and neurological effects, as well as the human epidemiology studies to be described in Assumption #6, occurred with much longer exposures to RF radiation than the exposure durations used in the acute studies in rats [10] and monkeys [11]. Consequently, the acute behavioral exposure studies that served as the basis for exposure limits to RF radiation established by the FCC and ICNIRP are inadequate to identify and characterize adverse effects of RF radiation after longer exposure durations. Neither the exposure limits established in the 1990s by the FCC [4] or by ICNIRP [9], nor those reaffirmed more recently by these groups [3, 5] address health risks associated with long-term exposure to RF radiation.

Assumption 4) *No additional effects would occur from RF radiation with co-exposure to other environmental agents.*

The current FCC/ICNIRP exposure limits do not take into consideration interactive effects of RF radiation with other environmental agents even though such effects have been documented. Interactions of RF radiation with other agents may result in antagonistic or synergistic effects, i.e., effects that are greater than the sum of each agent alone.

In the International Agency for Research on Cancer (IARC) evaluation of the carcinogenicity of RF-EMF [44], the expert working group noted that 4 of 6 co-carcinogenesis studies available at that time showed increased responses with exposure to RF-EMF. One of those studies reported co-carcinogenic effects of UMTS-modulated RF radiation at 4.8 W/m² in the liver and lung of mice that had been treated with the carcinogen ethylnitrosourea (ENU) in utero [95]; the incidence of liver and lung cancers were increased in mice exposed to ENU plus RF radiation compared to cage controls, sham controls and ENU alone. After the IARC evaluation, Lerchl et al. [96] replicated the experimental design of Tillmann et al. [95] by exposing mice to RF-EMF at whole-body SAR levels of 0 (sham), 0.04, 0.4, and 2 W/kg. Significant increases in lung adenomas and/or liver carcinomas were observed at all exposure levels. Lerchl et al. [96] concluded that their “findings are a very clear indication that tumor-promoting effects

of life-long RF-EMF exposure may occur at levels supposedly too low to cause thermal effects.” Thus, the reproducibility of the tumor-promoting effects of RFR at non-thermal exposure levels has been demonstrated.

Other examples of reported synergistic effects include the following study results. Synergistic effects on damage to human lymphocytes were observed with co-exposure to RFR (1.8 GHz RFR, SAR 3 W/kg) and 2 different mutagens, namely, mitomycin C or 4-nitroquinoline-1-oxide [97], or with co-exposure to ultraviolet (UVC) light [98]. A synergistic effect was found on DNA damage in human blood cells exposed to 2450 MHz radiation (5 mW/cm²) and then exposed to mitomycin C [99]. A potentiation effect on DNA damage was observed in cultured mammalian cells exposed to CDMA-modulated 835 MHz RF-EMF (SAR = 4 W/kg) and the clastogens cyclophosphamide or 4-nitroquinoline-1-oxide [100]. Gene expression was altered in neuronal and glial cells of rats pre-treated with lipopolysaccharide, a neuroinflammatory agent, and then exposed to 1800 MHz GSM modulated radiation (SAR = 3.22 W/kg) for 2 hr. [101]. In rats pre-treated with picrotoxin, a chemical that induces seizures, exposure to pulse-modulated 900 MHz GSM-modulated RF radiation of mobile phones increased regional changes in brain activity and c-Fos expression [102, 103].

Exposure limits based on exposure to only RF radiation will result in an underestimation of the true risk and inadequate protection of human health under conditions in which co-exposures to other toxic agents lead to synergistic adverse effects [104].

B. Factors affecting dosimetry

Assumption 5) *Health effects are dependent only on the time-averaged SAR value; carrier wave modulations, frequency, or pulsing do not matter except as they influence the SAR.*

The FCC's and ICNIRP's exposure limits to RFR are based on SARs for frequencies up to 6 GHz and on power densities for frequencies between 6 GHz and 300 GHz averaged over 6-minute or 30-minute intervals for local areas and whole-body exposures [3, 5]. However, time-averaged dosimetry does not capture the unique characteristics of modulated or pulsed RFR. For example, GSM modulation may involve as many as 8 voice channels with a duration of 0.577 msec for each channel. Thus, the exposure from GSM modulation can be 8-times higher during each time slot pulse compared to exposure to a continuous wave at equivalent time-averaged SARs. Also, as noted under assumption #14, repetitive pulses of data in bursts with short exposures to 5G can cause localized

temperature spikes in the skin [105]. The impact of pulsed radiation on biological activities at the molecular or cellular levels is not taken into consideration with time-averaged dosimetry.

Another issue not addressed by time-averaged dosimetry is the importance of low frequency modulations on biological systems. As discussed under assumption #2, increased production of ROS in living cells and DNA damage have been demonstrated with exposure to low frequency modulations of radiofrequency carrier waves [106]. Exposure limits based on time-averaged SAR dosimetry or power density, without consideration of the impact of amplitude or frequency modulations, do not adequately address potential health effects of real-world exposures to RFR. There is ample evidence that various effects of RFR exposure depend on carrier wave modulations, frequency, or pulsing [43, 107, 108]. In contrast to ICNIRP/FCC, the IARC monograph on RFR carcinogenicity noted that RFR effects may be influenced by such exposure characteristics as duration of exposure, carrier frequency, type of modulation, polarization, exposure intermittence, and background electromagnetic fields [44].

C. Human brain tumor risk

Assumption 6) *The multiple human studies that find associations between exposure to cell phone RF radiation and increases in brain tumor risk are flawed because of biases in the published case-control studies, and because brain cancer rates have remained steady since the time that use of wireless communication devices became widespread.*

Although claims have been made that “current limits for cell phones are acceptable for protecting the public health” because “even with frequent daily use by the vast majority of adults, we have not seen an increase in events like brain tumors” [109], the SEER (Surveillance, Epidemiology, and End Results Program) database shows an annual decrease of 0.3% for all brain tumors, but an increase of 0.3% per year for glioblastoma in the US between 2000 and 2018 (<https://seer.cancer.gov/explorer/>). Most concerning was that the annual increase for glioblastoma was 2.7% per year for people under 20 years of age. In addition, Zada et al. [110] reported that the incidence of glioblastoma multiforme (GBM) in the frontal lobe, temporal lobe, and cerebellum increased in the US between 1992 and 2006, and Philips et al. [111] likewise reported a statistically significant increasing incidence of GBM in the frontal and temporal lobes of the brain in the UK during 1995–2015. In Sweden, rates of brain tumors in the Swedish National Inpatient Register and the Swedish Cancer Register increased from 1998 to

2015 [112]. In addition, it should be realized that cumulative exposure, side-of-head use, and latency for tumor formation from RFR are not fully captured in national cancer registries. Thus, the claim that trends in brain cancer incidence rates have not increased since mobile phones were introduced is both wrong and misleading. The specificity of effect needs to be factored into such trend analyses.

Case-control studies, using sound scientific methods, have consistently found increased risks with long-term, heavy mobile phone use for brain tumors of the glioma type and acoustic neuroma. This association was evaluated at IARC in 2011 by 30 expert participants who concluded that radiofrequency (RF) radiation is a “possible” human carcinogen [44]. In contrast, the much-cited Danish cohort study on ‘mobile phone users’ [113] was disregarded by IARC due to serious methodological shortcomings in the study design, including exposure misclassifications [44, 114].

Results of meta-analyses of glioma risk and acoustic neuroma from Swedish case-control studies conducted by Hardell and coworkers [115, 116], the 13-nation Interphone study [117], and the French study by Coureau et al. [118] are shown in Table 1 as odds ratios (OR) with 95% confidence intervals. For glioma on any location in the head, a statistically significant increase of nearly two-fold was found, while for ipsilateral mobile phone use (tumor and phone use on the same side of the head) the risk was increased by 2.5-fold. These ORs are based on the groups in each study with the highest category of cumulative call time, which were ≥ 1640 hr. in the Interphone study [117, 119] and the Swedish studies [115, 116], and ≥ 896 hr. in the study by Coureau et al. [118]. Decreased survival among glioma cases, especially astrocytoma grade IV, was associated with long-term and high cumulative use of wireless phones [120]. Increased risk for the mutant

type of *p53* gene expression in the peripheral zone of astrocytoma grade IV was associated with use of mobile phones for ≥ 3 hours a day. Increase in this mutation was significantly correlated with shorter overall survival time [121].

For acoustic neuroma, risk was significantly increased with cumulative exposure and ipsilateral use by 2.7-fold. A random effects model, which was based on a test for heterogeneity, was used for the meta-analyses of these published studies. Tumor volume of acoustic neuroma increased per 100 hr. of cumulative use of wireless phones in the Swedish study and years of latency, indicating tumor promotion [115].

Other case-control studies of mobile phone use also reported increased risk of acoustic neuroma [122–124]. Those studies were not included in the meta-analysis because data on cumulative mobile phone use with numbers of cases and controls were not given or there were other shortcomings. It is also noteworthy that tumor risks were increased in subsets of the Interphone study; for example, there was nearly a 2-fold increase in the risk of acoustic neuroma for ≥ 10 y and ipsilateral use among the North European countries that participated in the Interphone study [125].

Claims have been made that associations between increases in brain cancer risk and exposure to cell phone RF radiation in the published case-control studies may be attributable to recall and/or selection biases [5, 109]. However, a re-analysis of the Canadian data that was included in the Interphone study showed that there was no effect on the risk of glioma after adjustments were made for selection and recall biases [126]. Odds ratios (OR) for glioma were increased significantly and to a similar extent when comparing the highest quartile of use to those who were not regular users whether or not adjustments for biases were made. In addition, Hardell

Table 1 Odds ratios (OR) with 95% confidence interval (CI) for glioma and acoustic neuroma in case-control studies in the highest category for cumulative mobile phone use in hours^a

	Glioma				Acoustic neuroma			
	All		Ipsilateral		All		Ipsilateral	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
Interphone [117, 119] Cumulative use ≥ 1640 hr	1.40	1.03–1.89	1.96	1.22–3.16	1.32	0.88–1.97	2.33	1.23–4.40
Coureau et al. [118] Cum use ≥ 896 hr	2.89	1.41–5.93	2.11	0.73–6.08				
Hardell et al. [115, 116] Cumulative use ≥ 1640 hr	2.13	1.61–2.82	3.11	2.18–4.44	2.40	1.39–4.16	3.18	1.65–6.12
Meta-analysis longest cumulative use	1.90	1.31–2.76	2.54	1.83–3.52	1.73	0.96–3.09	2.71	1.72–4.28

^a Note Hardell et al. [115, 116] also assessed use of cordless phones

and Carlberg [116] showed that the risk for glioma with mobile phone use was increased significantly even when compared to the risk for meningioma. Because risk of meningioma was not increased significantly, this tumor response could not be attributed to recall bias. Clearly, selection and recall biases do not explain the elevated brain tumor risk associated with the use of mobile phones. Thus, epidemiological evidence contradicts the opinions of the FCC and ICNIRP on brain tumor risk from RF radiation.

It should also be noted that the thyroid gland is a target organ for RFR from smartphones. A case-control study on mobile phone use suggested an increased risk for thyroid microcarcinoma associated with long-term cell phone use [127]. Peripheral lymphocyte DNA obtained from cases and controls was used to study genotype-environment interactions. The study showed that several genetic variants based on single nucleotide polymorphisms (SNPs) increased the risk of thyroid cancer with mobile phone use [128]. Increasing incidence of thyroid cancer in the Nordic countries, especially over the last two decades, has also been reported [129, 130]. In addition, a recent case-control study found significant increases in breast cancer risk among Taiwanese women based on their use of smartphones and distance between the breast and placement of their smartphone [131].

D. Individual variations in exposure and sensitivity to RF-EMF

Assumption 7) *There are no differences among individuals, including children, in the absorption of RF-EMF and susceptibility to this radiation.*

Differences between children and adults regarding the absorption of radiofrequency electromagnetic fields when mobile phones are operated close to the head have been demonstrated and widely documented [132–137]. The main factors accounting for these dissimilar absorption rates include differences in anatomy, tissue dielectric properties, and physiology. Through finite-difference time-domain (FDTD) simulations, employing detailed computational anthropomorphic models, it is possible to find differences relating to anatomy and to dimensions of the head.

Since EMF penetration into human tissues can be in the order of a few centimeters, depending on the wavelength, the inner tissues in the brain clearly will receive a significantly higher dose in the smaller heads of children compared to adults, despite the total absorption and the peak spatial SAR (psSAR) calculated across the whole head varying by smaller amounts [132, 133, 138]. Fernández et al. [136] estimated that the cell phone radiation psSAR in the hippocampus was 30-fold higher in

children compared to adults, while the psSAR in the eyes was 5-fold higher in children; these differences were due largely to closer proximity to the cell phone antennas. The thinner dimensions of children's skulls also contribute to this difference [135], resulting in a psSAR around 2-fold higher in children's brains [134–137, 139] compared to adults.

Additionally, tissues of young mammals have higher conductivity and electrical permittivity than those of mature animals [140]. This also contributes to greater EMF penetration and absorption, resulting in further increases in the psSAR. The psSAR in the skull bone marrow of children was estimated to increase by 10-fold due to higher conductivity in this tissue [137]. Distance between the mobile device and the body tissues is important in characterizing tissue dosimetry. The National Agency ANFR of France recently released cell phone SAR test data for 450 cell phones. Ten gram psSARs increased by 10–30% for each millimeter of proximal placement of the cell phone to the planar body phantom (<http://data.anfr.fr/explore/dataset/das-telephonie-mobile/?disjunctive.marque&disjunctive.modele&sort=marque>).

Finally, it is important to note that simulations of tissue dosimetry consider only the physical parameters of the tissues; they do not consider biological processes occurring in living tissues. While children are growing, developing organs and multi-organ systems are more susceptible to adverse effects of environmental agents; finite-difference time-domain (FDTD) simulations do not address differences in organ or system susceptibility for exposures occurring during child development.

Assumption 8) *There are no differences among individuals in their sensitivity to RF radiation-induced health effects.*

All life is “electrosensitive” to some degree as physiological processes are dependent on both subtle and substantial electromagnetic interactions at every level, from the molecular to the systemic. Responses to multiple types of electromagnetic exposure reveal that there is a far broader range of EMF sensitivity than previously assumed, and subgroups of extremely hypersensitive subjects exist [141–151]. Given the adverse health effects noted in Assumption #1, including cardiomyopathy, carcinogenicity and neurological effects, the acute, conscious symptoms manifesting in some individuals should not be unexpected. The term currently and most frequently used within the medical profession to describe those who are acutely, symptomatically sensitive to non-ionizing radiation exposures is Electromagnetic Hypersensitivity (EHS).

EHS is a multisystem, physical response characterized by awareness and/or symptoms triggered by EMF exposures. Common symptoms include (but are not limited to) headaches, dizziness, sleep disturbance, heart palpitations, tinnitus, skin rashes, visual disturbance, sensory disturbance, and mood disturbance [152, 153]. These symptoms are reported in response to even extremely low intensity (orders of magnitude below current safety levels) EMFs of multiple types (in terms of frequency, intensity and waveforms). Commonly noticed triggers of frequent and persistent EHS symptoms are pulse-modulated RF emissions, modulated at extremely low frequencies. Common triggering sources include mobile phones, DECT cordless landlines, Wi-Fi/Bluetooth-enabled computers, Wi-Fi routers, smart meters, base station antennas, and household electrical items. EMF avoidance/mitigation is found to be the most effective way to reduce symptoms [154].

Guidelines for EHS diagnosis and management have also been peer-reviewed and concur that the mainstay of medical management is avoidance of anthropogenic electromagnetic fields [152, 155, 156]. Case histories detailing clinical presentations, EMF measurements and mitigation are also published [157], and biomarkers including elevated markers of oxidative stress, inflammatory markers and changes in cerebral blood flow continue to be explored [152].

EHS has been proven to be a physical response under blinded conditions [145, 151, 158, 159] and, in addition to these studies, acute EMF-induced changes in cognition, behavior, and physiology reactions have been observed in studies involving animals [27, 30, 160–172]; plus further references under Assumption 13), which cannot be biased by media-cultivated fears. These studies provide further evidence which invalidates the nocebo response (physical symptoms induced by fear) as causal regarding symptoms.

It should not be expected that all provocation studies will reliably demonstrate adverse reactions; however, suggestions that the nocebo response may cause EHS symptoms were claimed from provocation studies which failed to show a relationship between the EMF exposure and the reported symptoms [173]. The failures of these studies are explainable given the very poor methodology in the majority of them. There were failures to account for a multitude of essential factors that must be tailored to the individual, such as variable symptom onset and offset, the necessity for adequate washout periods, specificity of trigger frequencies and intensities, requirement for complete EMF hygiene during sham exposures, requirement for life-like exposures (e.g., pulse-modulated information-carrying waves), etc. For example, it has been shown that various frequency channels from GSM/

UMTS mobile phones affect the same human cells differently [174–177]. Similarly, EHS has been shown to be frequency dependent [151]. As noted above, meaningful provocation studies need to take into consideration multiple physical parameters of exposure, including frequency, modulation, duration of exposure, and time after exposure [155]; however, most provocation studies that have failed to establish causative connection between RFR exposure and EHS symptoms [173] used only one or two conditions with short-term exposures.

There are many issues with the nocebo response as a cause of EHS, not least of which is also the absence of the required temporal link. For the nocebo response to be the cause of EHS, awareness and concern of negative health impacts from EMFs must precede symptoms. But, in the majority of EHS persons this is not the case [178]. As public risk communication improves, this will no longer be verifiable; however, this has been importantly observed at the only point in time when it could have been – prior to generalized awareness of health detriments from non-ionizing radiation (NIR).

While recognizing that some vulnerable groups may be more susceptible to effects of NIR exposure, ICNIRP [179] acknowledged that their guidelines may not safely accommodate these sensitive subgroups:

“Different groups in a population may have differences in their ability to tolerate a particular NIR [Non-Ionizing Radiation] exposure. For example, children, the elderly, and some chronically ill people might have a lower tolerance for one or more forms of NIR exposure than the rest of the population. Under such circumstances, it may be useful or necessary to develop separate guideline levels for different groups within the general population, but it may be more effective to adjust the guidelines for the general population to include such groups. Some guidelines may still not provide adequate protection for certain sensitive individuals nor for normal individuals exposed concomitantly to other agents, which may exacerbate the effect of the NIR exposure, an example being individuals with photosensitivity”.

In 2020, ICNIRP [23] also noted that biological effects are not easily discernible from adverse health effects, and that their guidelines:

“...are not intended to protect against biological effects as such (when compensatory mechanisms are overwhelmed or exhausted), unless there is also an associated adverse health effect. However, it is not always easy to draw a clear distinction between biological and adverse health effects, and indeed this can vary depending on individual susceptibility”.

to specific situations. An example is sensory effects from nonionizing radiation exposures under certain circumstances, such as a tingling sensation resulting from peripheral nerve stimulation by electric or magnetic fields; magnetophosphenes (light flickering sensations in the periphery of the visual field) resulting from stimulation of the retina by electric fields induced by exposure to low-frequency magnetic fields; and microwave hearing resulting from thermoelastic waves due to expansion of soft tissues in the head which travel via bone conduction to the inner ear. Such perceptions may sometimes lead to discomfort and annoyance. ICNIRP does not consider discomfort and annoyance to be adverse health effects by themselves, but, in some cases, annoyance may lead to adverse health effects by compromising well-being. The exposure circumstances under which discomfort and annoyance occur vary between individuals."

Trivializing "discomfort" which is the pre-cursor to pain is not in keeping with WHO recommendations quoted by the same ICNIRP [23] document: "Health is a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity."

Discomfort is a sign that an organism is experiencing something which is compromising optimal health and although in some cases this can be trivial and reversible, in other cases it may not be reversed. There is an extremely broad range of both pain tolerance and also of pain perception among humans, and to achieve meaningful preventative health care, "discomfort" must be taken seriously and mitigated whenever possible. This is especially true in this case where symptoms such as headaches are being reported in response to mobile phone exposures at the same time as increased brain tumor risk is noted from those same exposures (see Assumption 6).

In reality, people with EHS are reporting far more serious health disruption than "discomfort" or "annoyance" and in some cases these symptoms are disabling [180, 181]. Increasingly, EHS is being recognized as a disability by national courts in France, Sweden, and Spain, which amplifies the requirement for safety guidelines that are deliberately accommodating to this more susceptible group [180].

E. Applied safety factors for RF-EMF-RF workers and the general population

Assumption 9) *A 50-fold safety factor for whole body exposure to RF radiation is adequate for protecting the general population to any health risks from RF radiation.*

Public health agencies in the US and worldwide apply multiple uncertainty factors to health effects data to establish exposure levels that are considered safe for the great majority of exposed populations [182–184]. Although guidelines for the use of uncertainty factors were developed for chemicals, they are also pertinent to other toxic agents, such as RFR. The uncertainty factors needed for toxic effects of RFR based on studies that demonstrate a no-observed-adverse-effect level (NOAEL) in experimental animals include:

- 1) Animal-to-human extrapolation. When data are based on studies in experimental animals, a factor of 3–10 is applied (for potential species differences in tissue dosimetry and response) unless there are convincing data demonstrating equivalent sensitivity in animals and humans. However, there is no evidence showing that humans are equally or less sensitive to RFR than animals that were used in studies from which exposure limits were established by the FCC and ICNIRP.
- 2) Adjustment for human variability. A second factor of 10 is used to account for interindividual variability in susceptibility (for instance, due to differences in age, sex, genetic variation, pre-existing diseases) to the toxic agent among the general population. It has been recognized that a factor of 10 for human variability is likely inadequate for sensitive subpopulations and may require an additional adjustment.
- 3) Extrapolation from short-term studies to lifetime exposure. An additional factor of 10 is applied for short-term studies, such as those used to establish exposure limits to RF radiation, to provide lifetime protection from chronic exposure. This is of particular importance considering the remarkably short periods over which RFR toxicity was originally assessed [10, 11].
- 4) Database insufficiencies. Finally, an uncertainty factor of 3-to-10 is applied for database inadequacy, i.e., for incomplete characterization of an agent's toxicity. The behavioral studies [10, 11] that were used to establish the FCC and ICNIRP exposure limits to RFR do not provide a full characterization of the effects of this type of radiation nor did they identify the most sensitive adverse effect of RFR exposures.

Basing exposure limits to RFR on the behavioral studies in rats and monkeys [10, 11, 90, 91] would require the application of a composite uncertainty factor of about 900 to 10,000 to be consistent with approaches used by public health agencies to establish protective exposure limits for workers and the general population. Based on the size of the needed uncertainty/safety factor, the

data sets used by the FCC and ICNIRP are clearly inadequate to establish RF exposure limits with reasonable confidence. The arbitrarily selected safety factors of 10 for workers and 50 for the general population by the FCC and ICNIRP are woefully inadequate for protecting exposed populations.

When uncertainty/safety factors are applied to a misrepresented threshold exposure value for adverse effects, the resulting level does not provide assurance of health protection for the general population exposed to that agent. Studies cited above [18, 22, 91, 92, 96] show that the whole-body SAR of 4 W/kg is not a threshold level for adverse effects caused by RFR. In a recent quantitative analysis of various adverse health effects from the NTP study, Uche and Naidenko [185] showed that the permissible whole-body SAR of 0.08 W/kg (based on a 50-fold reduction of the assumed threshold SAR of 4 W/kg) was 20–40-fold higher than health protective SAR values derived by benchmark dose modelling of NTP data for cardiomyopathy (following application of 10-fold safety factors for interspecies and intraspecies variability). The approaches used by these authors are consistent with methodologies recommended by the US Environmental Protection Agency for quantifying health risks for toxic and carcinogenic environmental agents [1, 182]. Thus, a 50-fold reduction of the assumed threshold whole-body SAR of 4 W/kg is inadequate to protect the health of the general population from exposure to RF radiation.

Assumption 10) *A 10-fold safety factor for whole body exposure to RF radiation is adequate for protecting workers to any health risks from RF radiation.*

When RFR exposure limits were implemented in 1997, the rationale given for the difference in safety factors for the general population (50-fold) and for workers (10-fold) was “based on the exposure periods of the two populations, rounded to one digit (40 work hours per week/168 hours per week = ~0.2)” [6]. In addition to differences in exposure periods between workers and the general population, ICNIRP rationalizes the appropriateness of the lower safety factor for workers because “occupationally-exposed individuals can be considered a more homogeneous group than the general population,” they are, “in general, relatively healthy adults within a limited age range,” and “occupationally-exposed individuals should be operating under controlled conditions and be informed about the risks associated with non-ionizing radiation exposure for their specific situation and how to reduce these risks” [23]. In contrast, “the general public are, in most cases, unaware of their exposure to non-ionizing radiation and, without education, cannot

reasonably be expected to take precautions to minimize or avoid any adverse effects of exposure.”

The assumption that workers are trained in understanding health risks associated with exposure to RFR and in mitigating those risks to the greatest possible degree is not correct because neither the FCC nor the ICNIRP guidelines recognize any health effects from RFR at SARs below 4 W/kg, and the exposure limits authorized by the FCC and ICNIRP do not consider health effects from long-term exposures [3, 5]. The only health effect addressed by the FCC and ICNIRP is tissue damage due to excessive heating from acute exposures. Thus, the 10-fold reduction from the threshold whole-body SAR calculated from acute behavioral studies in rats and monkeys is inadequate for protecting the health of workers exposed long-term to RFR (see comments under assumption #9). There are no data demonstrating the adequacy of this arbitrarily chosen safety/uncertainty factor for occupationally-exposed workers, while on the contrary, excess cancer risks have been associated with exposure to RFR workers who operate radar and communication systems in military and occupational settings [186].

Assumption 11) *Exposure of any gram of cube-shaped tissue up to 1.6 W/kg, or 10g of cube-shaped tissue up to 2 W/kg, (duration not specified) will not increase the risk of that tissue to any toxic or carcinogenic effects in the general population.*

Tissue dosimetry was analysed in the NTP study of cell phone RF radiation in rats and mice [187]. In rats, whole body exposures during the 10-minute on cycles were 1.5, 3.0, or 6.0 W/kg, and the brain and heart SARs varied from the whole-body SARs by about 7% to under 2-fold for the brain and heart, respectively. A quantitative risk assessment of the NTP tumor incidence data is needed to evaluate organ-specific cancer risk. The FDA [19] nomination to the NTP recognized the need for “large well-planned animal experiments to provide the basis to assess the risk to human health of wireless communications devices.” However, more than 3 years after an external peer-review of the NTP studies found “clear evidence of carcinogenic activity,” the FDA [109] has continued to downplay the importance of these findings and avoid conducting a quantitative risk assessment of the tumor data that they (the FDA) originally requested. In contrast to the FDA, Uche and Naidenko [185] analysed the NTP data on cardiomyopathy by a benchmark dose approach and found that the 10% extra risk level for this effect was in the range of a whole-body SAR of 0.2 to 0.4 W/kg. Thus, there is an increased risk (greater than 10%) of developing cardiomyopathy at local tissue SARs below 1.6 or 2.0 W/kg.

The peak spatial specific absorption rate (psSAR), as used by ICNIRP and the FCC, is an inadequate dosimetric of RF radiation at frequencies above 1 GHz. The psSAR is calculated by averaging fixed cubic volumes containing a given amount of mass, and assumes a homogeneous material with a given mass density. The ICNIRP recommendation is to average cubic volumes containing 10 g of tissue (10 g-psSAR), while the FCC recommendation is to average cubic volumes containing 1 g of tissue (1 g-psSAR). Current recommendations limit the use of psSAR to frequencies up to 6 GHz [3, 5].

An evaluation of the utility of using psSAR as a dosimetric parameter at different frequencies ranging from 100 MHz to 26 GHz and with cube sizes ranging from 10 mg to 10 g is shown in Additional file 2: Appendix 2. For the smaller cubes and lower frequencies, averaging in the cube does not underestimate the maximum value on the cube surface, but at higher frequencies the psSAR averaged on larger cubes can be several-fold lower than the psSAR averaged on smaller cubes. For example, at 2.45 GHz, averaging over a 10-g cube underestimates by 4 dB (approximately 2.5-fold) the psSAR averaged in smaller cubes, while for 5.8 GHz, averaging over a 10-g cube underestimates the psSAR by 12 dB (approximately 16-fold) compared with averaging in a 10-mg cube, and by 6 dB (approximately 4-fold) compared with averaging over a 1-g cube. When the frequency is increased, the underestimation of the psSAR averaged in larger cubes (e.g. 10 g or 1 g) compared to smaller cubes (e.g. 100 mg and 10 mg) becomes more pronounced. Considering the 10-g cube, the difference between the psSAR for 5.8 GHz EMF compared to 0.9 GHz EMF is around 7 dB (or approximately 5-fold underestimation). These large differences are due to reduced penetration of EMFs at higher frequencies. Therefore, the ICNIRP's 10 g-psSAR and FCC's 1 g-psSAR recommendations do not provide reliable dosimetric parameters to evaluate EMF absorption above 1 GHz.

The SAR averaging over a 10-g cube is also flawed for assessing carcinogenicity because it is too large a volume to focus on stem cells and their important role in carcinogenesis. Human stem cells were more sensitive to RFR exposures from GSM and UMTS mobile phones than lymphocytes and fibroblasts [175]. Instead of a random distribution of targets for carcinogenesis, localized distribution of SAR in smaller volumes is needed to more accurately characterize relationships between SAR and tumor induction. From the point of view of stem cell organization, the volume of SAR determinations may be especially important for setting safety limits for children, because most stem cells and their niches are spatially and temporally transient during brain development [188].

Assumption 12) *Exposure of any gram of cube-shaped tissue up to 8 W/kg, or 10 g of cube-shaped tissue up to 10 W/kg, (duration not specified) will not increase the risk of that tissue to any toxic or carcinogenic effects in workers.*

Based on the analyses of tissue dosimetry in the NTP study [187], organ-specific toxic and carcinogenic effects were observed in rats at local tissue SARs that were much lower than 8 or 10 W/kg [18]. The tissue dosimetry in the NTP study and the inadequacy of the local SAR as specified by ICNIRP and the FCC is described in assumption #9.

F. Environmental exposure to RF radiation

Assumption 13) *There is no concern for environmental effects of RF radiation or for effects on wildlife or household pets.*

While background levels of RF-EMF are increasing in the environment, including rural remote areas [189], neither the FCC nor the ICNIRP take into consideration effects of this radiation on wildlife. The constant movement of most wildlife species in and out of varying artificial EMF can result in high exposures near communication structures, especially for flying species such as birds and insects. There is a substantial amount of scientific literature on the disrupting effects of RFR on wildlife (e.g., [190–206]).

Many nonhuman species use Earth's geomagnetic fields for activities such as orientation and seasonal migration, food finding, mating, nest and den building [190]. For example, migratory bird species [191, 192], honeybees [193], bats [194], fish [195–197], and numerous other species sense Earth's magnetic fields with specialized sensory receptors. Mechanisms likely involved in magneto-reception include magnetic induction of weak electric signals in specialized sensory receptors [198], magneto-mechanical interactions with the iron-based crystal magnetite [194], and/or free-radical interactions with cryptochrome photoreceptors [191, 192]. Each of these sensing processes shows extreme sensitivity to low intensity changes in electromagnetic fields. For a fuller description of the mechanisms by which non-human species use magneto-reception to perform essential life activities see Levitt et al. [190].

The following studies represent a few of the many examples of the disrupting effects of low-level exposures to RF-EMF on magneto-reception and the natural behavior of wildlife. Oscillating magnetic fields have been reported to disrupt the ability of migratory birds to orient and navigate in Earth's geomagnetic field [199–202].

Garden warblers became disoriented by exposure to a weak oscillating magnetic field of 1.403 MHz at an intensity as low as 2–3 nT [200]. The orientation of European robins that use Earth's magnetic field for compass orientation was completely disrupted by exposure to electromagnetic noise in the frequency range of 50 kHz to 5 MHz or a broadband noise-modulated ELF covering the range ~2 kHz to ~9 MHz [199, 201]. RFR in the low MHz range (7.0 MHz of 480 nT or 1.315 MHz of 15 nT) has been shown to disable the magneto-reception avian compass as long as the exposure was present [202].

In addition to effects on migratory birds, Landler et al. [203] found that exposure to a low-level magnetic field (1.43 MHz at an intensity of 30–52 nT) disrupted the natural orientation of juvenile turtles hatched on land. GSM-modulated 900 MHz RF radiation caused ants to lose their visual and olfactory memory for finding food [166]. Navigational abilities of trout were reduced when reared under conditions in which magnetic fields were spatially distorted [204].

Activities of honeybees are also disrupted by exposure to RF radiation. GSM-modulated cell phone radiation (900 MHz) caused a reduction in egg laying by queen bees and depletion of beehive pollen and honey counts [205]. GSM-modulated cell phone radiation (900 MHz) reduced hatching and altered pupal development of honey queen bee larvae [206].

The lack of consideration of chronic low-level RF radiation exposure on wildlife could result in dangerously disruptive effects on fragile ecosystems and on the behavior and survival of species that have long existed in Earth's natural environment.

G. 5G (5th generation wireless)

Assumption 14) *No health effects data are needed for exposures to 5G; safety is assumed because penetration is limited to the skin (“minimal body penetration”).*

Fifth generation (5G) wireless communication systems are being deployed worldwide to provide higher data transfer rates with shorter lag times between massive numbers of connected wireless devices. To provide faster transfer of large amounts of data (up to 20 gigabits per second peak data rates), the frequency range for 5G includes millimeter waves (30 to 300 GHz), in addition to carrier frequencies as low as 600 MHz. Extremely high frequency millimeter waves (MMW) that transmit large amounts of data to user devices are directed into narrow beams by line-of-sight transmission with beamforming antennas. Because millimeter waves do not penetrate solid structures such as building materials, hills, foliage, etc., and travel only short distances (a few hundred

meters), denser networks of base-stations with massive Multiple Input/Multiple Output (MIMO) transmitters and receivers in millions of small cell towers are being installed on structures such as utility poles. These features can lead to much closer proximity between humans and radiation-emitting antennas, and thereby change individual peak and average exposures to RFR.

For a 5G frequency of 26 GHz, EMF absorption is very superficial, which means that for typical human skin, more than 86% of the incident power is absorbed within the first millimeter. The skin penetration depth was computed as 1 mm based on the electrical conductivity of the skin and its electrical permittivity [5, 207]. This is expected to bring the SAR in this tissue well above the recommended limits ([208], and Additional file 2: Appendix 2). This is also expected to be harmful to very small species, such as birds and other small animals (e.g., insects) [209]. It is often claimed that because of its shallow penetration, exposure to high frequency 5G radiation is safe, and that the only effect is tissue heating [210]. However, this view ignores the deeper penetration of the ELF components of modulated RF signals, which are rated on the basis of heat alone, as well as the effects of short bursts of heat from pulsed signals [211, 212]. Within the first 1 mm of skin, cells divide to renew the stratum corneum (a consideration for skin cancer), and nerve endings in the dermis are situated within 0.6 mm (eyelids) to 3 mm (feet) of the surface (a consideration for neurological effects). Ultraviolet light, which exerts its action at a penetration depth of less than 0.1 mm [213, 214] is a recognized cause of skin cancer [87].

The higher the frequency of electromagnetic waves, the shorter the wavelength and the shallower the penetration of energy into exposed people or animals. For example, penetration depth in the human body is about 8 mm at 6 GHz and 0.92 mm at 30 GHz [5]. Because of the minimal depth of energy absorption at frequencies above 6 GHz, the FCC and ICNIRP have based exposure limits on power density instead of on SAR levels. The FCC [3] proposed a general localized power density exposure limit of 4 mW/cm² averaged over 1 cm² and not to exceed 30 minutes for 5G services up to 3000 GHz for the general population, claiming that this exposure is consistent with the peak spatial-average SAR of 1.6 W/kg averaged over any 1 g of tissue at 6 GHz. ICNIRP's [5] exposure limits for 5G are an absorbed power density of 200 W/m² (0.2 W/cm²) averaged over 4 cm² and a 6-minute interval for frequencies up to 30 GHz, and 400 W/m² (0.4 mW/cm²) averaged over 1 cm² and a 6-minute interval for frequencies of 30 GHz to 300 GHz.

Because of its minimal penetration, exposure to 5G radiation results in higher energy intensity on the skin and other directly-exposed body parts, such as the eye

cornea or lens. However, the skin, which is the largest organ in the human body, provides important functions such as acting as a protective physical and immunological barrier against mechanical injury, infection by pathogenic microorganisms, and entry of toxic substances. In addition, skin cancers, including basal cell carcinomas and squamous cell carcinomas, are the most prevalent human cancers, while melanomas are highly metastatic and increasing in prevalence. Although the high incidence of skin cancers are largely attributed to exposure to ultraviolet light, no studies have been reported on the effects of 5G radiation on (i) the skin's ability to provide protection from pathogenic microorganisms, (ii) the possible exacerbation of other skin diseases, (iii) promotion of sunlight-induced skin cancers, or (iv) initiation of skin cancer by itself. Information is also lacking on the effects of 5G radiation on nervous and immune systems which are also exposed even by the shallower penetration of MMW.

Another important factor is the maximum bandwidth with 5G radiation, which is up to 100 MHz in the frequency range of 450 MHz to 6 GHz, and up to 400 MHz in the ranges from 24 GHz to 52 GHz, compared to previous types of mobile communication where bandwidth is limited to 20 MHz. Because many studies indicated frequency-dependent, non-thermal RF effects from mobile communication RFR [43, 177] and for MMW effects [215, 216], the possibility of effective frequency windows for biological effects would increase with the increased bandwidth of 5G radiation.

Another consideration for effects of 5G exposures on human health is that radiation pulses created by extremely fast data transmission rates have the potential to generate bursts of energy that can travel much deeper than predicted by conventional models [217, 218]. Neufeld and Kuster [105] showed that repetitive pulses of data in bursts with short exposures to 5G can cause localized temperature spikes in the skin leading to permanent tissue damage even when the average power density values were within ICNIRP's acceptable safety limits. The authors urged the setting of new thermal safety standards to address the kind of health risks possible with 5G technology:

"The FIFTH generation of wireless communication technology (5G) promises to facilitate transmission at data rates up to a factor of 100 times higher than 4G. For that purpose, higher frequencies (including millimetre-wave bands), broadband modulation schemes, and thus faster signals with steeper rise and fall times will be employed, potentially in combination with pulsed operation for time domain multiple access...The thresholds for frequencies

above 10 MHz set in current exposure guidelines (ICNIRP 1998, IEEE 2005, 2010) are intended to limit tissue heating. However, short pulses can lead to important temperature oscillations, which may be further exacerbated at high frequencies (>10 GHz, fundamental to 5G), where the shallow penetration depth leads to intense surface heating and a steep, rapid rise in temperature..."

Areas of uncertainty and health concerns with 5G radiation include potential increase in skin cancer rates with (or possibly without) co-exposure to sunlight, exacerbation of skin diseases, greater susceptibility to pathogenic microorganisms, corneal damage or early development of cataracts, testicular effects, and possible resonant-enhanced absorption due to skin structures [219]. One of the complex technical challenges in relation to human exposure to 5G millimeter waves is that the unpredictable propagation patterns that could result in unacceptable levels of human exposure to electromagnetic radiation are not well understood [220]. Although MMW are almost completely absorbed within 1–2 mm in biologically-equivalent tissues, their effects may penetrate deeper in a live human body possibly by affecting signal transduction pathways. Thus, there are too many uncertainties with exposure to 5G to support an assumption of safety without adequate health effects data. There are no adequate studies on health effects from short-term or long-term exposures to 5G radiation in animal models or in humans.

Discussion

To develop health-based exposure limits for toxic and carcinogenic substances, regulatory agencies typically rely on available scientific evidence about the agent under review. In the mid- and late-1990s when the FCC [4] and the ICNIRP [9] initially established exposure limits for RFR, the prevailing assumptions were that any adverse effects from exposure to RFR were due to excessive heating because non-ionizing radiation did not have sufficient energy to break chemical bonds or damage DNA. However, non-thermal effects of RFR are demonstrated from studies that find different effects with exposure to continuous waves versus pulsed or modulated waves at the same frequency and the same SAR or power density, e.g., [221–226], and from studies that show adverse effects at very low exposure intensities, e.g., [78, 96].

Acute exposure studies conducted in rats and monkeys in the 1980s [10, 11] suggested that an SAR of 4 W/kg could be a threshold dose for behavioral effects. Because this SAR was associated with an approximate increase in body temperature of 1 °C, it was again assumed that no adverse health effects would occur if increases in core

body temperature were less than 1°C. From this putative threshold dose a “safety factor” of 10 was applied for occupational exposures and an additional factor of 5 (50x total) was applied for the general population, resulting in exposure limits in which the whole-body SAR was less than 0.4 W/kg for workers and 0.08 W/kg for the general population. However, realizing that local parts of the body could receive doses of RFR that were 10 to 20 times higher than the whole-body SARs, local peak exposure limits were set by the FCC at SARs 20-times higher than the whole-body SARs, i.e., 8 W/kg averaged over any 1-g of tissue for localized exposures for workers and 1.6 W/kg averaged over any 1-g for the general population [3, 4]. ICNIRP opted for partial body exposures that would not exceed 2.0 W/kg averaged over any 10g of cube-shaped tissue for the general population [5, 9]. To rationalize the smaller safety factor for workers (10-fold) versus the general population (50-fold), one claim made by ICNIRP [24] is that workers are informed about risks associated with non-ionizing radiation exposure and how to reduce these risks, whereas “the general public are, in most cases, unaware of their exposure to non-ionizing radiation and, without education, cannot reasonably be expected to take precautions to minimize or avoid any adverse effects of exposure.” From a public health perspective, the FCC and ICNIRP should make the public aware of their exposures to RFR and promote precautionary measures to minimize potential adverse effects, especially for children and pregnant women. Eight practical recommendations by the International EMF Scientist Appeal aimed at protecting and educating the public about potential adverse health effects from exposures to non-ionizing EMFs [227] are shown in Table 2.

The acute behavioral studies that provide the basis for the FCC’s and ICNIRP’s exposure limits lacked any information on potential effects of RF radiation that can occur after longer durations of exposure, and they did not address effects of carrier wave modulations used in wireless communications. Research on RFR conducted over

the past 25 years has produced thousands of scientific papers, with many demonstrating that acute behavioral studies are inadequate for developing health protective exposure limits for humans and wildlife, and that inherent assumptions underlying the FCC’s and ICNIRP’s exposure limits are not valid. First, 4 W/kg is not a threshold SAR for health effects caused by RFR exposures; experimental studies at lower doses and for longer durations of exposure demonstrated cardiomyopathy, carcinogenicity, DNA damage, neurological effects, increased permeability of the blood brain barrier, and sperm damage (see Assumptions 1–3). Multiple robust epidemiologic studies on cell phone radiation have found increased risks for brain tumors (Assumption 6), and these are supported by clear evidence of carcinogenicity of the same cell types (glial cell and Schwann cell) from animal studies. Even studies conducted by D’Andrea et al. [89, 90] before the limits were adopted found behavioral disruption in rats exposed to RFR for 14 or 16 weeks at mean SARs of 0.7 W/kg and at 1.23 W/kg. A combination of exposure duration and exposure intensity would be more appropriate for setting safety standards for exposure to RFR from mobile communication systems including mobile phones, base stations, and WiFi.

More than 120 studies have demonstrated oxidative effects associated with exposure to low intensity RFR (Additional file 1: Appendix 1). DNA damage that has been reported in studies of RFR was most likely caused by induction of oxidative stress, which is a key characteristic of human carcinogens [88], rather than by direct ionization (Assumption 2). The generation of reactive oxygen species has also been linked to DNA damage and the carcinogenicity of UVA radiation [87] and asbestos [228]. Despite the enormous amount of scientific evidence of low-dose effects of RFR, the IEEE [229] maintains that behavioral disruption is still the most sensitive and reproducible effect of RFR. It is this opinion that contributed to the FCC [3] and ICNIRP [5] reaffirming their previous exposure limits to RFR.

Table 2 Precautionary Measures Recommended by the International EMF Scientist Appeal

-
- 1) Priority should be given to protect children and pregnant women
 - 2) Guidelines and regulatory standards should be strengthened
 - 3) Manufacturers should be encouraged to develop safer technologies
 - 4) The public should be fully informed about the potential health risks from electromagnetic energy and taught harm reduction strategies
 - 5) Medical professionals need to be educated about the biological effects of electromagnetic energy and be provided training on treatment of patients with electromagnetic sensitivity
 - 6) Governments need to fund training and research on electromagnetic fields and health that is independent of industry
 - 7) The media should disclose experts’ financial relationships with industry when citing their opinions regarding health and safety aspects of EMF-emitting technologies
 - 8) Radiation-free areas need to be established, especially for individuals with EHS
-

Other concerns about the current exposure limits for RFR are that they do not consider potential synergistic effects due to co-exposure to other toxic or carcinogenic agents, the impact of pulsed radiation or frequency modulations, multiple frequencies, differences in levels of absorption or of susceptibility by children, or differences among individuals in their sensitivity to RFR (see Assumptions 4, 5, 7, 8). Currently, children's cumulative exposures are much higher than previous generations and they continue to increase [230]. ICNIRP [23, 179] acknowledged that their guidelines do not accommodate sensitive subgroups and admit to difficulties separating "biological effects" from "health effects." Neurological symptoms, some of which are acknowledged by ICNIRP and currently being experienced by persons with EHS, are most certainly non-thermal "health effects" that need to be mitigated by providing environments with reduced exposures to anthropogenic EMF for hypersensitive individuals.

The debilitating effects and restrictions suffered by adults and children with EHS constitutes a contravention of the 2010 Equalities Act, Human Rights Act and other ethical and legal frameworks. Failure to respond and appropriately safeguard this group is already causing preventable morbidity, mortality and economic deficit due to lost workdays, compensations for health damages and increased healthcare costs. Conversely, accommodating this group by, as suggested by ICNIRP [179], acting to 'adjust the guidelines for the general population to include such groups' would not only lessen the negative impacts for people with EHS, but would also improve public health more broadly, given the other NIR-related health concerns that are highlighted in this paper.

Basing local tissue exposure limits on 1-g [3] or 10-g [5] cubes substantially underestimates the peak spatial SAR compared to basing local tissue exposure limits on smaller cubes (e.g., 100 mg or 10 mg), and therefore are not reliable dosimetric parameters to evaluate EMF absorption at frequencies above 1 GHz (Assumptions 11, 12). The volumes specified by the FCC and ICNIRP for local tissue SAR limits are too large to focus on stem cells which are important targets for carcinogenesis. To reduce health risks from exposures to RFR, limits for localized distribution of the SAR should be based on 100 mg, or preferably 10 mg cubes.

Another important deficiency raised in this paper is that neither the FCC nor ICNIRP addresses concerns for environmental effects of RFR on wildlife, even though there is extensive literature demonstrating the disrupting effects of RFR on wildlife behavior (Assumption 13).

The arbitrarily selected uncertainty/safety factors applied to the putative threshold SAR for RFR are woefully inadequate for protecting public health

(Assumptions 9, 10). Based on the way the US Environmental Protection Agency, the International Council for Harmonization, and the National Institute for Occupational Safety and Health (US NIOSH) apply uncertainty/safety factors to a no-observed-adverse-effect level (NOAEL) in experimental animals [182–184], the safety factor for RFR would be at least 900 to 10,000, which is 18 to 200 times larger than the safety factor recommended by the FCC and ICNIRP for the general population. This large safety factor is based on adjustments for human variability, lifetime exposure from short-term studies, and database insufficiencies that include incomplete characterization of the toxicity of RFR. Clearly, the acute behavioral studies that served as the basis for the current exposure limits for RFR are not suitable for characterizing human health risks associated with long-term exposure to this type of radiation. The NCRP report from 1986 [6] and the ANSI/IEEE document from 1992 [7] recognized that when future studies on biological effects of RFR become available including effects of chronic exposures or evidence of non-thermal interactions there will be a need to evaluate and possibly revise exposure standards. When the FCC [3] and ICNIRP [5] reaffirmed their exposure limits from the 1990s, they dismissed the scientific evidence that invalidated the assumptions that underlie the basis for those exposure limits. An independent re-evaluation of RFR exposure limits based on the scientific knowledge gained over the past 25 years is needed and is long overdue. This evaluation should be performed by scientists and medical doctors who have no conflicting interests and who have expertise in RF-EMF exposure and dosimetry, toxicology, epidemiology, clinical assessment, and risk assessment. Special precautions should be taken to ensure that interpretations of health effects data and the setting of exposure limits for RFR are not influenced by the military or the telecommunications industry. In the meantime, manufacturers should be obliged to develop safer technologies [227].

Finally, we note our concern about the worldwide deployment of 5G communication networks for faster transfer of large amounts of data, but with no adequate health effects studies demonstrating the safety of high frequency millimeter waves. Because of limitations of the penetration and distance of travel of millimeter waves, dense networks of base stations are being mounted on structures such as utility poles in highly populated cities. Also, because the absorption of EMF at frequencies above 6 GHz is minimal, ICNIRP [5] has specified absorbed power density (S_{ab}) as the dosimetric parameter for "heating effects" at the higher frequencies. S_{ab} is a function of the incident power density (S_{inc}) and the input reflection coefficient (Γ). In near field scenarios, the S_{inc} does not have a singular value; this is largely due

to the heterogeneous nature of human body tissues and their relevant parameters (such as the permittivity, equivalent conductivity, mass density), which vary in different body regions and with frequency. Therefore, unless a powerful EMF simulation method together with realistic human models are used, the S_{inc} and the reflection coefficient values would be difficult to accurately estimate, making the resulting S_{ab} unreliable.

The assumption that 5G is safe at the power density limits recommended by ICNIRP (50 W/m² and 10 W/m² averaged over 6 min for occupational and 30 min for public exposures, respectively) because of its minimal penetration into the body does not justify the dismissal of the need for health effects studies prior to implementing 5G networks. The new communication networks will result in exposures to a form of radiation that has not been previously experienced by the public at large (Assumption 14). The implementation of 5G technology without adequate health effects information raises many questions, such as: Will exposure to 5G radiation: (i) compromise the skin's ability to provide protection from pathogenic microorganisms? (ii) will it exacerbate the development of skin diseases? (iii) will it increase the risk of sunlight-induced skin cancers? (iv) will it increase the risk of damage to the lens or cornea? (v) will it increase the risk of testicular damage? (vi) will it exert deeper tissue effects either indirectly following effects on superficial structures or more directly due to deeper penetration of the ELF components of modulated RF signals? (vii) will it adversely affect wildlife populations? Answers to these questions and others that are relevant to human and wildlife health should be provided *before* widespread exposures to 5G radiation occur, not afterwards. Based on lessons that should have been learned from studies on RFR at frequencies below 6 GHz, we should no longer rely on the untested assumption that current or future wireless technology, including 5G, is safe without adequate testing. To do otherwise is not in the best interest of either public or environmental health.

Abbreviations

ANSI: American National Standards Institute; CDMA: Code-division multiple access; dB: Decibel; DECT: Digital enhanced cordless technology; EHS: Electromagnetic hypersensitivity; ELF: Extremely low frequency; EMF: Electromagnetic field; FCC: Federal Communications Commission; FDA: Food and Drug Administration; GHz: Gigahertz; GBM: Glioblastoma multiforme brain cancer; GSM: Global system for mobile communication; IARC: International Agency for Research on Cancer; ICNIRP: International Commission on Non-Ionizing Radiation Protection; IEEE: Institute of Electrical and Electronics Engineers; LTE: Long Term Evolution (4G); MMW: Millimeter wave; NCRP: National Council on Radiation Protection and Measurements; NIR: Non-ionizing radiation; nT: Nanotesla; NTP: National Toxicology Program; 8-OHdG: 8-hydroxy-2'-deoxyguanosine; psSAR: Peak spatial specific absorption rate; RFR: Radiofrequency radiation; ROS: Reactive oxygen species; SAR: Specific absorption rate; UMTS: Universal mobile telecommunications service (3G); UVR: Ultraviolet radiation; 5G: 5th generation wireless.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12940-022-00900-9>.

Additional file 1: Appendix 1 Table 1. Studies demonstrating increased oxidative DNA damage and other indicators of oxidative stress at SAR < 4 W/kg.

Additional file 2: Appendix 2. On the inadequacy of the psSAR Dosimetric Parameter at Frequencies above 1 GHz. **Table 1.** Electric permittivity and electric conductivity of the gray matter. **Figure 1.** A block of gray matter radiated by different frequencies. The highlighted cubes are of 10 g, 1 g, 100 mg and 10 mg. **Fig. 2.** A block of gray matter radiated by different frequencies. The highlighted cubes are of 10 g, 1 g, 100 mg and 10 mg. **Fig. 3.** Electric field intensity averaged in each cube for different frequencies: in the left axis, the electric field is in dB and in the right axis the electric field is in V/m normalized to 100 V/m.

Acknowledgements

Igor Belyaev: Cancer Research Institute, Biomedical Research Center, Slovak Academy of Sciences, Slovakia
 Carl Blackman: US Environmental Protection Agency (retired), North Carolina, USA
 Kent Chamberlin: Department of Electrical and Computer Engineering, University of New Hampshire, USA
 Alvaro DeSalles: Graduate Program on Electrical Engineering (PPGEE), Federal University of Rio Grande do Sul (UFRGS), Porto Alegre, Brazil
 Suleyman Dasdag: Biophysics Department, Istanbul Medeniyet University, Medical School, Turkey
 Claudio Fernandez: Division of Electrical and Electronics Engineering, Federal Institute of Rio Grande do Sul (IFRS), Canoas, Brazil
 Lennart Hardell: Department of Oncology, Orebro University Hospital, Sweden (retired), The Environment and Cancer Research Foundation, Orebro, Sweden
 Paul Heroux: Epidemiology, Biostatistics and Occupational Health, Faculty of Medicine, McGill University, Canada
 Elizabeth Kelley: ICBE-EMF and International EMF Scientist Appeal, and Electromagnetic Safety Alliance, Arizona, USA
 Kavindra Kesari: Department of Applied Physics, School of Science, Aalto, University, Espoo, Finland
 Don Maisch: EMFacts Consultancy; The Oceanic Radiofrequency, Scientific Advisory Association; Tasmania, Australia
 Erica Mallory-Blythe: Physicians' Health Initiative for Radiation and Environment; British Society of Ecological Medicine; Oceania Radiofrequency Scientific Advisory Association, UK
 Ronald L. Melnick: National Toxicology Program, National Institute of Environmental Health Sciences (retired), Ron Melnick Consulting LLC, Logan, Utah, USA
 Anthony Miller: Dalla Lana School of Public Health (Professor Emeritus), University of Toronto, Ontario, Canada
 Joel M. Moskowitz: School of Public Health, University of California, Berkeley, California, USA
 Wenjun Sun: School of Public Health, Zhejiang University School of Medicine; Hangzhou, China
 Igor Yakymenko: National University of Food Technology, Kyiv Medical University, Ukraine

Authors' contributions

IB, AD, CF, LH, PH, KK, DM, EMB, RLM, and IY drafted the initial sections of this manuscript: by IB (factors affecting dosimetry), AD and CF (absorption in children versus adults, peak spatial specific absorption rate), LH (human brain cancer risk), KK (sperm damage), DM and DM (5G), EMB (electromagnetic hypersensitivity), RLM (cardiomyopathy, carcinogenicity, neurologic effects, safety factors), and IY (oxidative stress and DNA damage). IY prepared Appendix 1, and AD and CF prepared Appendix 2. The authors who drafted sections of the manuscript, as well as CB, KC, SD, EK, AM, JMM, and WS reviewed multiple manuscript drafts and made revisions. All authors reviewed and approved the final manuscript.

Funding

The Electromagnetic Safety Alliance provided funding for publication costs.

Availability of data and materials

All literature citations are available online.

Declarations**Ethics approval and consent to participate**

Not Applicable.

Consent for publication

Not Applicable.

Competing interests

IB, EMB, and AM have served as plaintiff's expert witnesses in cases involving radiofrequency radiation. All other authors declare they have no competing interests.

Received: 14 July 2022 Accepted: 8 September 2022

Published online: 18 October 2022

References

- US Environmental Protection Agency (US EPA). "Guidelines for carcinogen risk assessment", EPA/630/P-03/001F. Washington, DC; 2005. Available at https://www3.epa.gov/airtoxics/cancer_guidelines_final_3-25-05.pdf
- US Environmental Protection Agency (US EPA). "Supplemental guidance for assessing susceptibility for early-life exposure to carcinogens", EPA/630/R-03/003F. Washington, DC; 2005. Available at https://www.epa.gov/sites/production/files/2013-09/documents/childrens_supplement_final.pdf
- Federal Communications Commission (FCC). "Proposed Changes in the Commission's Rules Regarding Human Exposure to Radiofrequency Electromagnetic Fields; Reassessment of Federal Communications Commission Radiofrequency Exposure Limits and Policies", FCC19–126, 2019. <https://www.federalregister.gov/documents/2020/04/06/2020-06966/human-exposure-to-radiofrequency-electromagnetic-fields>
- Federal Communications Commission (FCC). "Evaluating Compliance with FCC Guidelines for Human Exposure to Radiofrequency Electromagnetic Fields", 1997. OET Bulletin 65. https://transition.fcc.gov/Bureaus/Engineering_Technology/Documents/bulletins/oet65/oet65.pdf
- International Commission on Non-Ionizing Radiation Protection (ICNIRP). Guidelines for limiting exposure to electromagnetic fields (100 kHz to 300 GHz). *Health Phys.* 2020;118:483–524.
- National Council on Radiation Protection and Measurements (NCRP). "Biological Effects and Exposure Criteria for Radiofrequency Electromagnetic Fields", NCRP Report No. 86, 1986. <https://ncrponline.org/publications/reports/ncrp-report-86/>
- American National Standards Institute (ANSI). "Safety Levels with Respect to Human Exposure to Radio Frequency Electromagnetic Fields, 3 kHz to 300 GHz," ANSI/IEEE C95.1–1992. <https://emfguide.itu.int/pdfs/c95.1-2005.pdf>
- D'Andrea JA, Adair ER, de Lorge JO. Behavioral and cognitive effects of microwave exposure. *Bioelectromagnetics Suppl.* 2003;6:S39–62.
- International Commission on Non-Ionizing Radiation Protection (ICNIRP). ICNIRP guidelines for limiting exposure to time-varying electric, magnetic and electromagnetic fields (up to 300 GHz). *Health Phys.* 1998;74:494–522.
- De Lorge JO, Ezell CS. Observing-responses of rats exposed to 1.28- and 5.62-GHz microwaves. *Bioelectromagnetics.* 1980;1:183–98.
- De Lorge JO. Operant behavior and colonic temperature of *Macaca mulatta* exposed to radio frequency fields at and above resonant frequencies. *Bioelectromagnetics.* 1984;5:233–46.
- Lotz WG. Hyperthermia in radiofrequency-exposed rhesus monkeys: a comparison of frequency and orientation effects. *Radiat Res.* 1985;102:59–70.
- Stuchly MA. Potentially hazardous microwave radiation source—a review. *J Microw Power.* 1977;12(4):369–81.
- Adair RK. Biophysical limits on athermal effects of RF and microwave radiation. *Bioelectromagnetics.* 2003;24:39–48.
- Prohofsky EW. RF absorption involving biological macromolecules. *Bioelectromagnetics.* 2004;25:441–51.
- International Commission on Non-Ionizing Radiation Protection (ICNIRP). In: Vecchia P, Matthes R, Ziegelberger G, Lin J, Saunders R, Swerdlow, editors. Exposure to high frequency electromagnetic fields, biological effects and health consequences (100 kHz–300 GHz); 2009. <https://www.icnirp.org/en/publications/article/hf-review-2009.html>.
- Food and Drug Administration (FDA). 1999. FDA's nomination of RF radiation in 1999 for the NTP study. Available at https://ntp.niehs.nih.gov/ntp/htdocs/chem_background/exsumpdf/wireless051999_508.pdf
- National Toxicology Program (NTP). NTP technical report on the toxicology and carcinogenesis studies in Hsd:Sprague Dawley SD rats exposed to whole-body radio frequency radiation at a frequency (900 MHz) and modulations (GSM and CDMA) used by cell phones, Technical report series no. 595. Research Triangle Park: National Institutes of Health, Public Health Service, U.S. Department of Health and Human Services; 2018. https://ntp.niehs.nih.gov/ntp/htdocs/lt_rpts/tr595_508.pdf?utm_source=direct&utm_medium=prod&utm_campaign=ntpgolinks&utm_term=tr595
- National Toxicology Program (NTP). NTP technical report on the toxicology and carcinogenesis studies in B6C3F1/N mice exposed to whole-body radio frequency radiation at a frequency (1,900 MHz) and modulations (GSM and CDMA) used by cell phones, Technical report series no. 596. Research Triangle Park: National Institutes of Health, Public Health Service, U.S. Department of Health and Human Services; 2018. https://ntp.niehs.nih.gov/ntp/htdocs/lt_rpts/tr596_508.pdf?utm_source=direct&utm_medium=prod&utm_campaign=ntpgolinks&utm_term=tr596
- Chou CK, Guy AW, Kunz LL, Johnson RB, Crowley JJ, Krupp JH. Long-term, low-level microwave irradiation of rats. *Bioelectromagnetics.* 1992;13:469–96.
- National Toxicology Program (NTP). National Toxicology Program peer review of the draft NTP technical reports on cell phone radiofrequency radiation. Research Triangle Park: National Institute of Environmental Health Sciences; 2018. Available at https://ntp.niehs.nih.gov/ntp/about_ntp/trpanel/2018/march/peerreview20180328_508.pdf
- Falcioni L, Bua L, Tibaldi E, Lauriola M, DeAngelis L, Gnudi F, et al. Report of final results regarding brain and heart tumors in Sprague-Dawley rats exposed from prenatal life until natural death to mobile phone radio-frequency field representative of a 1.8 GHz base station environmental emission. *Environ Res.* 2018;165:496–503.
- International Commission on Non-Ionizing Radiation Protection (ICNIRP). Principles for non-ionizing radiation protection. *Health Phys.* 2020;118:477–82.
- International Commission on Non-Ionizing Radiation Protection (ICNIRP). ICNIRP note: critical evaluation of two radiofrequency electromagnetic field animal carcinogenicity studies published in 2018. *Health Phys.* 2020;118:525–32.
- Melnick R. Regarding ICNIRP's evaluation of the National Toxicology Program's carcinogenicity studies of radiofrequency electromagnetic fields. *Health Phys.* 2020;118:678–82.
- Wyde M, Horn R, Capstick MH, Ladbury JM, Koepke G, Wilson PF, et al. Effect of cell phone radiofrequency radiation on body temperature in rodents: pilot studies of the National Toxicology Program's reverberation chamber exposure system. *Bioelectromagnetics.* 2018;39:190–9.
- Fragopoulou AF, Miltiadous P, Stamatakis A, Stylianopoulou F, Kousoulakos SL, Margaritis LH. Whole body exposure with GSM 900-MHz affects spatial memory in mice. *Pathophysiology.* 2010;17:179–87.
- Li Y, Shi C, Lu G, Xu Q, Liu S. Effects of electromagnetic radiation on spatial memory and synapses in rat hippocampal CA1. *Neural Regen Res.* 2012;7:1248–55.
- Narayanan SN, Kumar RS, Karun KM, Nayak SB, Bhat PG. Possible cause for altered spatial cognition of prepubescent rats exposed to chronic radiofrequency electromagnetic radiation. *Metab Brain Dis.* 2015;30:1193–206.

30. Razavinasab M, Moazzami K, Shabani M. Maternal mobile phone exposure alters intrinsic electrophysiological properties of CA1 pyramidal neurons in rat offspring. *Toxicol Ind Health*. 2016;32:968–79.
31. Schneider J, Stangassinger M. Nonthermal effects of lifelong high-frequency electromagnetic field exposure on social memory performance in rats. *Behav Neurosci*. 2014;128:633–7.
32. Tang J, Zhang Y, Yang L, Chen Q, Tan L, Zuo S, et al. Exposure to 900 MHz electromagnetic fields activates the mep-1/ERK pathway and causes blood-brain barrier damage and cognitive impairment in rats. *Brain Res*. 2015;1601:92–101.
33. Lai H. A summary of recent literature (2007–2017) on neurobiological effects of radiofrequency radiation. In: Markov M, editor. *Mobile communications and public health*. Boca Raton: CRC press; 2018. p. 187–222. <https://www.taylorfrancis.com/chapters/edit/10.1201/b22486-8/summary-recent-literature-2007-2017-neurobiological-effects-radio-frequency-radiation-henry-lai>.
34. Hardell L, Söderqvist F, Carlberg M, Zetterberg H, Hansson-Mild K. Exposure to wireless phone emissions and serum beta-trace protein. *Int J Mol Med*. 2010;26:301–6.
35. Frey AH, Feld SR, Frey B. Neural function and behavior: defining the relationship. *Ann NY Acad Sci*. 1975;247:433–9.
36. Persson BR, Salford LG, Brun A, Eberhardt JL, Malmgren L. Increased permeability of the blood-brain barrier induced by magnetic and electromagnetic fields. *Ann NY Acad Sci*. 1992;649:356–8.
37. Salford LG, Brun A, Stureson K, Eberhardt JL, Persson BR. Permeability of the blood-brain barrier induced by 915 MHz electromagnetic radiation, continuous wave and modulated at 8, 16, 50, and 200 Hz. *Microsc Res Tech*. 1994;15:535–42.
38. Eberhardt JL, Persson BR, Brun AE, Salford LG, Malmgren LO. Blood-brain barrier permeability and nerve cell damage in rat brain 14 and 28 days after exposure to microwaves from GSM mobile phones. *Electromagn Biol Med*. 2008;27:215–29.
39. Nittby H, Brun A, Eberhardt J, Malmgren L, Persson BR, Salford LG. Increased blood-brain barrier permeability in mammalian brain 7 days after exposure to the radiation from a GSM- 900 mobile phone. *Pathophysiology*. 2009;16:103–12.
40. Sirav B, Seyhan N. Effects of radiofrequency radiation exposure on blood-brain barrier permeability in male and female rats. *Electromagn Biol Med*. 2011;30:253–60.
41. Sirav B, Seyhan N. Effects of GSM modulated radio-frequency electromagnetic radiation on permeability of blood-brain barrier in male & female rats. *J Chem Neuroanat*. 2016;75:123–7.
42. Schuermann D, Mevissen M. Manmade electromagnetic fields and oxidative stress – biological effects and consequences for health. *Int J Mol Sci*. 2021;22:3772. <https://doi.org/10.3390/ijms22073772>.
43. Belyaev IY. 2010. Dependence of non-thermal biological effects of microwaves on physical and biological variables: implications for reproducibility and safety standards. *Eur J Oncol – Library*. 2010;5:187–218.
44. International Agency for Research on Cancer (IARC). IARC monograph on the evaluation of carcinogenic risks to humans: non-ionizing radiation, part 2: radiofrequency electromagnetic fields. Lyon, France, 102; 2013. p. 1–460. <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Non-ionizing-Radiation-Part-2-Radiofrequency-Electromagnetic-Fields-2013>
45. Prausnitz S, Susskind C. Effects of chronic microwave irradiation on mice. *Ire Trans Biomed Electron*. 1962;9:104–8.
46. La Vignera S, Condorelli RA, Vicari E, D'Agata R, Calogero AE. Effects of the exposure to mobile phones on male reproduction: a review of the literature. *J Androl*. 2012;33:350–6.
47. Kesari KK, Kumar S, Nirala J, Siddiqui MH, Behari J. Biophysical evaluation of radiofrequency electromagnetic field effects on male reproductive pattern. *Cell Biochem Biophys*. 2013;65:85–96.
48. Kesari KK, Agarwal A, Henkel R. Radiations and male fertility. *Reprod Biol Endocrinol*. 2018;16:118. <https://doi.org/10.1186/s12958-018-0431-1>.
49. Zha XD, Wang WW, Xu S, Shang XJ. Impacts of electromagnetic radiation from cellphones and Wi-fi on spermatogenesis. *Zhonghua Nan Ke Xue*. 2019;25:451–45.
50. Yadav H, Rai U, Singh R. Radiofrequency radiation: a possible threat to male fertility. *Reprod Toxicol*. 2021;100:90–100.
51. Agarwal A, Desai NR, Makker K, Varghese A, Mouradi R, Sabanegh E, et al. Effects of radiofrequency electromagnetic waves (RF-EMW) from cellular phones on human ejaculated semen: an in vitro pilot study. *Fertil Steril*. 2009;92:1318–25.
52. Adams JA, Galloway TS, Mondal D, Esteves SC, Mathews F. Effect of mobile telephones on sperm quality: a systematic review and meta-analysis. *Environ Int*. 2014;70:106–12.
53. Dama MS, Bhat MN. Mobile phones affect multiple sperm quality traits: a meta-analysis. *F100Res*. 2013;2:40. <https://doi.org/10.12688/f1000research.2-40.v1>.
54. Kim S, Han D, Ryu J, Kim K, Kim YH. Effects of mobile phone usage on sperm quality - no time-dependent relationship on usage: a systematic review and updated meta-analysis. *Environ Res*. 2021;202:111784. <https://doi.org/10.1016/j.envres.2021.111784>.
55. Yu G, Bai Z, Song C, Cheng Q, Wang G, Tang Z, et al. Current progress on the effect of mobile phone radiation on sperm quality: an updated systematic review and meta-analysis of human and animal studies. *Environ Pollut*. 2021;282:116592. <https://doi.org/10.1016/j.envpol.2021.116592>.
56. Zilberlicht A, Wiener-Megnazi Z, Sheinfeld Y, Grach B, et al. Habits of cell phone usage and sperm quality - does it warrant attention? *Reprod BioMed Online*. 2015;31:421–6.
57. Zalata A, El-Samanoudy AZ, Shaalan D, El-Baiomy Y, Mostafa T. In vitro effect of cell phone radiation on motility, DNA fragmentation and clusterin gene expression in human sperm. *Int J Fertil Steril*. 2015;9:129–36.
58. De Iuliis GN, Newey RJ, King BV, Aitken RJ. Mobile phone radiation induces reactive oxygen species production and DNA damage in human spermatozoa in vitro. *PLoS One*. 2009;4:e6446. <https://doi.org/10.1371/journal.pone.0006446>.
59. Kesari K, Kumar S, Behari J. Mobile phone usage and male infertility in Wistar rats. *Indian J Exp Biol*. 2010;48:987–92.
60. Alkis ME, Akdag MZ, Dasdag S, Yegin K, Akpolat V. Single-strand DNA breaks and oxidative changes in rat testes exposed to radiofrequency radiation emitted from cellular phones. *Biotechnol Biotechnol Equip*. 2019;33:1733–40.
61. Gautam R, Singh KV, Nirala J, Murmu NN, et al. Oxidative stress-mediated alterations on sperm parameters in male Wistar rats exposed to 3G mobile phone radiation. *Andrologia*. 2019;51:e13201. <https://doi.org/10.1111/and.13201>.
62. Yu G, Tang Z, Chen H, Chen Z, Wang L, Cao H, et al. Long-term exposure to 4G smartphone radiofrequency electromagnetic radiation diminished male reproductive potential by directly disrupting Spock3-MMP2-BTB axis in the testes of adult rats. *Sci Total Environ*. 2020;698:133860. <https://doi.org/10.1016/j.scitotenv.2019.133860>.
63. Andrášková S, Holovská K, Ševčíková Z, Andrejčáková Z, et al. The potential adverse effect of 2.45 GHz microwave radiation on the testes of prenatally exposed peripubertal male rats. *Histol Histopathol*. 2021;18402. <https://doi.org/10.14670/HH-18-402>.
64. Houston BJ, Nixon B, McEwan KE, Martin JH, King BV, Aitken RJ, et al. Whole-body exposures to radiofrequency-electromagnetic energy can cause DNA damage in mouse spermatozoa via an oxidative mechanism. *Sci Rep*. 2019;9:17478. <https://doi.org/10.1038/s41598-019-53983-9>.
65. Houston BJ, Nixon B, King B, Aitken RJ, De Iuliis GN. Probing the origins of 1,800 MHz radio frequency electromagnetic radiation induced damage in mouse immortalized germ cells and spermatozoa *in vitro*. *Front Public Health*. 2018;6:270. <https://doi.org/10.3389/fpubh.2018.00270>.
66. Kesari KK, Behari J. Evidence for mobile phone radiation exposure effects on reproductive pattern of male rats: role of ROS. *Electromagn Biol Med*. 2012;31:213–22.
67. Kumar S, Behari J, Sisodia R. Influence of electromagnetic fields on reproductive system of male rats. *Int J Radiat Biol*. 2013;89:147–54.
68. Pandey N, Giri S, Das S, Upadhaya P. Radiofrequency radiation (900 MHz)-induced DNA damage and cell cycle arrest in testicular germ cells in Swiss albino mice. *Toxicol Ind Health*. 2017;33:373–84.
69. Smith-Roe SL, Wyde ME, Stout MD, Winters JW, et al. Evaluation of the genotoxicity of cell phone radiofrequency radiation in male and female rats and mice following subchronic exposure. *Environ Mol Mutagen*. 2020;61:276–90.

70. Akdag M, Dasdag S, Canturk F, Akdag MZ. Exposure to non-ionizing electromagnetic fields emitted from mobile phones induced DNA damage in human ear canal hair follicle cells. *Electromagn Biol Med*. 2018;37:66–75.
71. Lai H. Genetic effects of non-ionizing electromagnetic fields. *Electromagn Biol Med*. 2021;40:264–73.
72. Yakymenko I, Tsybulin O, Sidorik E, Henshel D, et al. Oxidative mechanisms of biological activity of low-intensity radiofrequency radiation. *Electromagn Biol Med*. 2016;35:186–202.
73. Barnes FS, Greenebaum B. The effects of weak magnetic fields on radical pairs. *Bioelectromagnetics*. 2015;36:45–54.
74. Panagopoulos DJ, Karabarbounis A, Margaritis LH. Mechanism for action of electromagnetic fields on cells. *Biochem Biophys Res Commun*. 2002;298:95–102.
75. Belyaev I. Biophysical mechanisms for nonthermal microwave effects. In: Markov MS, editor. *Electromagnetic fields in biology and medicine*. Boca Raton, London, New York: CRC Press; 2015. p. 49–68. <https://www.taylorfrancis.com/chapters/mono/10.1201/b18148-9/biophysical-mechanisms-nonthermal-microwave-effects-marko-markov>.
76. Friedman J, Kraus S, Hauptman Y, Schiff Y, Seger R. Mechanism of short-term ERK activation by electromagnetic fields at mobile phone frequencies. *Biochem J*. 2007;405:559–68.
77. Inoue M, Sato EF, Nishikawa N, Park A-M, et al. Mitochondrial generation of reactive oxygen species and its role in aerobic life. *Curr Med Chem*. 2003;10:2495–505.
78. Yakymenko I, Burlaka A, Tsybulin I, Brieieva I, et al. Oxidative and mutagenic effects of low intensity GSM 1800 MHz microwave radiation. *Exp Oncol*. 2018;40:282–7.
79. Burlaka A, Tsybulin O, Sidorik E, Lukin S, et al. Overproduction of free radical species in embryonic cells exposed to low intensity radiofrequency radiation. *Exp Oncol*. 2013;35:219–25.
80. Alkis ME, Bilgin HM, Akpolat V, Dasdag S, et al. Effect of 900-, 1800-, and 2100-MHz radiofrequency radiation on DNA and oxidative stress in brain. *Electromagn Bio Med*. 2019;38:32–47.
81. Ding S-S, Sun P, Zhang Z, Liu X, et al. Moderate dose of trolox preventing the deleterious effects of Wi-fi radiation on spermatozoa in vitro through reduction of oxidative stress damage. *Chin Med J*. 2018;131:402–12.
82. Khalil AM, Gagaa MH, Alshamali AM. 8-Oxo-7, 8-dihydro-2'-deoxyguanosine as a biomarker of DNA damage by mobile phone radiation. *Hum Exp Toxicol*. 2012;31:734–40.
83. Xu S, Zhou Z, Zhang L, Yu Z, et al. Exposure to 1800 MHz radiofrequency radiation induces oxidative damage to mitochondrial DNA in primary cultured neurons. *Brain Res*. 2010;1311:189–96.
84. Güler G, Tomruk A, Ozjür E, Sahin D, et al. The effect of radiofrequency radiation on DNA and lipid damage in female and male infant rabbits. *Int J Radiat Biol*. 2012;88:367–73.
85. Bektas H, Dasdag S, Bektas MS. Comparison of effects of 2.4 GHz Wi-fi and mobile phone exposure on human placenta and cord blood. *Biotechnol Biotechnol Equip*. 2020;34:154–62.
86. Halliwell B. Biochemistry of oxidative stress. *Biochem Soc Trans*. 2007;35:1147–50.
87. International Agency for Research on Cancer (IARC). IARC monograph, a review of human carcinogens: radiation. Lyon, France, volume 100D; 2012. p. 1–363. <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Radiation-2012>
88. Smith MT, Guyton KZ, Gibbons CF, Fritz JM, Portier CJ, Rusyn I, et al. Key characteristics of carcinogens as a basis for organizing data on mechanisms of carcinogenesis. *Environ Health Perspect*. 2016;124:713–21.
89. D'Andrea JA, Gandhi OP, Lords JL. Behavioral and thermal effects of microwave radiation at resonant and nonresonant wavelengths. *Radio Sci*. 1977;12:251–6.
90. D'Andrea JA, Thomas A, Hatcher DJ. Rhesus monkey behavior during exposure to high-peak-power 5.62-GHz microwave pulses. *Bioelectromagnetics*. 1994;15:163–72.
91. D'Andrea JA, Gandhi OP, Lords JL, Durney CH, Johnson CC, Astle L. Physiological and behavioral effects of chronic exposure to 2450-MHz microwaves. *J Microw Power*. 1979;14:351–62.
92. D'Andrea JA, DeWitt JR, Emmerson RY, Bailey C, Gandhi OP. Intermittent exposure of rats to 2450 MHz microwaves at 2.5 mW/cm²: behavioral and physiological effects. *Bioelectromagnetics*. 1986;7:315–28.
93. Belyaev I. Duration of exposure and dose in assessing nonthermal biological effects of microwaves. In: Markov M, editor. *Dosimetry in bioelectromagnetics*. Boca Raton, London, New York: CRC Press; 2017. p. 171–84. <https://www.taylorfrancis.com/chapters/edit/10.1201/9781315154572-9/duration-exposure-dose-assessing-nonthermal-biological-effects-microwaves-igor-belyaev>.
94. Belyaev IY, Alipov YD, Shcheglov VS, Polunin VA, Aizenberg OA. Cooperative response of *Escherichia coli* cells to the resonance effect of millimeter waves at super low intensity. *Electro- Magnetobiol*. 1994;13:53–66.
95. Tillmann T, Ernst H, Streckert J, Zhou Y, Taugner F, Hansen V, et al. Indication of cocarcinogenic potential of chronic UMTS-modulated radiofrequency exposure in an ethylnitrosourea mouse model. *Int J Radiat Biol*. 2010;86:529–41.
96. Lerchl A, Klose M, Grote K, Wilhelm AF, Spathmann O, Fiedler T, et al. Tumor promotion by exposure to radiofrequency electromagnetic fields below exposure limits for humans. *Biochem Biophys Res Commun*. 2015;459:585–90.
97. Baohong W, Jiliang H, Lifan J, et al. Studying the synergistic damage effects induced by 1.8 GHz radiofrequency field radiation (RFR) with four chemical mutagens on human lymphocyte DNA using comet assay in vitro. *Mutat Res*. 2005;578:149–57.
98. Baohong W, Lifan J, Lanjuan L, et al. Evaluating the combinative effects on human lymphocyte DNA damage induced by ultraviolet ray C plus 1.8 GHz microwaves using comet assay in vitro. *Toxicol*. 2007;232:311–6.
99. Zhang MB, He JL, Jin LF, et al. Study of low-intensity 2450-MHz microwave exposure enhancing the genotoxic effects of mitomycin C using micronucleus test and comet assay in vitro. *Biomed Environ Sci*. 2002;15:283–90.
100. Kim JY, Hong SY, Lee YM, et al. In vitro assessment of clastogenicity of mobile-phone radiation (835 MHz) using the alkaline comet assay and chromosomal aberration test. *Environ Toxicol*. 2008;23:319–27.
101. Lameth J, Arnaud-Cormos D, Lévêque P, et al. Effects of a single head exposure to GSM-1800 MHz signals on the transcriptome profile in the rat cerebral cortex: enhanced gene responses under proinflammatory conditions. *Neurotox Res*. 2020;38:105–23.
102. López-Martin E, Bregains J, Relova-Quinteiro JL, et al. The action of pulse-modulated GSM radiation increases regional changes in brain activity and c-Fos expression in cortical and subcortical areas in a rat model of picrotoxin-induced seizure proneness. *J Neurosci Res*. 2009;87:1484–99.
103. Carballo-Quintás M, Martínez-Silva I, Cardarso-Suárez C, et al. A study of neurotoxic biomarkers, c-fos and GFAP after acute exposure to GSM radiation at 900 MHz in the picrotoxin model of rat brains. *Neurotoxicology*. 2011;32:478–94.
104. Kostoff RN, Heroux P, Aschner M, Tsatsakis A. Adverse health effects of 5G mobile networking technology under real-life conditions. *Toxicol Lett*. 2020;323:35–40.
105. Neufeld E, Kuster N. Systematic derivation of safety limits for time-varying 5G radiofrequency exposure based on analytical models and thermal dose. *Health Phys*. 2018;115:705–11.
106. Panagopoulos DJ, Karabarbounis A, Yakymenko I, Chrousos GP. Human-made electromagnetic fields: ion forced-oscillation and voltage-gated ion channel dysfunction, oxidative stress and DNA damage (review). *Int J Oncol*. 2021;59(92). <https://doi.org/10.3892/ijo.2021.5272>.
107. Pakhomov AG, Murphy MB. Comprehensive review of the research on biological effects of pulsed radiofrequency radiation in Russia and the former Soviet Union. In: Lin JC, editor. *Advances in electromagnetic fields in living system*, vol. 3. New York: Kluwer Academic/Plenum Publishers; 2000. p. 265–90. https://link.springer.com/chapter/10.1007/978-1-4615-4203-2_7.
108. Blackman CF. Cell phone radiation: evidence from ELF and RF studies supporting more inclusive risk identification and assessment. *Pathophysiology*. 2009;16:205–16.
109. Food and Drug Administration (FDA). Review of published literature between 2008 and 2018 of relevance to radiofrequency radiation and

- cancer; 2020. Available at <https://www.fda.gov/media/135043/download>
110. Zada G, Bond AE, Wang Y-P, Giannotta SL, Deapne D. Incidence trends in the anatomic location of primary malignant brain tumors in the United States: 1992–2006. *World Neurosurg*. 2012;77:518–24.
 111. Philips A, Henshaw DL, Lamburn G, O'Carroll MJ. Brain Tumours: rise in Glioblastoma Multiforme incidence in England 1995–2015 suggests an adverse environmental or lifestyle factor. *J Environ Public Health*. 2018;7910754. <https://doi.org/10.1155/2018/7910754>.
 112. Hardell L, Carlberg M. Mobile phones, cordless phones and rates of brain tumors in different age groups in the Swedish National Inpatient Register and the Swedish cancer register during 1998–2015. *PLoS One*. 2017;12:e0185461. <https://doi.org/10.1371/journal.pone.0185461>.
 113. Johansen C, Boice J, McLaughlin J, Olsen J. Cellular telephones and cancer—a nationwide cohort study in Denmark. *J Natl Cancer Inst*. 2001;93:203–7.
 114. Söderqvist F, Carlberg M, Hardell L. Review of four publications on the Danish cohort study on mobile phone subscribers and risk of brain tumors. *Rev Environ Health*. 2012;27:51–8.
 115. Hardell L, Carlberg M, Söderqvist F, Hansson MK. Pooled analysis of case-control studies on acoustic neuroma diagnosed 1997–2003 and 2007–2009 and use of mobile and cordless phones. *Int J Oncol*. 2013;43:1036–44.
 116. Hardell L, Carlberg M. Mobile phone and cordless phone use and the risk for glioma – analysis of pooled case-control studies in Sweden, 1997–2003 and 2007–2009. *Pathophysiology*. 2015;22:1–13.
 117. Interphone Study Group. Brain tumour risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Int J Epidemiol*. 2010;39:675–94.
 118. Coureau G, Bouvier G, Lebaillly P, Fabbro-Peray P, Gruber A, Leffondre K, et al. Mobile phone use and brain tumours in the CERENAT case-control study. *Occup Environ Med*. 2014;71:514–22.
 119. Interphone Study Group. Acoustic neuroma risk in relation to mobile telephone use: results of the INTERPHONE international case-control study. *Cancer Epidemiol*. 2011;35:453–64.
 120. Hardell L, Carlberg M. Use of mobile and cordless phones and survival of patients with glioma. *Neuroepidemiology*. 2013;40:101–8.
 121. Akhavan-Sigari R, Baf MM, Ariabod V, Rohde V, Rahighi S. Connection between cell phone use, p53 gene expression in different zones of glioblastoma multiforme and survival prognoses. *Rare Tumors*. 2014;6:5350. <https://doi.org/10.4081/rt.2014.5350>.
 122. Moon IS, Kim BG, Kim J, Lee JD, Lee WS. Association between vestibular schwannomas and mobile phone use. *Tumour Biol*. 2014;35:581–7.
 123. Sato Y, Akiba S, Kubo O, Yamaguchi N. A case-case study of mobile phone use and acoustic neuroma risk in Japan. *Bioelectromagnetics*. 2011;32:85–93.
 124. Pettersson D, Mathiesen T, Prochazka M, Bergenheim T, Florentzon R, Harder H, et al. Long-term mobile phone use and acoustic neuroma risk. *Epidemiology*. 2014;25:233–41.
 125. Schoemaker MJ, Swerdlow AJ, Ahlborn A, Avinen A, Blaasaas KG, Cardis E, et al. Mobile phone use and risk of acoustic neuroma: results of the Interphone case-control study in five north European countries. *Br J Cancer*. 2005;93:842–8.
 126. Momoli F, Siemiatycki J, McBride ML, Parent ME, Richardson L, Bedard D, et al. Probabilistic multiple-bias modelling applied to the Canadian data from the INTERPHONE study of mobile phone use and risk of glioma, meningioma, acoustic neuroma, and parotid gland tumors. *Am J Epidemiol*. 2017;186:885–93.
 127. Luo J, Deziel NC, Huang H, Chen Y, Ni X, Ma S, et al. Cell phone use and risk of thyroid cancer: a population-based case-control study in Connecticut. *Ann Epidemiol*. 2019;29:39–45.
 128. Luo J, Li H, Deziel NC, Huang H, Zhao N, Ma S, et al. Genetic susceptibility may modify the association between cell phone use and thyroid cancer: a population-based case-control study in Connecticut. *Environ Res*. 2020;182:109013. <https://doi.org/10.1016/j.envres.2019.109013>.
 129. Carlberg M, Hedendahl L, Ahonen M, Koppel T, Hardell L. Increasing incidence of thyroid cancer in the Nordic countries with main focus on Swedish data. *BMC Cancer*. 2016;16:426. <https://doi.org/10.1186/s12885-016-2429-4>.
 130. Carlberg M, Koppel T, Hedendahl LK, Hardell L. Is the increasing incidence of thyroid cancer in the Nordic countries caused by use of mobile phones? *Int J Environ Res Public Health*. 2020;17(23):9129. <https://doi.org/10.3390/ijerph17239129>.
 131. Shih YW, Hung CS, Huang CC, Chou KR, Niu SF, et al. The association between smartphone use and breast cancer risk among Taiwanese women: a case-control study. *Cancer Manag Res*. 2020;12:10799–807. <https://doi.org/10.2147/CMAR.S267415>.
 132. Gandhi OP, Lazzi G, Furse CM. Electromagnetic absorption in the human head and neck for mobile telephones at 835 and 1900 MHz. *IEEE Trans Microw Theory Tech*. 1996;44:1884–97.
 133. Gandhi OP, Morgan L, de Salles AA, Han YY, Herberman RB, Davis DL. Exposure limits: the underestimation of absorbed cell phone radiation, especially in children. *Electromagn Biol Med*. 2012;31:34–51.
 134. Fernández-Rodríguez CE, de Salles AA, Davis DL. Dosimetric simulations of brain absorption of mobile phone radiation—the relationship between psSAR and age. *IEEE Access*. 2015;3:2425–30.
 135. Fernández-Rodríguez C, de Salles AA. On the sensitivity of the skull thickness for the SAR assessment in the intracranial tissues, 2016 IEEE MTT-S Latin America microwave conference (LAMC); 2016. <https://doi.org/10.1109/LAMC.2016.7851256>.
 136. Fernández C, de Salles AA, Sears ME, Morris RD, Davis DL. Absorption of wireless radiation in the child versus adult brain and eye from cell phone conversation or virtual reality. *Environ Res*. 2018;167:694–9. <https://doi.org/10.1016/j.envres.2018.05.013>.
 137. Christ A, Gosselin MC, Christopoulou M, Kühn S, Kuster N. Age-dependent tissue-specific exposure of cell phone users. *Phys Med Biol*. 2010;55:1767–83.
 138. Foster KR, Chou CK. Response to "children absorb higher doses of radio frequency electromagnetic radiation from mobile phones than adults" and "yes the children are more exposed to radiofrequency energy from mobile telephones than adults". *IEEE Access*. 2016;4:5322–6.
 139. de Salles AA, Bulla G, Fernández-Rodríguez CE. Electromagnetic absorption in the head of adults and children due to mobile phone operation close to the head. *Electromagn Biol Med*. 2006;25:349–60.
 140. Peyman A, Gabriel C, Gran EH, Vermeeren G, Martens L. Variation of the dielectric properties of tissues with age: the effect on the values of SAR in children when exposed to walkie-talkie devices. *Phys Med Biol*. 2009;2009(54):227–41.
 141. Blondin JP, Nguyen DH, Sbeghen J, Goulet D, et al. Human perception of electric fields and ion currents associated with high-voltage DC transmission lines. *Bioelectromagnetics*. 1996;17:230–41.
 142. Leitgeb N, Schroettner J. Electric current perception study challenges electric safety limits. *J Med Eng Technol*. 2002;26:168–72.
 143. Leitgeb N, Schroettner J, Cech RJ. Electric current perception of children: the role of age and gender. *Med. Eng Technol*. 2006;30:306–9.
 144. Leitgeb N, Schröttner J, Cech R. Perception of ELF electromagnetic fields: excitation thresholds and inter-individual variability. *Health Phys*. 2007;92:591–5.
 145. McCarty DE, Carrubba S, Chesson AL, Filot C, et al. Electromagnetic hypersensitivity: evidence for a novel neurological syndrome. *Int J Neurosci*. 2011;121:670–6.
 146. Hinrikus H, Parts M, Lass J, Tuulik V. Changes in human EEG caused by low level modulated microwave stimulation. *Bioelectromagnetics*. 2004;2004(25):431–40.
 147. Hinrikus H, Bachmann M, Lass J, et al. Effect of low frequency modulated microwave exposure on human EEG: individual sensitivity. *Bioelectromagnetics*. 2008;29:527–38.
 148. Mueller CH, Krueger H, Schierz C. Project NEMESIS: perception of a 50 Hz electric and magnetic field at low intensities (laboratory experiment). *Bioelectromagnetics*. 2002;23:26–36.
 149. Legros A, Beuter A. Individual subject sensitivity to extremely low frequency magnetic field. *Neurotoxicology*. 2006;27:534–46.
 150. Kimata H. Microwave radiation from cellular phones increases allergen-specific IgE production. *Allergy*. 2005;60:838–9.
 151. Rea WJ, Pan Y, Fenyves EJ, Sujisawa I, et al. Electromagnetic field sensitivity. *J Bioelectricity*. 1991;10:241–56.
 152. Belpomme D, Irigaray P. Electrohypersensitivity as a newly identified and characterized neurologic pathological disorder: how to

- diagnose, treat, and prevent it. *Int J Mol Sci.* 2020;21:1915. <https://doi.org/10.3390/ijms21061915>.
153. Stein Y, Udasin IG. Electromagnetic hypersensitivity (EHS, microwave syndrome) - review of mechanisms. *Environ Res.* 2020;186:109445. <https://doi.org/10.1016/j.envres.2020.109445>.
 154. Hagström M, Auranen J, Ekman R. Electromagnetic hypersensitive Finns: symptoms, perceived sources and treatments, a questionnaire study. *Pathophysiology.* 2013;20:117–22.
 155. Belyaev I, Dean A, Eger H, Hubmann G, et al. European EMF guideline 2016 for the prevention, diagnosis and treatment of EMF-related health problems and illness. *Rev Environ Health.* 2016;31:363–97.
 156. Austrian Medical Association. Guideline of the Austrian medical association for the diagnosis and treatment of EMF-related health problems and illnesses (EMF syndrome); 2012. Available at <https://vagbrytaren.org/Guideline%20%20AG-EMF.pdf>
 157. Hardell L, Koppel T. Electromagnetic hypersensitivity close to mobile phone base stations - a case study in Stockholm, Sweden. *Rev Environ Health.* 2022. <https://doi.org/10.1515/reveh-2021-0169>.
 158. Havas M. Radiation from wireless technology affects the blood, the heart, and the autonomic nervous system. *Rev Environ Health.* 2013;2013(28):75–84.
 159. Leitgeb N, Schröttner J. Electrosensitivity and electromagnetic hypersensitivity. *Bioelectromagnetics.* 2003;24:387–94.
 160. Deshmukh PS, Banerjee BD, Abegaonkar MP, Megha K, et al. Effect of low level microwave radiation exposure on cognitive function and oxidative stress in rats. *Indian J Biochem Biophys.* 2013;50:114–9.
 161. Everaert J, Bauwens D. A possible effect of electromagnetic radiation from mobile phone base stations on the number of breeding house sparrows (*Passer domesticus*). *Electromagn Biol Med.* 2007;26:63–72.
 162. Megha K, Deshmukh PS, Banerjee BD, et al. Microwave radiation induced oxidative stress, cognitive impairment and inflammation in brain of Fischer rats. *Indian J Exp Biol.* 2012;50:889–96.
 163. Narayanan SN, Kumar RS, Potu BK, Nayak S. Effect of radio-frequency electromagnetic radiations (RF-EMR) on passive avoidance behaviour and hippocampal morphology in Wistar rats. *Ups J Med Sci.* 2010;115:91–6.
 164. Narayanan SN, Kumar RS, Paval J, Kedage V, et al. Analysis of emotionality and locomotion in radio-frequency electromagnetic radiation exposed rats. *Neurol Sci.* 2013;34:1117–24.
 165. Narayanan SN, Kumar RS, Kedage V, Nalini K, et al. Evaluation of oxidant stress and antioxidant defense in discrete brain regions of rats exposed to 900 MHz radiation. *Bratisl Lek Listy.* 2014;115:260–6.
 166. Cammaerts MC, De Doncker P, Patris X, Bellens F, Rachidi Z, Cammaerts D. GSM 900 MHz radiation inhibits ants' association between food sites and encountered cues. *Electromagn Biol Med.* 2012;31:151–65.
 167. Balmori A, Hallberg O. The urban decline of the house sparrow (*Passer domesticus*): a possible link with electromagnetic radiation. *Electromagn Biol Med.* 2007;26:141–51.
 168. Balmori A. Mobile phone mass effects on common frog (*Rana temporaria*) tadpoles: the city turned into a laboratory. *Electromagn Biol Med.* 2010;29:31–5.
 169. Aldad TS, Gan G, Gao XB, Taylor HS. Fetal radiofrequency radiation exposure from 800-1900 MHz-rated cellular telephones affects neurodevelopment and behavior in mice. *Sci Rep.* 2012;2:312. <https://doi.org/10.1038/srep00312>.
 170. Nittby H, Grafström G, Tian DP, Malmgren L, et al. Cognitive impairment in rats after long-term exposure to GSM-900 mobile phone radiation. *Bioelectromagnetics.* 2008;29:219–32.
 171. Ntzouni MP, Stamatakis A, Stylianopoulou F, Margaritis LH. Short-term memory in mice is affected by mobile phone radiation. *Pathophysiology.* 2011;18:193–9.
 172. Saikhedkar N, Bhatnagar M, Jain A, Sukhwai P, et al. Effects of mobile phone radiation (900 MHz radiofrequency) on structure and functions of rat brain. *Neurol Res.* 2014;36:1072–9.
 173. Rubin GJ, Nieto-Hernandez R, Wessely S. Idiopathic environmental intolerance attributed to electromagnetic fields (formerly 'electromagnetic hypersensitivity'): an updated systematic review of provocation studies. *Bioelectromagnetics.* 2010;31:1–11.
 174. Markova E, Hillert L, Malmgren L, Persson BRR, Belyaev IY. Microwaves from GSM mobile telephones affect 53BP1 and gamma-H2AX foci in human lymphocytes from hypersensitive and healthy persons. *Environ Health Perspect.* 2005;113:1172–7.
 175. Markova E, Malmgren LO, Belyaev IY. Microwaves from mobile phones inhibit 53BP1 focus formation in human stem cells more strongly than in differentiated cells: possible mechanistic link to cancer risk. *Environ Health Perspect.* 2010;118:394–9.
 176. Belyaev IY, Markova E, Hillert L, Malmgren LOG, Persson BRR. Microwaves from UMTS/GSM mobile phones induce long-lasting inhibition of 53BP1/gamma-H2AX DNA repair foci in human lymphocytes. *Bioelectromagnetics.* 2009;2009(30):129–41.
 177. Gulati S, Kosik P, Durdik M, Skorvaga M, et al. Effects of different mobile phone UMTS signals on DNA, apoptosis and oxidative stress in human lymphocytes. *Environ Pollut.* 2020;267:115632. <https://doi.org/10.1016/j.envpol.2020.115632>.
 178. Dieudonné M. Does electromagnetic hypersensitivity originate from nocebo responses? Indications from a qualitative study. *Bioelectromagnetics.* 2016;37:14–24.
 179. International Commission on Non-Ionizing Radiation Protection (ICNIRP). General approach to protection against non-ionizing radiation. *Health Phys.* 2002;82:540–8.
 180. World Health Organization (WHO). Electromagnetic fields and public health. Electromagnetic hypersensitivity; 2005. <https://web.archive.org/web/20220423095028/https://www.who.int/teams/environment-climate-change-and-health/radiation-and-health/non-ionizing/el-sensitivitiy>
 181. Havas M. Electrohypersensitivity (EHS) is an environmentally-induced disability that requires immediate attention. *J Sci Discov.* 2019;3(1):jsd18020. <https://doi.org/10.24262/jsd.3.1.18020>.
 182. US Environmental Protection Agency (US EPA). A review of the reference dose (RfD) and reference concentration (RfC) process. Risk assessment forum. EPA/630/P-02/002F. Washington, DC; 2002. Available at: <https://www.epa.gov/sites/default/files/2014-12/documents/rfd-final.pdf>
 183. International Council for Harmonization (ICH). Impurities: guidelines for residual solvents Q3C(R7); 2018. Available at: <https://www.pmda.go.jp/files/000231003.pdf>
 184. Dankovic DA, Naumann BD, Maier A, Dourson ML, Levy LS. The scientific basis of uncertainty factors used in setting occupational exposure limits. *J Occup Environ Hyg.* 2015;12:555–68.
 185. Uche UI, Naidenko OV. Development of health-based exposure limits for radiofrequency radiation from wireless devices using a benchmark dose approach. *Environ Health.* 2021;20:84. <https://doi.org/10.1186/s12940-021-00768-1>.
 186. Peleg M, Naativ O, Richter ED. Radio frequency radiation-related cancer: assessing causation in the occupational/military setting. *Environ Res.* 2018;163:123–33.
 187. Gong Y, Capstick M, McCormick DL, Gauger JR, Horn T, Wilson P, et al. Life time dosimetric assessment for mice and rats exposed to cell phone radiation. *IEEE Trans Electromagn Compat.* 2017;59:1798–808.
 188. Alvarez-Buylla A, Lim DA. For the long run: maintaining germinal niches in the adult brain. *Neuron.* 2004;41:683–6.
 189. Levitt BB, Lai HC, Manville AM. Effects of non-ionizing electromagnetic fields on flora and fauna, part 1. Rising ambient EMF levels in the environment. *Rev Environ Health.* 2021. <https://doi.org/10.1515/reveh-2021-0026>.
 190. Levitt BB, Lai HC, Manville AM. Effects of non-ionizing electromagnetic fields on flora and fauna, part 2 impacts: how species interact with natural and man-made EMF. *Rev Environ Health.* 2021. <https://doi.org/10.1515/reveh-2021-0050>.
 191. Moller A, Sagasser S, Wiltshko W, Schierwater B. Retinal cryptochrome in a migratory passerine bird: a possible transducer for the avian magnetic compass. *Naturwissenschaften.* 2004;91:585–8.
 192. Heyers D, Manns M, Luksch H, Güntürkün O, Mouritsen H. A visual pathway links brain structures active during magnetic compass orientation in migratory birds. *PLoS One.* 2007;2:e937. <https://doi.org/10.1371/journal.pone.0000937>.
 193. Collett TS, Barron J. Biological compasses and the coordinate frame of landmark memories in honeybees. *Nature.* 1994;386:137–40.
 194. Holland RA, Kirschvink JL, Doak TG, Wikelski M. Bats use magnetoreception to detect the earth's magnetic field. *PLoS One.* 2008;3:e1676. <https://doi.org/10.1371/journal.pone.0001676>.

195. Putman NF, Scanlan MM, Billman EJ, O'Neil JP, Couture RB, Quinn TP, et al. An inherited magnetic map guides ocean navigation in juvenile Pacific salmon. *Curr Biol*. 2014;24:446–50.
196. Putman NF, Williams CR, Gallagher EP, Dittman AH. A sense of place: pink salmon use a magnetic map for orientation. *J Exp Biol*. 2020;223:218735. <https://doi.org/10.1242/jeb.218735>.
197. Quinn TP, Merrill RT, Brannon EL. Magnetic field detection in sockeye salmon. *J Exp Zool*. 1981;217:137–42.
198. Kalmijn AJ. Electric and magnetic field detection in elasmobranch fishes. *Science*. 1982;198(218):916–8.
199. Engels S, Schneider NL, Lefeldt N, Hein CM, Zapka M, Michalik A, et al. Anthropogenic electromagnetic noise disrupts magnetic compass orientation in a migratory bird. *Nature*. 2014;509:353–6.
200. Pakhomov A, Bojarinova J, Cherbunin R, Chetverikova R, Grigoryev PS, Kavokin K, et al. Very weak oscillating magnetic field disrupts the magnetic compass of songbird migrants. *J R Soc Interface*. 2017;14:20170364. <https://doi.org/10.1098/rsif.2017.0364>.
201. Schwarze S, Schneibder NL, Reichl T, Dreyer D, Lefeldt N, Engels S, et al. Weak broadband electromagnetic fields are more disruptive to magnetic compass orientation in a night-migratory songbird (*Erethacus rubecula*) than strong narrow-band fields. *Front Behav Neurosci*. 2016;10:55. <https://doi.org/10.3389/fnbeh.2016.00055>.
202. Wiltchko R, Thalau P, Gehring D, Nießner C, Ritz T, Wiltchko W. Magnetoreception in birds: the effect of radio-frequency fields. *J R Soc Interface*. 2015;12:20141103. <https://doi.org/10.1098/rsif.2014.1103>.
203. Landler L, Painter MS, Youmans PW, Hopkins WA, Phillips JB. Spontaneous magnetic alignment by yearling snapping turtles: rapid association of radio frequency dependent pattern of magnetic input with novel surroundings. *PLoS One*. 2015;10:e0124728. <https://doi.org/10.1371/journal.pone.0124728>.
204. Putman NF, Meinke AM, Noakes DL. Rearing in a distorted magnetic field disrupts the 'map sense' of juvenile steelhead trout. *Biol Lett*. 2014;10:20140169. <https://doi.org/10.1098/rsbl.2014.0169>.
205. Sharma VP, Kumar NR. Changes in honeybee behaviour and biology under the influence of cellphone radiations. *Curr Sci*. 2010;98:1376–8.
206. Odemer R, Odemer F. Effects of radiofrequency electromagnetic radiation (RF-EMF) on honey bee queen development and mating success. *Sci Total Environ*. 2019;661:553–62.
207. Gabriel C, Lau RW, Gabriel S. The dielectric properties of biological tissues: II. Measurements in the frequency range 10 Hz to 20 GHz. *Phys Med Biol*. 1996;41:2251–69.
208. Gandhi O, Riaz A. Absorption of millimeter waves by human beings and its biological implications. *IEEE Trans Microw Theory Tech*. 1986;34:228–35.
209. Thielens A, Bell D, Mortimore DB, Greco MK, Martens L, Joseph W. Exposure of insects to radio-frequency electromagnetic fields from 2 to 120 GHz. *Sci Rep*. 2018;8(1):3924. <https://doi.org/10.1038/s41598-018-22271-3>.
210. Pretz K. Will 5G be bad for our health? *IEEE Spectr*. 2019; <https://pectrum.ieee.org/will-5g-be-bad-for-our-health>.
211. Neufeld E, Carrasco E, Murbach M, Balzano Q, Christ A, Kuster N. Theoretical and numerical assessment of maximally allowable power-density averaging area for conservative electromagnetic exposure assessment above 6 GHz. *Bioelectromagnetics*. 2018;39:617–30.
212. Foster KR, Ziskin MC, Balzano Q. Thermal response of human skin to microwave energy: a critical review. *Health Phys*. 2016;111:528–41.
213. Anderson RR, Parrish JA. The optics of human skin. *J Invest Dermatol*. 1981;77:13–9.
214. Meinhardt M, Kerbs R, Anders A, Heinrich U, Tronnier H. Wavelength-dependent penetration depths of ultraviolet radiation in human skin. *J Biomed Opt*. 2008;13:044030. <https://doi.org/10.1117/1.2957970>.
215. Pakhomov AG, Akyl Y, Pakhomova ON, Stuck BE, Murphy MR. Current state and implications of research on biological effects of millimeter waves: a review of the literature. *Bioelectromagnetics*. 1998;19:393–413.
216. Belyaev IY, Shcheglov VS, Alipov ED, Ushakov VD. Nonthermal effects of extremely high-frequency microwaves on chromatin conformation in cells in vitro - dependence on physical, physiological, and genetic factors. *IEEE Trans Microw Theory Tech*. 2000;48:2172–9.
217. Albanese R, Blaschak J, Medina R, Penn J. Ultrashort electromagnetic signals: biophysical questions, safety issues, and medical opportunities. *Aviat Space Environ Med*. 1994;65:A116–20.
218. Oughstun KE. Optimal pulse penetration in Lorentz-model dielectrics using the Sommerfeld and Brillouin precursors. *Opt Express*. 2015;23:26604–16.
219. Wood AW. What is the current status of research on mm-wave frequencies? - in relation to health; 2018. <https://slideplayer.com/slide/14592262/>
220. Blackman C, Forge S. 5G deployment: state of play in Europe, USA, and Asia. European Parliament; 2019. [http://www.europarl.europa.eu/RegData/etudes/IDAN/2019/631060/IPOL_IDA\(2019\)631060_EN.pdf](http://www.europarl.europa.eu/RegData/etudes/IDAN/2019/631060/IPOL_IDA(2019)631060_EN.pdf)
221. Regel SJ, Gottselig JM, Schuderer J, Tinguely G, et al. Pulsed radio frequency radiation affects cognitive performance and the waking electroencephalogram. *NeuroReport*. 2007;18:803–7.
222. Thomas JR, Schrot J, Banvard RA. Comparative effects of pulsed and continuous-wave 2.8-GHz microwaves on temporally defined behavior. *Bioelectromagnetics*. 1982;3:227–35.
223. Creighton MO, Larsen LE, Stewart-DeHaan PJ, Jacobi JH, et al. In vitro studies of microwave-induced cataract. II. Comparison of damage observed for continuous wave and pulsed microwaves. *Exp Eye Res*. 1987;45:357–73.
224. Czerska EM, Elson EC, Davis CC, Swicord ML, Czerski P. Effects of continuous and pulsed 2450-MHz radiation on spontaneous lymphoblastoid transformation of human lymphocytes in vitro. *Bioelectromagnetics*. 1992;13:247–59.
225. El Khoueiry C, Moretti D, Renom R, Camera F, Orlacchio R, Garenne A, et al. Decreased spontaneous electrical activity in neuronal networks exposed to radiofrequency 1,800 MHz signals. *J Neurophysiol*. 2018;120:2719–29.
226. Mohammed HS, Fahmy HM, Radwan NM, Elsayed AA. Non-thermal continuous and modulated electromagnetic radiation fields effects on sleep EEG of rats. *J Adv Res*. 2013;4:181–7.
227. Blank M, Havas M, Kelley E, Lai H, Moskowitz J. International appeal: scientists call for protection from non-ionizing electromagnetic field exposure. *Eur J Oncol Environ Health*. 2015;20:180–2 Available from: <https://mattioli1885journals.com/index.php/EJOEH/article/view/4971>.
228. International Agency for Research on Cancer (IARC). IARC monograph, a review of human carcinogens: arsenic, metals, fibres, and dusts. Lyon, France, volume 100C; 2012. p. 1–527. <https://publications.iarc.fr/Book-And-Report-Series/Iarc-Monographs-On-The-Identification-Of-Carcinogenic-Hazards-To-Humans/Arsenic-Metals-Fibres-And-Dusts-2012>
229. Institute of Electrical and Electronics Engineers. IEEE standard for safety levels with respect to human exposure to electric, magnetic, and electromagnetic fields, 0 Hz to 300 GHz. IEEE Std C95.1™. New York: IEEE; 2019. <https://ieeexplore.ieee.org/document/8859679>
230. Bandara P, Carpenter DO. Planetary electromagnetic pollution: it is time to assess its impact. *Lancet Planet Health*. 2018;2:e512–4. [https://doi.org/10.1016/S2542-5196\(18\)30221-3](https://doi.org/10.1016/S2542-5196(18)30221-3).

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

